



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶: **C12N 15/13, C07K 19/00, A61K 47/48, C07K 16/24, C12N 15/85, 5/10** **A1** (11) International Publication Number: **WO 99/37779**
(43) International Publication Date: **29 July 1999 (29.07.99)**

(21) International Application Number: **PCT/US99/01081**(22) International Filing Date: **19 January 1999 (19.01.99)**

(30) Priority Data:

09/012,116	22 January 1998 (22.01.98)	US
PCT/US98/03337	20 February 1998 (20.02.98)	US
09/122,513	24 July 1998 (24.07.98)	US
09/121,952	24 July 1998 (24.07.98)	US

(71) Applicant (for all designated States except US): **GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).**


(72) Inventors; and

(75) Inventors/Applicants (for US only): **HSEI, Vanessa [US/US]; 5047 Capistrano Avenue, San Jose, CA 95129 (US). KOUMENIS, Iphigenia [CY/US]; Apartment 6, 3820 Park Boulevard, Palo Alto, CA 94306 (US). LEONG, Steven, J. [US/US]; 1914 Eldorado Avenue, Berkeley, CA 94707 (US). PRESTA, Leonard, G. [US/US]; 1900 Gough Street #206, San Francisco, CA 94109 (US). SHAHROKH, Zahra [US/US]; 24 Sotelo Avenue, San Francisco, CA 94116 (US). ZAPATA, Gerardo, A. [US/US]; 785 Widgeon Street, Foster City, CA 94404 (US).**(74) Agents: **LOVE, Richard, B. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).**(81) Designated States: **AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: **ANTIBODY FRAGMENT-POLYMER CONJUGATES AND HUMANIZED ANTI-IL-8 MONOCLONAL ANTIBODIES AND USES OF SAME**

(57) Abstract

Humanized anti-IL-8 monoclonal antibodies and variants thereof are described for use in diagnostic applications and in the treatment of inflammatory disorders. Also described is a conjugate formed by an antibody fragment covalently attached to a non-proteinaceous polymer, wherein the apparent size of the conjugate is at least about 500 kD. The conjugate exhibits substantially improved half-life, mean residence time, and/or clearance rate in circulation as compared to the underivatized parental antibody fragment.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ANTIBODY FRAGMENT-POLYMER CONJUGATES AND HUMANIZED ANTI-IL-8 MONOCLONAL ANTIBODIES AND USES OF SAME

FIELD OF THE INVENTION

5 This application relates to the field of antibody fragments derivatized with polymers, and in particular to the use of such derivatization to increase the circulation half-lives of antibody fragment-polymer conjugates. This application also relates to the field of inflammatory diseases and asthma, and in particular to anti-IL-8 antibody treatment of inflammatory diseases and asthmatic diseases. This application further relates to humanized anti-interleukin-8 (IL-8) antibodies and to high affinity variants of such antibodies.

10 BACKGROUND

Modification of proteins with polyethylene glycol ("PEGylation") has the potential to increase residence time and reduce immunogenicity in vivo. For example, Knauf et al., J. Biol. Chem., 263: 15064-15070 (1988) reported a study of the pharmacodynamic behavior in rats of various polyoxylated glycerol and polyethylene glycol modified species of interleukin-2. Despite the known advantage of PEGylation, 15 PEGylated proteins have not been widely exploited for clinical applications. In the case of antibody fragments, PEGylation has not been shown to extend serum half-life to useful levels. Delgado et al., Br. J. Cancer, 73: 175-182 (1996), Kitamura et al., Cancer Res., 51: 4310-4315 (1991), Kitamura et al., Biochem. Biophys. Res. Comm., 171: 1387-1394 (1990), and Pedley et al., Br. J. Cancer, 70: 1126-1130 (1994) reported studies characterizing blood clearance and tissue uptake of certain anti-tumor antigen antibodies or antibody fragments 20 derivatized with low molecular weight (5 kD) PEG. Zapata et al., FASEB J., 9: A1479 (1995) reported that low molecular weight (5 or 10 kD) PEG attached to a sulfhydryl group in the hinge region of a Fab' fragment reduced clearance compared to the parental Fab' molecule.

Interleukin-8 (IL-8) is neutrophil chemotactic peptide secreted by a variety of cells in response to inflammatory mediators (for a review see Hebert et al. Cancer Investigation 11(6):743 (1993)). IL-8 can play 25 an important role in the pathogenesis of inflammatory disorders, such as adult respiratory distress syndrome (ARDS), septic shock, and multiple organ failure. Immune therapy for such inflammatory disorders can include treatment of an affected patient with anti-IL-8 antibodies.

Sticherling et al. (J. Immunol. 143:1628 (1989)) disclose the production and characterization of four monoclonal antibodies against IL-8. WO 92/04372, published March 19, 1992, discloses polyclonal antibodies 30 which react with the receptor-interacting site of IL-8 and peptide analogs of IL-8, along with the use of such antibodies to prevent an inflammatory response in patients. St. John et al. (Chest 103:932 (1993)) review immune therapy for ARDS, septic shock, and multiple organ failure, including the potential therapeutic use of anti-IL-8 antibodies. Sekido et al. (Nature 365:654 (1993)) disclose the prevention of lung reperfusion injury in rabbits by a monoclonal antibody against IL-8. Mulligan et al. (J. Immunol. 150:5585 (1993)), disclose 35 protective effects of a murine monoclonal antibody to human IL-8 in inflammatory lung injury in rats.

WO 95/23865 (International Application No. PCT/US95/02589 published September 8, 1995) demonstrates that anti-IL-8 monoclonal antibodies can be used therapeutically in the treatment of other inflammatory disorders, such as bacterial pneumonias and inflammatory bowel disease.

Anti-IL-8 antibodies are additionally useful as reagents for assaying IL-8. For example, Sticherling et al. (Arch. Dermatol. Res. 284:82 (1992)), disclose the use of anti-IL-8 monoclonal antibodies as reagents in 40

immun hist chemical studies. Ko *et al.* (*J. Immunol. Methods* 149:227 (1992)) disclose the use of anti-IL-8 monoclonal antibodies as reagents in an enzyme-linked immunoabsorbent assay (ELISA) for IL-8.

SUMMARY OF THE INVENTION

One aspect of the invention is a conjugate consisting essentially of one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein the apparent size of the conjugate is at least about 500 kD.

Another aspect of the invention is a conjugate formed by one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein the apparent size of the conjugate is at least about 500 kD, and wherein the covalent structure of the conjugate is free of any matter other than the antibody fragment and nonproteinaceous polymer molecules.

Yet another aspect of the invention is a conjugate formed by the one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein the covalent structure of the conjugate further incorporates one or more nonproteinaceous labels, wherein the covalent structure of the conjugate is free of any matter other than the antibody fragment, nonproteinaceous polymer and nonproteinaceous label molecules, and wherein the apparent size of the conjugate is at least about 500 kD.

An additional aspect of the invention is a method of treating an inflammatory disorder in a mammal comprising administering to the mammal an effective amount of a conjugate consisting essentially of one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein at least one antibody fragment comprises an antigen binding site that binds to human interleukin-8 (IL-8), and wherein the apparent size of the conjugate is at least about 500 kD.

A further aspect of the invention is a method of treating an asthmatic disorder in a mammal comprising administering to the mammal an effective amount of a conjugate consisting essentially of one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein at least one antibody fragment comprises an antigen binding site that binds to human interleukin-8 (IL-8), and wherein the apparent size of the conjugate is at least about 500 kD.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a graph depicting the blocking of IL-8 mediated elastase release from neutrophils by anti-IL-8 monoclonal antibody 5.12.14.

Figure 2 is a graph depicting the inhibition of I-IL-8 binding to neutrophils by unlabeled IL-8.

Figure 3 demonstrates that a isotype matched negative control Fab (denoted as "4D5 Fab") does not inhibit the binding of ¹²⁵I-IL-8 to human neutrophils.

Figure 4 is a graph depicting the inhibition of binding of ¹²⁵I-IL-8 to human neutrophils by chimeric 5.12.14 Fab with an average IC₅₀ of 1.6 nM.

Figure 5 is a graph depicting the inhibition of binding of ¹²⁵I-IL-8 to human neutrophils by chimeric 6G.4.25 Fab with an average IC₅₀ of 7.5 nM.

Figure 6 demonstrates the inhibition of human IL-8 mediated neutrophil chemotaxis by chimeric 6G4.2.5 Fab and chimeric 5.12.14 Fab.

Figure 7 demonstrates the relative abilities of chimeric 6G4.2.5 Fab and chimeric 5.12.14 Fab to inhibit rabbit IL-8 mediated neutrophil chemotaxis.

Figure 8 depicts the stimulation of elastase release from human neutrophils by various concentrations of human and rabbit IL-8. The relative extent of elastase release was quantitated by measurement of absorbance at 405 nm. The data represent mean \pm SEM of triplicate samples.

Figure 9 is a graph depicting the ability of chimeric 6G4.2.5 Fab and chimeric 5.12.14 Fab to inhibit elastase release from human neutrophils stimulated by human IL-8. The results were normalized to reflect the percentage of elastase release elicited by 100 nM IL-8 alone. The data represent the mean \pm SEM of three separate experiments performed on different days with different blood donors. IC₅₀ values were calculated by four parameter fit.

Figure 10 is a graph depicting the relative abilities of chimeric 6G4.2.5 Fab and chimeric 5.12.14 Fab to inhibit elastase release from human neutrophils stimulated by rabbit IL-8. The results were normalized to reflect the percentage of elastase release elicited by 100 nM IL-8 alone. The data represent the mean \pm SEM of three separate experiments performed on different days with different blood donors. IC₅₀ values were calculated by four parameter fit.

Figures 11A-11J are a set of graphs depicting the following parameters in a rabbit ulcerative colitis model: Figure 11A depicts myeloperoxidase levels in tissue; Figure 11B depicts IL-8 levels in tissue; Figure 11C depicts colon weight; Figure 11D depicts gross inflammation; Figure 11E depicts edema; Figure 11F depicts extent of necrosis; Figure 11G depicts severity of necrosis; Figure 11H depicts neutrophil margination; Figure 11I depicts neutrophil infiltration; and Figure 11J depicts mononuclear infiltration.

Figure 12 is a graph depicting the effect of anti-IL-8 monoclonal antibody treatment on the number of neutrophils in bronchoalveolar lavage (BAL) fluid in animals infected with Streptococcus pneumoniae, Escherichia coli, or Pseudomonas aeruginosa. Treatment with 6G4.2.5 significantly reduced the number of neutrophils present in the BAL fluid compared to animals treated with isotype control mouse IgG (Figure 12).

Figure 13 depicts the DNA sequences (SEQ ID NOS: 1-6) of three primers designed for each of the light and heavy chains. Multiple primers were designed in order to increase the chances of primer hybridization and efficiency of first strand cDNA synthesis for cloning the variable light and heavy regions of monoclonal antibody 5.12.14.

Figure 14 depicts the DNA sequences (SEQ ID NOS: 7-10) of one forward primer and one reverse primer for the 5.12.14 light chain variable region amplification.

Figure 15 depicts the DNA sequences (SEQ ID NOS: 11-15) of one forward primer and one reverse primer for the 5.12.14 heavy chain variable region amplification.

Figure 16 depicts the DNA sequence (SEQ ID NO: 16) and the amino acid sequence (SEQ ID NO: 17) of the 5.12.14 light chain variable region and partial murine constant light region. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). Important restriction sites are indicated in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable-light region is amino acids 1 to 109. The partial murine constant light region is amino acids 110 to 123 (in italics).

Figure 17 depicts the DNA sequence (SEQ ID NO: 18) and the amino acid sequence (SEQ ID NO: 19) of the 5.12.14 heavy chain variable region and partial murine constant heavy region. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). Important restriction sites are indicated in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable heavy region is amino acids 1 to 120. The partial murine constant heavy region is amino acids 121 to 130.

Figure 18 depicts the DNA sequences (SEQ ID NOS: 20-23) of amplification primers used to convert murine light and heavy chain constant region residues to their human equivalents.

Figure 19 depicts the DNA sequence (SEQ ID NO: 24) and the amino acid sequence (SEQ ID NO: 25) for the 5.12.14 light chain variable region and the human IgG1 light chain constant region. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). The human constant region is denoted in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable light region is amino acids 1 to 109. The human constant light region is amino acids 110 to 215.

Figures 20A-20B depict the DNA sequence (SEQ ID NO: 26) and the amino acid sequence (SEQ ID NO: 27) for the 5.12.14 heavy chain variable region and the heavy chain constant region of human IgG1. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). The human constant region is denoted in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable heavy region is amino acids 1 to 120. The human constant heavy region is amino acids 121 to 229.

Figure 21 depicts the DNA sequences (SEQ ID NOS: 1-6) of three primers designed for each of the light and heavy chains. Multiple primers were designed in order to increase the chances of primer hybridization and efficiency of first strand cDNA synthesis for cloning the variable light and heavy regions of monoclonal antibody 6G4.2.5.

Figure 22 depicts the DNA sequences (SEQ ID NOS: 28-31) of one forward primer and one reverse primer for the 6G4.2.5 light chain variable region amplification.

Figure 23 depicts the DNA sequences (SEQ ID NOS: 32,33,11,15,14, and 13) of one forward primer and one reverse primer for the 6G4.2.5 heavy chain variable region amplification.

Figure 24 depicts the DNA sequence (SEQ ID NO: 34) and the amino acid sequence (SEQ ID NO: 35) of the 6G4.2.5 light chain variable region and partial murine constant light region. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). Useful cloning sites are in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable light region is amino acids 1 to 114. The partial murine constant light region is amino acids 115 to 131.

Figure 25 depicts the DNA sequence (SEQ ID NO: 36) and the amino acid sequence (SEQ ID NO: 37) of the 6G4.2.5 heavy chain variable region and partial murine constant heavy region. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). Useful cloning sites are in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable heavy region is amino acids 1 to 122. The partial murine constant heavy region is amino

acids 123 to 135.

Figure 26 depicts the DNA sequences (SEQ ID NOS: 38-40) of primers to convert the murine light chain and heavy chain constant regions to their human equivalents.

Figures 27A-27B depict the DNA sequence (SEQ ID NO: 41) and the amino acid sequence (SEQ ID NO: 42) for the chimeric 6G4.2.5 light chain. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). The human constant region is denoted in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable heavy region is amino acids 1 to 114. The human constant heavy region is amino acids 115 to 220.

Figures 28A-28B depict the DNA sequence (SEQ ID NO: 43) and the amino acid sequence (SEQ ID NO: 44) for the chimeric 6G4.2.5 heavy chain. CDRs are indicated by either X-ray crystallography (underlined amino acids) or by Kabat sequence comparison (amino acids denoted with asterisk). The human constant region is denoted in italics. The signal peptide of STII is amino acids -23 to -1. The murine variable heavy region is amino acids 1 to 122. The human constant heavy region is amino acids 123 to 231.

Fig. 29 depicts an amino acid sequence alignment of murine 6G425 light chain variable domain (SEQ ID NO: 45), humanized 6G425 F(ab)-1 light chain variable domain (SEQ ID NO: 46), and human light chain κ I consensus framework (SEQ ID NO: 47) amino acid sequences, and an amino acid sequence alignment of murine 6G425 heavy chain variable domain (SEQ ID NO: 48), humanized 6G425 F(ab)-1 heavy chain variable domain (SEQ ID NO: 49), and human IgG1 subgroup III heavy chain variable domain (SEQ ID NO: 50) amino acid sequences, used in the humanization of 6G425. Light chain CDRs are labeled L1, L2, L3; heavy chain CDRs are labeled H1, H2, and H3. = and + indicate CDR sequences as defined by X-ray crystallographic contacts and sequence hypervariability, respectively. # indicates a difference between the aligned sequences. Residue numbering is according to Kabat *et al.* Lower case lettering denotes the insertion of an amino acid residue relative to the humIII consensus sequence numbering.

Figs. 30A, 30B and 30C are graphs depicting the ability of F(ab)-9 (humanized 6G4V11 Fab) to inhibit human wild type IL-8, human monomeric IL-8, and rhesus IL-8 mediated neutrophil chemotaxis, respectively. Fig. 30A presents inhibition data for F(ab)-9 samples at concentrations of 0.06 nM, 6.25 nM, 12.5 nM, 25 nM, 50 nM, and 100 nM, for an isotype control antibody (denoted "4D5") sample at a concentration of 100 nM, and for a no antibody control sample, in the presence of 2nM human wild type IL-8. Fig. 30B presents inhibition data for F(ab)-9 samples at concentrations of 6.25 nM, 12.5 nM, 25 nM, and 50 nM, for an isotype control antibody (denoted "4D5") sample at a concentration of 100 nM, and for a no antibody control sample, in the presence of 4 nM human monomeric IL-8 (denoted as "BD59" and as "monomeric IL-8"). Fig. 30C presents inhibition data for F(ab)-9 samples at concentrations of 1 nM, 12.5 nM, 25 nM, and 50 nM, for an isotype control antibody (denoted "4D5") sample at a concentration of 100 nM, and for a no antibody control sample, in the presence of 2 nM rhesus IL-8. In addition, Figs. 30A-30C each presents data for a no IL-8 buffer control sample (denoted as "Buffer") in the respective inhibition assay.

Fig. 31A depicts the amino acid sequences of the humanized anti-IL-8 6G4.2.5V11 light chain in an N-terminal fusion with the STII leader peptide (SEQ ID NO: 51), the humanized anti-IL-8 6G4.2.5V11 heavy chain in an N-terminal fusion with the STII leader peptide (SEQ ID NO: 52), and a peptide-linker in a C-terminal fusion with M13 phage gene-III coat protein (SEQ ID NO: 53).

Fig. 31B depicts the nucleic acid sequence (SEQ ID NO: 54) and the translated amino acid sequence (SEQ ID NO: 51) of the humanized anti-IL-8 6G4.2.5V11 light chain in an N-terminal fusion with the STII leader peptide.

Fig. 31C depicts the amino acid sequences of the humanized anti-IL-8 6G4.2.5V19 light chain in an N-terminal fusion with the STII leader peptide (SEQ ID NO: 51), and the humanized anti-IL-8 6G4.2.5V19 heavy chain in an N-terminal fusion with the STII leader peptide (SEQ ID NO: 55).

Fig. 32 is a three dimensional computer model of the humanized anti-IL-8 6G4.2.5V11 antibody. Heavy chain CDR loops and variable domain regions appear in purple, and CDR-H3 side chain residues appear in yellow. Heavy chain constant domain regions appear in red. Light chain CDR loops and variable domain regions appear in off-white, and the Asn residue at amino acid position 35 (N35) in CDR L1 appears in green. Light chain constant domain regions appear in amber.

Fig. 33 is a Scatchard plot depicting the inhibition of 125 I-IL-8 binding to human neutrophils exhibited by intact murine 6G4.2.5 antibody (denoted 6G4 murine mAb), 6G4.2.5 murine-human chimera Fab (denoted 6G4 chimera), humanized 6G4.2.5 Fab versions I and II (denoted VI and VII), and variant 6G4.2.5V11N35A Fab (denoted V11N35A).

Figs. 34A, 34B, 34C and 34D are graphs depicting the ability of 6G4.2.5V11N35A Fab to inhibit human wild type IL-8, human monomeric IL-8, rabbit IL-8, and rhesus IL-8 mediated neutrophil chemotaxis, respectively. Fig. 34A presents inhibition data for 6G4.2.5V11N35A Fab samples at concentrations of 0.5, 1, 2, 4, 8, 16, and 33 nM, for an isotype control antibody (denoted "4D5") sample at a concentration of 33 nM, and for a no antibody control (denoted "HuIL-8") sample, in the presence of 2 nM human wild type IL-8. Fig. 34B presents inhibition data for 6G4.2.5V11N35A Fab samples at concentrations of 0.5, 1, 2, 4, 8, 16, and 33 nM, for an intact 6G4.2.5 mAb sample at a concentration of 33 nM, for an isotype control antibody (denoted as "4D5") sample at a concentration of 33 nM, and for a no antibody control (denoted "BD59") sample, in the presence of 2 nM human monomeric IL-8. Fig. 34C presents inhibition data for 6G4.2.5V11N35A Fab samples at concentrations of 0.5, 1, 2, 4, 8, 16, and 33 nM, for an intact 6G4.2.5 mAb sample at a concentration of 33 nM, for an isotype control antibody (denoted "4D5") sample at a concentration of 33 nM, and for a no antibody control (denoted "Rab IL-8") sample, in the presence of 2 nM rabbit IL-8. Fig. 34D presents inhibition data for 6G4.2.5V11N35A Fab samples at concentrations of 0.5, 1, 2, 4, 8, 16, and 33 nM, for an intact 6G4.2.5 mAb sample at a concentration of 33 nM, for an isotype control antibody (denoted as "4D5") sample at a concentration of 33 nM, and for a no antibody control (denoted "Rhe IL-8") sample, in the presence of 2 nM rhesus IL-8. In addition, Figs. 34B-34D each presents data for human wild type IL-8 control (denoted "HuIL-8") samples at a concentration of 2 nM in the respective assay, and Figs. 34A-34D each presents data for a no IL-8 buffer control (denoted "Buffer") sample in the respective assay.

Fig. 35 depicts the amino acid sequences of the humanized anti-IL-8 6G4.2.5V11N35A light chain in an N-terminal fusion with the STII leader peptide (SEQ ID NO: 56), the humanized anti-IL-8 6G4.2.5V11N35A heavy chain in an N-terminal fusion with the STII leader peptide (SEQ ID NO: 52), and the GCN4 leucine zipper peptide (SEQ ID NO: 57). The Ala residue (substituted for the wild type Asn residue) at amino acid position 35 in the 6G4.2.5V11N35A light chain appears in bold case. A putative pepsin cleavage

site in the GCN4 leucine zipper sequence is underlined.

Fig. 36 depicts the DNA sequence (SEQ ID NO: 58) and the amino acid sequence (SEQ ID NO: 56) of the humanized anti-IL-8 6G4.2.5V11N35A light chain in an N-terminal fusion with the STII leader peptide. Complementarity determining regions L1, L2, and L3 are underlined

5 Figs. 37A-37B depict the DNA sequence (SEQ ID NO: 59) and the amino acid sequence (SEQ ID NO: 60) of the humanized anti-IL-8 6G4.2.5V11N35A heavy chain in an N-terminal fusion with the STII leader peptide and in a C-terminal fusion with the GCN4 leucine zipper sequence. Complementarity determining regions H1, H2, and H3 are underlined.

Fig. 38 is a Scatchard plot depicting the inhibition of 125 I-IL-8 binding to human neutrophils 10 exhibited by 6G4.2.5V11N35A Fab (denoted Fab), 6G4.2.5V11N35A F(ab')₂ (denoted F(ab')₂), and human wild type IL-8 control (denoted IL-8).

Fig. 39 is a graph depicting a comparison of the wild type human IL-8 mediated neutrophil chemotaxis inhibition activities of the 6G4.2.5V11N35A F(ab')₂ and 6G4.2.5V11N35A Fab. Inhibition data are presented for 6G4.2.5V11N35A Fab samples (denoted "N35A Fab") and 6G4.2.5V11N35A F(ab')₂ 15 samples (denoted N35A F(ab')₂) at concentrations of 0.3, 1, 3, 10, 30, and 100 nM, for an isotype control antibody (denoted as "4D5") sample at a concentration of 100 nM, and for a no antibody control sample, in the presence of 2 nM human wild type IL-8. In addition, inhibition data are presented for no IL-8 buffer control samples (denoted "Buffer").

Fig. 40 is a graph depicting the ability of 6G4.2.5V11N35A F(ab')₂ to inhibit human monomeric IL- 20 8, rhesus IL-8, and rabbit IL-8 mediated neutrophil chemotaxis. Human monomeric IL-8 mediated neutrophil chemotaxis data are presented for 6G4.2.5V11N35A F(ab')₂ samples at concentrations of 0.3, 1, 3, and 10 nM, for an isotype control antibody (denoted as "4D5") sample at a concentration of 100 nM, and for a no antibody control sample (denoted as "BD59"), in the presence of human monomeric IL-8 (denoted as "BD59") at a concentration of 0.5 nM. Rhesus IL-8 mediated neutrophil chemotaxis data are presented for 25 6G4.2.5V11N35A F(ab')₂ samples at concentrations of 0.3, 1, 3, and 10 nM, and for a no antibody control sample, in the presence of rhesus IL-8 at a concentration of 2 nM. Rabbit IL-8 mediated neutrophil chemotaxis data are presented for 6G4.2.5V11N35A F(ab')₂ samples at concentrations of 0.3, 1, 3, and 10 nM, and for a no antibody control sample, in the presence of rabbit IL-8 at a concentration of 2 nM. In addition, inhibition data are presented for a no IL-8 buffer control sample (denoted as "Buffer") and for a 2 nM human wild type 30 IL-8 (denoted as "HuIL-8").

Figs. 41A-41V depict the nucleic acid sequence (SEQ ID NO: 61) of the p6G4V11N35A.F(ab')₂ vector.

Fig. 42 depicts the nucleic acid sequences of the stop template primer (SEQ ID NO: 63) and the NNS randomization primer (SEQ ID NO: 64) used for random mutagenesis of amino acid position 35 in variable 35 light chain CDR-L1 of humanized antibody 6G4V11.

Fig. 43A is a table of data describing the frequencies of different phage display clones obtained from the randomization of amino acid position 35 in variable light chain CDR-L1 of humanized antibody 6G4V11.

Figs. 43B, 43C, 43D and 43E are graphs of displacement curves depicting the inhibition of 125 I-IL-8 binding to neutrophils exhibited by the 6G4V11N35A, 6G4V11N35D, 6G4V11N35E and 6G4V11N35G Fab's.

Fig. 44 contains a graph depicting the typical kinetics of an anti-IL-8 antibody fragment (6G4V11N35A F(ab')₂) binding to IL-8. Fig. 44 also contains a table of data providing the equilibrium constant for 6G4V11N35A Fab binding to IL-8 (rate constants were not determined "ND"), and the equilibrium and rate constants for 6G4V11N35A F(ab')₂ and 6G4V11N35E Fab binding to IL-8.

Fig. 45 depicts the DNA sequence (SEQ ID NO: 65) and amino acid sequence (SEQ ID NO: 62) of the 6G4V11N35E light chain in an N-terminal fusion with the STII leader peptide. Complementarity determining regions L1, L2 and L3 are underlined.

Fig. 46 is a graph depicting the ability of 6G4V11N35E Fab to inhibit human IL-8 (dark columns) and rabbit IL-8 (light columns) mediated neutrophil chemotaxis. Data are presented for 6G4V11N35E Fab samples at concentrations of 0.4, 1.2, 3.7, 11 and 33 nM, and for an isotype control antibody (4D5) sample at a concentration of 100 nM, in the presence of 2 nM human IL-8 or 2 nM rabbit IL-8. In addition, inhibition data are presented for a no IL-8 buffer control sample (denoted "Buffer") and for human and rabbit IL-8 control samples (denoted "IL-8").

Fig. 47 depicts the DNA sequence of the sense (SEQ ID NO: 66) and anti-sense (SEQ ID NO: 67) strands of a PvuII-XhoI synthetic nucleotide encoding amino acids Leu4 to Phe29 of the 6G4V11N35A heavy chain.

Figs. 48A-48T depict the DNA sequence (SEQ ID NO: 68) of plasmid p6G4V11N35A.choSD9.

Figs. 49A, 49B, 49C and 49D are graphs of displacement curves depicting the inhibition of 125 I-IL-8 binding to neutrophils exhibited by IL-8 control, intact murine 6G4.2.5 antibody, the full length IgG1 form of variant 6G4V11N35A, and the full length IgG1 form of variant 6G4V11N35E, respectively.

Figs. 50A-50B are graphs depicting the ability of full length 6G4V11N35A IgG1 and 6G4V11N35E IgG1 to inhibit human IL-8 (Fig. 50A) and rabbit IL-8 (Fig. 50B) mediated neutrophil chemotaxis.

Fig. 51 contains a graph depicting the typical kinetics of a full length anti-IL8 antibody (6G4V11N35A IgG1) binding to IL-8. Fig. 51 also contains a table of data providing the equilibrium and rate constants for full length murine 6G4.2.5 IgG2a, 6G4V11N35A IgG1 and 6G4V11N35E IgG1 binding to IL-8.

Figs. 52A and 52B are graphs of displacement curves depicting the results of an unlabeled IL-8/ 125 I-IL-8 competition radioimmunoassay performed with full length 6G4V11N35A IgG1 and 6G4V11N35E IgG1, respectively.

Fig. 53 depicts the DNA sequence (SEQ ID NO: 69) and amino acid sequence (SEQ ID NO: 70) of the 6G4V11N35A Fab' heavy chain (6G4V11N35A Fab heavy chain modified to contain a cysteine residue in the hinge region).

Figs. 54A-54C contain graphs of displacement curves depicting the IL-8 binding and IC₅₀'s for PEG-

maleimide modified 6G4V11N35A Fab' molecules.

Figs. 55A-55C are graphs depicting the ability of PEG-maleimide modified 6G4V11N35A Fab' molecules to inhibit human IL-8 and rabbit IL-8 mediated neutrophil chemotaxis.

Figs. 56A-56C are graphs depicting the ability of PEG-maleimide modified 6G4V11N35A Fab' molecules to inhibit IL-8 mediated release of β -glucuronidase from neutrophils.

Figs. 57A-57B contain graphs of displacement curves depicting the inhibition of 125 I-IL-8 binding to neutrophils exhibited by PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules.

Figs. 58A-58B are graphs depicting the ability of PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules to inhibit human IL-8 mediated neutrophil chemotaxis.

Figs. 59A-59B are graphs depicting the ability of PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules to inhibit human IL-8 mediated release of β -glucuronidase from neutrophils.

Fig. 60 is a graph depicting the theoretical molecular weight (dotted bars) and effective size (solid bars) of PEG-maleimide modified 6G4V11N35A Fab' molecules as determined by SEC-HPLC.

Figs. 61A and 61B are SDS-PAGE gels depicting the electrophoretic mobility of various PEG-maleimide modified 6G4V11N35A Fab' molecules under reducing and non-reducing conditions, respectively.

Fig. 62 contains size exclusion chromatograms (SEC-HPLC) depicting the retention times and effective (hydrodynamic) sizes of various PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules.

Fig. 63 is a graph depicting the theoretical molecular weight (open columns), effective size determined by SEC-HPLC (solid columns), and the actual molecular weight determined by SEC-light scattering (shaded columns) for various PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules.

Fig. 64 is an SDS-PAGE gel depicting the electrophoretic mobility of various PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules. From left to right, lane 1 contains unmodified F(ab')₂, lane 2 contains F(ab')₂ coupled to two 40 kD branched PEG-succinimide molecules (denoted "Br(2)-40kD(N)-F(ab')₂"), lane 3 contains F(ab')₂ coupled to one 40 kD branched PEG-succinimide molecule (denoted "Br(1)-40kD(N)-Fab'2"), lane 4 contains a mixture of F(ab')₂ coupled to four 20 kD linear PEG-succinimide molecules and F(ab')₂ coupled to five 20 kD linear PEG-succinimide molecules (denoted "L(4+5)-20kD-(N)-Fab'2"), lane 5 contains F(ab')₂ coupled to one 20 kD linear PEG-succinimide molecule (denoted "L(1)-20kD-(N)-Fab'2"), and lane 6 contains molecular weight standards.

Figs. 65A and 65B are graphs comparing the serum concentration vs. time profiles of various PEG-maleimide modified 6G4V11N35A Fab' molecules (Fig. 65A) and various PEG-succinimide modified 6G4V11N35A F(ab')₂ molecules (Fig. 65B) in rabbits. In Fig. 65A, "bran.(1)40K(s)Fab' " denotes 6G4V11N35A Fab' coupled to one 40 kD branched PEG-maleimide molecule, "lin.(1)40K(s)Fab' " denotes 6G4V11N35A Fab' coupled to one 40 kD linear PEG-maleimide molecule, "lin.(1)30K(s)Fab' " denotes 6G4V11N35A Fab' coupled to one 30 kD linear PEG-maleimide molecule, "lin.(1)20K(s)Fab' " denotes

6G4V11N35A Fab' coupled to one 20 kD linear PEG-maleimide molecule. In Fig. 65B, "bran.(2)40K(N)Fab'2" denotes 6G4V11N35A F(ab')₂ coupled to two 40 kD branched PEG-succinimide molecules, "bran.(1)40K(N)Fab'2" denotes 6G4V11N35A F(ab')₂ coupled to one 40 kD branched PEG-succinimide molecule, and "Fab'2" denotes unmodified 6G4V11N35A F(ab')₂. In both Figs. 65A and 65B, "IgG" denotes a full length IgG1 equivalent of the human-murine chimeric anti-rabbit IL-8 Fab described in Example F below.

Fig. 66 contains graphs comparing the serum concentration vs. time profiles of 6G4V11N35A Fab' coupled to one 40 kD branched PEG-maleimide molecule (denoted as "bran.(1)40K(s)Fab'"), 6G4V11N35A F(ab')₂ coupled to one 40 kD branched PEG-succinimide molecule (denoted as "bran.(1)40K(N)Fab'2"), unmodified 6G4V11N35A F(ab')₂ (denoted as "Fab'2"), unmodified 6G4V11N35A Fab' (denoted as "Fab'"), and a full length IgG1 (denoted as "IgG") equivalent of the human-murine chimeric anti-rabbit IL-8 Fab described in Example F below.

Fig. 67 is a graph depicting the effect of 6G4V11N35A Fab' coupled to one 40 kD branched PEG-maleimide molecule (denoted as "PEG 40 Kd") and murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 (full length IgG2a) (denoted as "6G4.2.5") on gross weight of entire lung in an ARDS rabbit model.

Fig. 68 is a graph depicting the effect of 6G4V11N35A Fab' coupled to one branched 40 kD PEG-maleimide molecule (denoted as "PEG 40-Kd") and murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 (full length IgG2a) (denoted as "6G4.2.5") on BAL total leukocyte (light columns) and polymorphonuclear cell (dark columns) counts in an ARDS rabbit model. Untreated (no therapeutics) control animal data is denoted as "Control".

Fig. 69 is a graph depicting the effect of 6G4V11N35A Fab' coupled to one branched 40 kD PEG-maleimide molecule (denoted as "PEG 40 Kd") and murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 (full length IgG2a) (denoted as "6G4.2.5") on PaO₂/FiO₂ ratio at 24 hours-post treatment (light columns) and 48 hours post-treatment (dark columns) in an ARDS rabbit model. Untreated (no therapeutics) control animal data is denoted as "Control".

Fig. 70A is a graph depicting PaO₂/FiO₂ ratios obtained in 100% oxygen at 24 hours after acid instillation for: (1) rabbits (n=5) treated with 7 mg/kg IV 20 kD linear PEG-6G4V11N35E Fab' at 10 minutes before and 6 hours after acid instillation, (2) rabbits (n=7) treated with 5 mg/kg IV full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 at 10 minutes before acid instillation, (3) rabbits (n=3) treated with 5 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, (4) rabbits (n=2) treated with 20 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, and (5) rabbits (n=25) treated with 5 ml IV saline at 10 minutes before and 6 hours after acid instillation.

Fig. 70B is a graph depicting PaO₂/FiO₂ ratios obtained in 100% oxygen at 48 hours after acid instillation for: (1) rabbits (n=5) treated with 7 mg/kg IV 20 kD linear PEG-6G4V11N35E Fab' at 10 minutes before and 6 hours after acid instillation, (2) rabbits (n=7) treated with 5 mg/kg IV full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 at 10 minutes before acid instillation, (3) rabbits (n=3) treated with 5 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, (4) rabbits (n=2)

treated with 20 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, and (5) rabbits (n=16) treated with 5 ml IV saline at 10 minutes before and 6 hours after acid instillation.

Fig. 70C is a graph depicting gross lung weight (in grams)/body weight (in kilograms) ratios (denoted as "GLW/BW Ratio") obtained at 72 hours post reperfusion for: (1) rabbits (n=5) treated with 7 mg/kg IV 20 kD linear PEG-6G4V11N35E Fab' at 10 minutes before and 6 hours after acid instillation, (2) rabbits (n=7) treated with 5 mg/kg IV full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 at 10 minutes before acid instillation, (3) rabbits (n=3) treated with 5 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, (4) rabbits (n=3) treated with 20 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, and (5) rabbits (n=29) treated with 5 ml IV saline at 10 minutes before and 6 hours after acid instillation.

Fig. 70D is a graph depicting total leukocyte (WBC) count in BAL fluid (represented in millions of cells counted in 20 ml BAL fluid) obtained at 72 hours post reperfusion for: (1) rabbits (n=5) treated with 7 mg/kg IV 20 kD linear PEG-6G4V11N35E Fab' at 10 minutes before and 6 hours after acid instillation, (2) rabbits (n=7) treated with 5 mg/kg IV full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 at 10 minutes before acid instillation, (3) rabbits (n=3) treated with 5 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, (4) rabbits (n=3) treated with 20 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, and (5) rabbits (n=11) treated with 5 ml IV saline at 10 minutes before and 6 hours after acid instillation.

Fig. 70E is a graph depicting total polymorphonuclear (PMN) cell count in BAL fluid (represented in millions of cells counted in 20 ml BAL fluid) obtained at 72 hours post reperfusion for: (1) rabbits (n=5) treated with 7 mg/kg IV 20 kD linear PEG-6G4V11N35E Fab' at 10 minutes before and 6 hours after acid instillation, (2) rabbits (n=7) treated with 5 mg/kg IV full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 at 10 minutes before acid instillation, (3) rabbits (n=3) treated with 5 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, (4) rabbits (n=3) treated with 20 mg/kg IV 40 kD branched PEG-6G4V11N35A Fab' at 10 minutes before acid instillation, and (5) rabbits (n=9) treated with 5 ml IV saline at 10 minutes before and 6 hours after acid instillation.

Fig. 71 is a graph depicting the effect of pegylated anti-IL-8 Fab' (as measured by percent change in ear volume at 1, 2 and 3 days post reperfusion) in a rabbit ear model of ischemia reperfusion injury. The data points from animals treated with empty vehicle (n=11), full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 (n=4), 20 kD linear PEG-6G4V11N35E Fab' (n=3), 30 kD linear PEG-6G4V11N35E Fab' (n=3), and 40 kD branched PEG-6G4V11N35E Fab' (n=3) are denoted by open boxes, open diamonds, open circles, open triangles, and crossed boxes, respectively.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. DEFINITIONS

In general, the following words or phrases have the indicated definition when used in the description, examples, and claims.

"Polymerase chain reaction" or "PCR" refers to a procedure or technique in which minute amounts of a specific piece of nucleic acid, RNA and/or DNA, are amplified as described in U.S. Patent No. 4,683,195 issued 28 July 1987. Generally, sequence information from the ends of the region of interest or beyond needs

to be available, such that oligonucleotide primers can be designed; these primers will be identical or similar in sequence to opposite strands of the template to be amplified. The 5' terminal nucleotides of the two primers can coincide with the ends of the amplified material. PCR can be used to amplify specific RNA sequences, specific DNA sequences from total genomic DNA, and cDNA transcribed from total cellular RNA, bacteriophage or plasmid sequences, etc. See generally Mullis *et al.*, Cold Spring Harbor Symp. Quant. Biol. 51:263 (1987); Erlich, ed., PCR Technology (Stockton Press, NY, 1989). As used herein, PCR is considered to be one, but not the only, example of a nucleic acid polymerase reaction method for amplifying a nucleic acid test sample comprising the use of a known nucleic acid as a primer and a nucleic acid polymerase to amplify or generate a specific piece of nucleic acid.

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules which lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

"Native antibodies and immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains (Clothia *et al.*, J. Mol. Biol. 186:651 (1985); Novotny and Haber, Proc. Natl. Acad. Sci. U.S.A. 82:4592 (1985)).

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, Sequences of Proteins of Immunological Interest, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability

to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. In a two-chain Fv species, this region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv species (scFv), one heavy- and one light-chain variable domain can be covalently linked by a flexible peptide linker such that the light and heavy chains can associate in a "dimeric" structure analogous to that in a two-chain Fv species. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site. For a review of scFv see Pluckthun, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these can be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "antibody" is used in the broadest sense and specifically covers single monoclonal antibodies (including agonist and antagonist antibodies) and antibody compositions with polyepitopic specificity.

"Antibody fragment", and all grammatical variants thereof, as used herein are defined as a portion of an intact antibody comprising the antigen binding site or variable region of the intact antibody, wherein the portion is free of the constant heavy chain domains (i.e. CH2, CH3, and CH4, depending on antibody isotype) of the Fc region of the intact antibody. Examples of antibody fragments include Fab, Fab', Fab'-SH, $F(ab')_2$, and Fv fragments; diabodies; any antibody fragment that is a polypeptide having a primary structure consisting of one uninterrupted sequence of contiguous amino acid residues (referred to herein as a "single-chain antibody

fragment" or "single chain polypeptide"), including without limitation (1) single-chain Fv (scFv) molecules (2) single chain polypeptides containing only one light chain variable domain, or a fragment thereof that contains the three CDRs of the light chain variable domain, without an associated heavy chain moiety and (3) single chain polypeptides containing only one heavy chain variable region, or a fragment thereof containing the three CDRs of the heavy chain variable region, without an associated light chain moiety; and multispecific or multivalent structures formed from antibody fragments. In an antibody fragment comprising one or more heavy chains, the heavy chain(s) can contain any constant domain sequence (e.g. CH1 in the IgG isotype) found in a non-Fc region of an intact antibody, and/or can contain any hinge region sequence found in an intact antibody, and/or can contain a leucine zipper sequence fused to or situated in the hinge region sequence or the constant domain sequence of the heavy chain(s). Suitable leucine zipper sequences include the jun and fos leucine zippers taught by Kostelny et al., *J. Immunol.*, **148**: 1547-1553 (1992) and the GCN4 leucine zipper described in the Examples below.

Unless specifically indicated to the contrary, the term "conjugate" as described and claimed herein is defined as a heterogeneous molecule formed by the covalent attachment of one or more antibody fragment(s) to one or more polymer molecule(s), wherein the heterogeneous molecule is water soluble, i.e. soluble in physiological fluids such as blood, and wherein the heterogeneous molecule is free of any structured aggregate.

In the context of the foregoing definition, the term "structured aggregate" refers to (1) any aggregate of molecules in aqueous solution having a spheroid or spheroid shell structure, such that the heterogeneous molecule is not in a micelle or other emulsion structure, and is not anchored to a lipid bilayer, vesicle or liposome; and (2) any aggregate of molecules in solid or insolubilized form, such as a chromatography bead matrix, that does not release the heterogeneous molecule into solution upon contact with an aqueous phase. Accordingly, the term "conjugate" as defined herein encompasses the aforementioned heterogeneous molecule in a precipitate, sediment, bioerodible matrix or other solid capable of releasing the heterogeneous molecule into aqueous solution upon hydration of the solid.

Unless specifically indicated to the contrary, the terms "polymer", "polymer molecule", "nonproteinaceous polymer", and "nonproteinaceous polymer molecule" are used interchangeably and are defined as a molecule formed by covalent linkage of two or more monomers, wherein none of the monomers is contained in the group consisting of alanine (Ala), cysteine (Cys), aspartic acid (Asp), glutamic acid (Glu), phenylalanine (Phe), glycine (Gly), histidine (His), isoleucine (Ile), lysine (Lys), leucine (Leu), methionine (Met), asparagine (Asn), proline (Pro), glutamine (Gln), arginine (Arg), serine (Ser), threonine (Thr), valine (Val), tryptophan (Trp), and tyrosine (Tyr) residues.

The term "monoclonal antibody" (mAb) as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts.

Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each mAb is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they can be synthesized by hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal"

indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567 to Cabilly *et al.*). The "monoclonal antibodies" also include clones of antigen-recognition and binding-site containing antibody fragments (Fv clones) isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein include hybrid and recombinant antibodies produced by splicing a variable (including hypervariable) domain of an anti-IL-8 antibody with a constant domain (e.g. "humanized" antibodies), or a light chain with a heavy chain, or a chain from one species with a chain from another species, or fusions with heterologous proteins, regardless of species of origin or immunoglobulin class or subclass designation, as well as antibody fragments (e.g., Fab, F(ab')₂, and Fv), so long as they exhibit the desired biological activity. (See, e.g., U.S. Pat. No. 4,816,567 to Cabilly *et al.*; Mage and Lamoyi, in Monoclonal Antibody Production Techniques and Applications, pp. 79-97 (Marcel Dekker, Inc., New York, 1987).)

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (Cabilly *et al.*, *supra*; Morrison *et al.*, Proc. Natl. Acad. Sci. U.S.A. 81:6851 (1984)).

"Humanized" forms of non-human (e.g., murine) antibodies are specific chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂, or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies can comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details see Jones *et al.*, *Nature* 321:522 (1986); Reichmann *et al.*, *Nature* 332:323 (1988); and Presta, Curr. Op. Struct. Biol. 2:593 (1992).

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in which the disorder is to be prevented.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc. Preferably, the mammal herein is human.

As used herein, protein, peptide and polypeptide are used interchangeably to denote an amino acid polymer or a set of two or more interacting or bound amino acid polymers.

As used herein, the term "inflammatory disorders" refers to pathological states resulting in inflammation, typically caused by neutrophil chemotaxis. Examples of such disorders include inflammatory skin diseases including psoriasis and atopic dermatitis; systemic scleroderma and sclerosis; responses associated with inflammatory bowel disease (such as Crohn's disease and ulcerative colitis); ischemic reperfusion disorders including surgical tissue reperfusion injury, myocardial ischemic conditions such as myocardial infarction, cardiac arrest, reperfusion after cardiac surgery and constriction after percutaneous transluminal coronary angioplasty, stroke, and abdominal aortic aneurysms; cerebral edema secondary to stroke; cranial trauma; hypovolemic shock; asphyxia; adult respiratory distress syndrome; acute lung injury; Behçet's Disease; dermatomyositis; polymyositis; multiple sclerosis; dermatitis; meningitis; encephalitis; uveitis; osteoarthritis; lupus nephritis; autoimmune diseases such as rheumatoid arthritis, Sjorgen's syndrome, vasculitis; diseases involving leukocyte diapedesis; central nervous system (CNS) inflammatory disorder, multiple organ injury syndrome secondary to septicemia or trauma; alcoholic hepatitis; bacterial pneumonia; antigen-antibody complex mediated diseases including glomerulonephritis; sepsis; sarcoidosis; immunopathologic responses to tissue/organ transplantation; inflammations of the lung, including pleurisy, alveolitis, vasculitis, pneumonia, chronic bronchitis, bronchiectasis, diffuse panbronchiolitis, hypersensitivity pneumonitis, idiopathic pulmonary fibrosis (IPF), and cystic fibrosis; etc. The preferred indications include acute lung injury, adult respiratory distress syndrome, ischemic reperfusion (including surgical tissue reperfusion injury, myocardial ischemia, and acute myocardial infarction), hypovolemic shock, asthma, bacterial pneumonia and inflammatory bowel disease such as ulcerative colitis.

As used herein, the terms "asthma", "asthmatic disorder", "asthmatic disease", and "bronchial asthma" refer to a condition of the lungs in which there is widespread narrowing of lower airways. "Atopic asthma" and "allergic asthma" refer to asthma that is a manifestation of an IgE-mediated hypersensitivity reaction in the lower airways, including, e.g., moderate or severe chronic asthma, such as conditions requiring the frequent or constant use of inhaled or systemic steroids to control the asthma symptoms. A preferred indication is allergic asthma.

The terms "hydrodynamic size", "apparent size", "apparent molecular weight", "effective size" and "effective molecular weight" of a molecule are used synonymously herein refer to the size of a molecule as determined by comparison to a standard curve produced with globular protein molecular weight standards in a size exclusion chromatography system, wherein the standard curve is created by mapping the actual molecular weight of each standard against its elution time observed in the size exclusion chromatography system. Thus, the apparent size of a test molecule is derived by using the molecule's elution time to extrapolate a putative

molecular weight from the standard curve. Preferably, the molecular weight standards used to create the standard curve are selected such that the apparent size of the test molecule falls within the linear portion of the standard curve.

II. MODES FOR CARRYING OUT THE INVENTION

5 In one part, the invention arises from the surprising and unexpected discovery that antibody fragment-polymer conjugates having an effective or apparent size significantly greater than the antibody fragment-polymer conjugates described in the art confers an increase in serum half-life, an increase in mean residence time in circulation (MRT), and/or a decrease in serum clearance rate over underivatized antibody fragment which far exceed the modest changes in such biological property or properties obtained with the art-known
10 antibody fragment-polymer conjugates. The present inventors have determined for the first time that increasing the effective size of an antibody fragment to at least about 500,000 D, or increasing the effective size of an antibody fragment by at least about 8 fold over the effective size of the parental antibody fragment, or derivatizing an antibody fragment with a polymer of at least about 20,000 D in molecular weight, yields a molecule with a commercially useful pharmacokinetic profile. The greatly extended serum half-life, extended
15 MRT, and/or reduced serum clearance rate of the conjugates of the invention makes such conjugates viable alternatives to intact antibodies used for therapeutic treatment of many disease indications. Antibody fragments provide significant advantages over intact antibodies, notably the fact that recombinant antibody fragments can be made in bacterial cell expression systems. Bacterial cell expression systems provide several advantages over mammalian cell expression systems, including reduced time and cost at both the research and
20 development and manufacturing stages of a product.

In another part, the present invention also arises from the humanization of the 6G4.2.5 murine anti-rabbit IL-8 monoclonal antibody ("6G4.2.5") described in WO 95/23865 (PCT/US95/02589 published September 8, 1995), the entire disclosure of which is specifically incorporated herein by reference. The hybridoma producing antibody 6G4.2.5 was deposited on September 28, 1994 with the American Type Culture
25 Collection and assigned ATCC Accession No. HB 11722 as described in the Examples below. In one aspect, the invention provides a humanized derivative of the 6G4.2.5 antibody, variant 11 (referred to herein as "6G4.2.5v11"), in which the murine CDRs of 6G4.2.5 are grafted onto a consensus framework for human light chain κ 1 and human IgG1 heavy chain subgroup III, followed by importing three framework residues from the murine 6G4.2.5 parent heavy chain variable domain sequence into analogous sites in the heavy chain variable
30 domain of the human template sequence, as described in the Examples below. In another aspect, the invention provides variants of the 6G4.2.5v11 antibody with certain amino acid substitution(s) yielding increased affinity for human IL-8 and/or promoting greater efficiency in recombinant manufacturing processes.

It will be understood that in the context of this Section (II) and all subsections thereof, every reference to "an antibody fragment" or "the antibody fragment" contained in a conjugate shall be a reference to one or
35 more antibody fragment(s) in the conjugate (consistent with the definition of the term "conjugate" set forth in Section (I) above), except where the number of antibody fragment(s) in the conjugate is expressly indicated. It will be understood that in the context of this Section (II) and all subsections thereof, every reference to "a polymer", "a polymer molecule", "the polymer", or "the polymer molecule" contained in a conjugate shall be a reference to one or more polymer molecule(s) in the conjugate (consistent with the definition of the term

"conjugate" set forth in Section (I) above), except where the number of polymer molecule(s) in the conjugate is expressly indicated.

1. LARGE EFFECTIVE SIZE ANTIBODY FRAGMENT-POLYMER CONJUGATES

In one aspect, the invention provides an antibody fragment covalently attached to a polymer to form a conjugate having an effective or apparent size of at least about 500,000 Daltons (D). In another aspect, the invention provides an antibody fragment covalently attached to a polymer to form a conjugate having an apparent size that is at least about 8 fold greater than the apparent size of the parental antibody fragment. In yet another aspect, the invention provides an antibody fragment covalently attached to a polymer of at least about 20,000 D in molecular weight (MW). It will be appreciated that the unexpectedly and surprisingly large increase in antibody fragment serum half-life, increase in MRT, and/or decrease in serum clearance rate can be achieved by using any type of polymer or number of polymer molecules which will provide the conjugate with an effective size of at least about 500,000 D, or by using any type of polymer or number of polymer molecules which will provide the conjugate with an effective size that is at least about 8 fold greater than the effective size of the parental antibody fragment, or by using any type or number of polymers wherein each polymer molecule is at least about 20,000 D in MW. Thus, the invention is not dependent on the use of any particular polymer or molar ratio of polymer to antibody fragment in the conjugate.

In addition, the beneficial aspects of the invention extend to antibody fragments without regard to antigen specificity. Although variations from antibody to antibody are to be expected, the antigen specificity of a given antibody will not substantially impair the extraordinary improvement in serum half-life, MRT, and/or serum clearance rate for antibody fragments thereof that can be obtained by derivatizing the antibody fragments as taught herein.

In one embodiment, the conjugate has an effective size of at least about 500,000 D, or at least about 800,000 D, or at least about 900,000 D, or at least about 1,000,000 D, or at least about 1,200,000 D, or at least about 1,400,000 D, or at least about 1,500,000 D, or at least about 1,800,000 D, or at least about 2,000,000 D, or at least about 2,500,000 D.

In another embodiment, the conjugate has an effective size of at or about 500,000 D to at or about 10,000,000 D, or an effective size of at or about 500,000 D to at or about 8,000,000 D, or an effective size of at or about 500,000 D to at or about 5,000,000 D, or an effective size of at or about 500,000 D to at or about 4,000,000 D, or an effective size of at or about 500,000 D to at or about 3,000,000 D, or an effective size of at or about 500,000 D to at or about 2,500,000 D, or an effective size of at or about 500,000 D to at or about 2,000,000 D, or an effective size of at or about 500,000 D to at or about 1,800,000 D, or an effective size of at or about 500,000 D to at or about 1,600,000 D, or an effective size of at or about 500,000 D to at or about 1,500,000 D, or an effective size of at or about 500,000 D to at or about 1,000,000 D.

In another embodiment, the conjugate has an effective size of at or about 800,000 D to at or about 10,000,000 D, or an effective size of at or about 800,000 D to at or about 8,000,000 D, or an effective size of at or about 800,000 D to at or about 5,000,000 D, or an effective size of at or about 800,000 D to at or about 4,000,000 D, or an effective size of at or about 800,000 D to at or about 3,000,000 D, or an effective size of at or about 800,000 D to at or about 2,500,000 D, or an effective size of at or about 800,000 D to at or about 2,000,000 D, or an effective size of at or about 800,000 D to at or about 1,800,000 D, or an effective size of at

or about 800,000 D to at or about 1,600,000 D, or an effective size of at or about 800,000 D to at or about 1,500,000 D, or an effective size of at or about 800,000 D to at or about 1,000,000 D.

In another embodiment, the conjugate has an effective size of at or about 900,000 D to at or about 10,000,000 D, or an effective size of at or about 900,000 D to at or about 8,000,000 D, or an effective size of at or about 900,000 D to at or about 5,000,000 D, or an effective size of at or about 900,000 D to at or about 4,000,000 D, or an effective size of at or about 900,000 D to at or about 3,000,000 D, or an effective size of at or about 900,000 D to at or about 2,500,000 D, or an effective size of at or about 900,000 D to at or about 2,000,000 D, or an effective size of at or about 900,000 D to at or about 1,800,000 D, or an effective size of at or about 900,000 D to at or about 1,600,000 D, or an effective size of at or about 900,000 D to at or about 1,500,000 D.

In another embodiment, the conjugate has an effective size of at or about 1,000,000 D to at or about 10,000,000 D, or an effective size of at or about 1,000,000 D to at or about 8,000,000 D, or an effective size of at or about 1,000,000 D to at or about 5,000,000 D, or an effective size of at or about 1,000,000 D to at or about 4,000,000 D, or an effective size of at or about 1,000,000 D to at or about 3,000,000 D, or an effective size of at or about 1,000,000 D to at or about 2,500,000 D, or an effective size of at or about 1,000,000 D to at or about 2,000,000 D, or an effective size of at or about 1,000,000 D to at or about 1,800,000 D, or an effective size of at or about 1,000,000 D to at or about 1,600,000 D, or an effective size of at or about 1,000,000 D to at or about 1,500,000 D.

In a further embodiment, the conjugate has an effective size that is at least about 8 fold greater, or at least about 10 fold greater, or at least about 12 fold greater, or at least about 15 fold greater, or at least about 18 fold greater, or at least about 20 fold greater, or at least about 25 fold greater, or at least about 28 fold greater, or at least about 30 fold greater, or at least about 40 fold greater, than the effective size of the parental antibody fragment.

In another embodiment, the conjugate has an effective size that is about 8 fold to about 100 fold greater, or is about 8 fold to about 80 fold greater, or is about 8 fold to about 50 fold greater, or is about 8 fold to about 40 fold greater, or is about 8 fold to about 30 fold greater, or is about 8 fold to about 28 fold greater, or is about 8 fold to about 25 fold greater, or is about 8 fold to about 20 fold greater, or is about 8 fold to about 18 fold greater, or is about 8 fold to about 15 fold greater, than the effective size of the parental antibody fragment.

In another embodiment, the conjugate has an effective size that is about 12 fold to about 100 fold greater, or is about 12 fold to about 80 fold greater, or is about 12 fold to about 50 fold greater, or is about 12 fold to about 40 fold greater, or is about 12 fold to about 30 fold greater, or is about 12 fold to about 28 fold greater, or is about 12 fold to about 25 fold greater, or is about 12 fold to about 20 fold greater, or is about 12 fold to about 18 fold greater, or is about 12 fold to about 15 fold greater, than the effective size of the parental antibody fragment.

In another embodiment, the conjugate has an effective size that is about 15 fold to about 100 fold greater, or is about 15 fold to about 80 fold greater, or is about 15 fold to about 50 fold greater, or is about 15 fold to about 40 fold greater, or is about 15 fold to about 30 fold greater, or is about 15 fold to about 28 fold greater, or is about 15 fold to about 25 fold greater, or is about 15 fold to about 20 fold greater, or is about 15

fold to about 18 fold greater, than the effective size of the parental antibody fragment.

5 In another embodiment, the conjugate has an effective size that is about 18 fold to about 100 fold greater, or is about 18 fold to about 80 fold greater, or is about 18 fold to about 50 fold greater, or is about 18 fold to about 40 fold greater, or is about 18 fold to about 30 fold greater, or is about 18 fold to about 28 fold greater, or is about 18 fold to about 25 fold greater, or is about 18 fold to about 20 fold greater, than the effective size of the parental antibody fragment.

10 In another embodiment, the conjugate has an effective size that is about 20 fold to about 100 fold greater, or is about 20 fold to about 80 fold greater, or is about 20 fold to about 50 fold greater, or is about 20 fold to about 40 fold greater, or is about 20 fold to about 30 fold greater, or is about 20 fold to about 28 fold greater, or is about 20 fold to about 25 fold greater, than the effective size of the parental antibody fragment.

15 In another embodiment, the conjugate has an effective size that is about 25 fold to about 100 fold greater, or is about 25 fold to about 80 fold greater, or is about 25 fold to about 50 fold greater, or is about 25 fold to about 40 fold greater, or is about 25 fold to about 30 fold greater, or is about 25 fold to about 28 fold greater, than the effective size of the parental antibody fragment.

20 In another embodiment, the conjugate has an effective size that is about 28 fold to about 100 fold greater, or is about 28 fold to about 80 fold greater, or is about 28 fold to about 50 fold greater, or is about 28 fold to about 40 fold greater, or is about 28 fold to about 30 fold greater, than the effective size of the parental antibody fragment.

25 In another embodiment, the conjugate has an effective size that is about 30 fold to about 100 fold greater, or is about 30 fold to about 80 fold greater, or is about 30 fold to about 50 fold greater, or is about 30 fold to about 40 fold greater, than the effective size of the parental antibody fragment.

30 In another embodiment, the conjugate has an effective size that is about 40 fold to about 100 fold greater, or is about 40 fold to about 80 fold greater, or is about 40 fold to about 50 fold greater, than the effective size of the parental antibody fragment.

25 In still another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW of at least about 20,000 D.

In a further embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW of at least about 30,000 D.

30 In yet another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW of at least about 40,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW that is at or about 20,000 D to at or about 300,000 D, or is at or about 30,000 D to at or about 300,000 D, or is at or about 40,000 D to at or about 300,000 D.

35 In another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW that is at or about 20,000 D to at or about 100,000 D, or is at or about 30,000 D to at or about 100,000 D, or is at or about 40,000 D to at or about 100,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW that is at or about 20,000 D to at or about 70,000 D, or is at or about 30,000 D to at or about 70,000 D, or is at or about 40,000 D to at or about 70,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW that is at or about 20,000 D to at or about 50,000 D, or is at or about 30,000 D to at or about 50,000 D, or is at or about 40,000 D to at or about 50,000 D.

5 In another embodiment, the conjugate is an antibody fragment covalently attached to at least one polymer having an actual MW that is at or about 20,000 D to at or about 40,000 D, or is at or about 30,000 D to at or about 40,000 D.

The conjugates of the invention can be made using any suitable technique now known or hereafter developed for derivatizing antibody fragments with polymers. It will be appreciated that the invention is not limited to conjugates utilizing any particular type of linkage between an antibody fragment and a polymer.

10 The conjugates of the invention include species wherein a polymer is covalently attached to a non-specific site or non-specific sites on the parental antibody fragment, i.e. polymer attachment is not targeted to a particular region or a particular amino acid residue in the parental antibody fragment. In such embodiments, the coupling chemistry can, for example, utilize the free epsilon amino groups of lysine residues in the parental antibody as attachment sites for the polymer, wherein such lysine residue amino groups are randomly
15 derivatized with polymer.

In addition, the conjugates of the invention include species wherein a polymer is covalently attached to a specific site or specific sites on the parental antibody fragment, i.e. polymer attachment is targeted to a particular region or a particular amino acid residue or residues in the parental antibody fragment. In such
20 embodiments, the coupling chemistry can, for example, utilize the free sulfhydryl group of a cysteine residue not in a disulfide bridge in the parental antibody fragment. In one embodiment, one or more cysteine residue(s) is (are) engineered into a selected site or sites in the parental antibody fragment for the purpose of providing a specific attachment site or sites for polymer. The polymer can be activated with any functional group that is capable of reacting specifically with the free sulfhydryl or thiol group(s) on the parental antibody, such as maleimide, sulfhydryl, thiol, triflate, tesa[®], aziridine, epoxide, and 5-pyridyl functional groups. The
25 polymer can be coupled to the parental antibody fragment using any protocol suitable for the chemistry of the coupling system selected, such as the protocols and systems described in Section (II)(1)(b) or in Section (T) of the Examples below.

In another embodiment, polymer attachment is targeted to the hinge region of the parental antibody fragment. The location of the hinge region varies according to the isotype of the parental antibody. Typically,
30 the hinge region of IgG, IgD and IgA isotype heavy chains is contained in a proline rich peptide sequence extending between the C_H1 and C_H2 domains. In a preferred embodiment, a cysteine residue or residues is (are) engineered into the hinge region of the parental antibody fragment in order to couple polymer specifically to a selected location in the hinge region.

In one aspect, the invention encompasses a conjugate having any molar ratio of polymer to antibody
35 fragment that endows the conjugate with an apparent size in the desired range as taught herein. The apparent size of the conjugate will depend in part upon the size and shape of the polymer used, the size and shape of the antibody fragment used, the number of polymer molecules attached to the antibody fragment, and the location of such attachment site(s) on the antibody fragment. These parameters can easily be identified and maximized

to obtain the a conjugate with the desired apparent size for any type of antibody fragment, polymer and linkage system.

In another aspect, the invention encompasses a conjugate with a polymer to antibody fragment molar ratio of no more than about 10:1, or no more than about 5:1, or no more than about 4:1, or no more than about 3:1, or no more than about 2:1, or no more than 1:1.

In yet another aspect, the invention encompasses a conjugate wherein the antibody fragment is attached to about 10 or fewer polymer molecules, each polymer molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In another embodiment, the conjugate contains an antibody fragment attached to about 5 or fewer polymer molecules, each polymer molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In still another embodiment, the conjugate contains an antibody fragment attached to about 4 or fewer polymer molecules, each polymer molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In a further embodiment, the conjugate contains an antibody fragment attached to about 3 or fewer polymer molecules, each polymer molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In an additional embodiment, the conjugate contains an antibody fragment attached to about 2 or fewer polymer molecules, each polymer molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. Also provided herein is a conjugate containing an antibody fragment attached to a single polymer molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D.

In still another aspect, the invention encompasses a conjugate wherein every polymer molecule in the conjugate has a molecular weight that is at or about 20,000 D to at or about 300,000 D, or is at or about 30,000 D to at or about 300,000 D, or is at or about 40,000 D to at or about 300,000 D, and wherein the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

In still another aspect, the invention encompasses a conjugate wherein every polymer molecule in the conjugate has a molecular weight that is at or about 20,000 D to at or about 100,000 D, or is at or about 30,000 D to at or about 100,000 D, or is at or about 40,000 D to at or about 100,000 D, and wherein the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

In still another aspect, the invention encompasses a conjugate wherein every polymer molecule in the conjugate has a molecular weight that is at or about 20,000 D to at or about 70,000 D, or is at or about 30,000 D to at or about 70,000 D, or is at or about 40,000 D to at or about 70,000 D, and wherein the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

In still another aspect, the invention encompasses a conjugate wherein every polymer molecule in the

conjugate has a molecular weight that is at or about 20,000 D to at or about 50,000 D, or is at or about 30,000 D to at or about 50,000 D, or is at or about 40,000 D to at or about 50,000 D, and wherein the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

In still another aspect, the invention encompasses a conjugate wherein every polymer molecule in the conjugate has a molecular weight that is at or about 20,000 D to at or about 40,000 D, or is at or about 30,000 D to at or about 40,000 D, and wherein the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

It is believed that the serum half-life, MRT and/or serum clearance rate of any antibody fragment can be greatly improved by derivatizing the antibody fragment with polymer as taught herein. In one embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH, Fv, scFv and F(ab')₂.

In a preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

In yet another preferred embodiment, the conjugate contains a F(ab')₂ antibody fragment attached to no more than about 2 polymer molecules, wherein every polymer molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In a further embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule and the polymer is coupled to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In an additional embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate is at least about 20,000 D in molecular weight, or at least about 30,000 in molecular weight, or at least about 40,000 D in molecular weight, every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment, and the

conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

5 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment, and the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

10 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment, and the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

20 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment, and the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

25 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment, and the conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

30 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, every polymer molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, every polymer molecule in the conjugate is attached to the hinge region of the antibody fragment, and the

conjugate contains no more than about 10 polymer molecules, or no more than about 5 polymer molecules, or no more than about 4 polymer molecules, or no more than about 3 polymer molecules, or no more than about 2 polymer molecules, or no more than 1 polymer molecule.

In a further embodiment, the conjugate contains a $F(ab')_2$ antibody fragment attached to no more than about 2 polymer molecules, wherein every polymer molecule in the conjugate is at least about 20,000 D in molecular weight, or at least about 30,000 D in molecular weight, or at least about 40,000 D in molecular weight, and wherein every polymer molecule in the conjugate is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains a $F(ab')_2$ antibody fragment attached to no more than about 2 polymer molecules, wherein every polymer molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, and wherein every polymer molecule in the conjugate is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains a $F(ab')_2$ antibody fragment attached to no more than about 2 polymer molecules, wherein every polymer molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, and wherein every polymer molecule in the conjugate is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains a $F(ab')_2$ antibody fragment attached to no more than about 2 polymer molecules, wherein every polymer molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, and wherein every polymer molecule in the conjugate is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains a $F(ab')_2$ antibody fragment attached to no more than about 2 polymer molecules, wherein every polymer molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, and wherein every polymer molecule in

the conjugate is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains a $F(ab')_2$ antibody fragment attached to no more than
5 about 2 polymer molecules, wherein every polymer molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, and wherein every polymer molecule in the conjugate is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the
10 corresponding cysteine residue in the opposite chain.

In yet another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at least about 20,000 D in molecular weight, or at least about 30,000 D in molecular weight, or at least about 40,000 D in molecular weight, wherein the polymer molecule
15 is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, wherein the polymer molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine,
25 for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, wherein the polymer molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine,
30 for the corresponding cysteine residue in the opposite chain.

In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, wherein the polymer molecule is attached to a cysteine residue in the light or
35

heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

5 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, wherein the polymer molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

10 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, wherein the polymer molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

15 In still another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at least about 20,000 D in molecular weight, or at least about 30,000 D in molecular weight, or at least about 40,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

20 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

25 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

30 In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

70,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

In another embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 polymer molecule, wherein the polymer molecule is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, and wherein the polymer molecule is attached to the hinge region of the antibody fragment.

Although any type of polymer is contemplated for use in constructing the conjugates of the invention, including the polymers and chemical linkage systems described in Section (II)(1)(b) below, polyethylene glycol (PEG) polymers are preferred for use herein.

In one embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW of at least about 20,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW of at least about 30,000 D.

In yet another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW of at least about 40,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW that is at or about 20,000 D to at or about 300,000 D, or is at or about 30,000 D to at or about 300,000 D, or is at or about 40,000 D to at or about 300,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW that is at or about 20,000 D to at or about 100,000 D, or is at or about 30,000 D to at or about 100,000 D, or is at or about 40,000 D to at or about 100,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW that is at or about 20,000 D to at or about 70,000 D, or is at or about 30,000 D to at or about 70,000 D, or is at or about 40,000 D to at or about 70,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW that is at or about 20,000 D to at or about 50,000 D, or is at or about 30,000 D to at or about 50,000 D, or is at or about 40,000 D to at or about 50,000 D.

In another embodiment, the conjugate is an antibody fragment covalently attached to at least one PEG having an actual MW that is at or about 20,000 D to at or about 40,000 D, or is at or about 30,000 D to at or about 40,000 D.

In another aspect, the invention encompasses a conjugate with a PEG to antibody fragment molar ratio of no more than about 10:1, or no more than about 5:1, or no more than about 4:1, or no more than about 3:1,

or no more than about 2:1, or no more than 1:1.

In yet another aspect, the invention encompasses a conjugate wherein the antibody fragment is attached to about 10 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In another embodiment, the conjugate contains an antibody fragment attached to about 5 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In still another embodiment, the conjugate contains an antibody fragment attached to about 4 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In a further embodiment, the conjugate contains an antibody fragment attached to about 3 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In an additional embodiment, the conjugate contains an antibody fragment attached to about 2 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. Also provided herein is a conjugate containing an antibody fragment attached to a single PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D.

In another aspect, the invention encompasses a conjugate wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In still another aspect, the invention encompasses a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH and F(ab')₂, wherein the antibody fragment is attached to about 10 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In another embodiment, the foregoing conjugate contains an antibody fragment attached to about 5 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In still another embodiment, the foregoing conjugate contains an antibody fragment attached to about 4 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In a further embodiment, the foregoing conjugate contains an antibody fragment attached to about 3 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. In an additional embodiment, the foregoing conjugate contains an antibody fragment attached to about 2 or fewer PEG molecules, each PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. Also provided herein is the foregoing conjugate that contains an antibody fragment attached to a single PEG molecule having a molecular weight of at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D.

In another aspect, the invention encompasses a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH and F(ab')₂, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH and F(ab')₂, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate containing an antibody fragment selected

from the group consisting of Fab, Fab', Fab'-SH and F(ab')₂, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH and F(ab')₂, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another aspect, the invention encompasses a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', Fab'-SH and F(ab')₂, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In a preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG having a molecular weight of at least about 20,000D, or at least about 30,000D, or at least about 40,000D, and wherein every PEG molecule in the conjugate is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG having a molecular weight that is at or about 20,000 D to about 300,000 D, or is at or about 30,000 D to at or about 300,000 D, or is at or about 40,000 D to at or about 300,000 D, and wherein every PEG molecule in the conjugate is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG having a molecular weight that is at or about 20,000 D to about 100,000 D, or is at or about 30,000 D to at or about 100,000 D, or is at or about 40,000 D to at or about 100,000 D, and wherein every PEG molecule in the conjugate is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG having a molecular weight that is at or about 20,000 D to about 70,000 D, or is at or about 30,000 D to at or about 70,000 D, or is at or about 40,000 D to at or about 70,000 D, and wherein every PEG molecule in the conjugate

is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG having a molecular weight that is at or about 20,000 D to about 50,000 D, or is at or about 30,000 D to at or about 50,000 D, or is at or about 40,000 D to at or about 50,000 D, and wherein every PEG molecule in the conjugate is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG having a molecular weight that is at or about 20,000 D to about 40,000 D, or is at or about 30,000 D to at or about 40,000 D, and wherein every PEG molecule in the conjugate is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at least about 20,000D in molecular weight, or at least about 30,000D in molecular weight, or at least about 40,000D in molecular weight, wherein every PEG molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, wherein every PEG molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, wherein every PEG molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about

70,000 D in molecular weight, wherein every PEG molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

5 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, wherein every PEG molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

10 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, wherein every PEG molecule in the conjugate molecule is attached to the hinge region of the antibody fragment, and wherein the conjugate contains no more than about 10 PEG molecules, or no more than about 5 PEG molecules, or no more than about 4 PEG molecules, or no more than about 3 PEG molecules, or no more than about 2 PEG molecules, or no more than 1 PEG molecule.

20 In yet another preferred embodiment, the conjugate contains a $F(ab')_2$ antibody fragment derivatized with PEG, wherein every PEG molecule in the conjugate is at least about 20,000D in molecular weight, or at least about 30,000D in molecular weight, or at least about 40,000D in molecular weight, wherein the antibody fragment is attached to no more than about 2 PEG molecules, and wherein every PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

25 In another preferred embodiment, the conjugate contains a $F(ab')_2$ antibody fragment derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, wherein the antibody fragment is attached to no more than about 2 PEG molecules, and wherein every PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

30 In another preferred embodiment, the conjugate contains a $F(ab')_2$ antibody fragment derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about

40,000 D to at or about 100,000 D in molecular weight, wherein the antibody fragment is attached to no more than about 2 PEG molecules, and wherein every PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains a $F(ab')_2$ antibody fragment derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, wherein the antibody fragment is attached to no more than about 2 PEG molecules, and wherein every PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains a $F(ab')_2$ antibody fragment derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, wherein the antibody fragment is attached to no more than about 2 PEG molecules, and wherein every PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains a $F(ab')_2$ antibody fragment derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, wherein the antibody fragment is attached to no more than about 2 PEG molecules, and wherein every PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In still another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at least about 20,000 D in molecular weight, or at least about 30,000 in molecular weight, or at least about 40,000 D in molecular weight, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein

every PEG molecule in the conjugate is at or about 20,000 D to at or about 300,000 D in molecular weight, or is at or about 30,000 D to at or about 300,000 D in molecular weight, or is at or about 40,000 D to at or about 300,000 D in molecular weight, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 100,000 D in molecular weight, or is at or about 30,000 D to at or about 100,000 D in molecular weight, or is at or about 40,000 D to at or about 100,000 D in molecular weight, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 70,000 D in molecular weight, or is at or about 30,000 D to at or about 70,000 D in molecular weight, or is at or about 40,000 D to at or about 70,000 D in molecular weight, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 50,000 D in molecular weight, or is at or about 30,000 D to at or about 50,000 D in molecular weight, or is at or about 40,000 D to at or about 50,000 D in molecular weight, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is attached to a cysteine residue in the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is derivatized with PEG, wherein every PEG molecule in the conjugate is at or about 20,000 D to at or about 40,000 D in molecular weight, or is at or about 30,000 D to at or about 40,000 D in molecular weight, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is attached to a cysteine residue in the light or

heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains, wherein the disulfide bridge is avoided by substituting another amino acid, such as serine, for the corresponding cysteine residue in the opposite chain.

5 It will be appreciated that all of the above-described embodiments of the invention utilizing PEG polymers include conjugates wherein the PEG polymer(s) is (are) linear or branched. In a preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and at least about 40,000 D in molecular weight. In a particularly surprising and unexpected finding, the inventors discovered that the foregoing conjugate exhibits a serum half-life, MRT and serum clearance rate approaching that of full length antibody as shown in Example X below.

10 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 300,000 D.

15 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 100,000 D.

20 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 70,000 D.

25 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 50,000 D.

30 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and at least 40,000D in molecular weight, and the PEG molecule is attached to the hinge region of the antibody fragment.

35 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 300,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 100,000 D, and the PEG molecule is attached to the hinge region of the antibody

fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 70,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 40,000 D to at or about 50,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In a preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and at least about 40,000 D in molecular weight.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 300,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 100,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 70,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 50,000 D.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and at least 40,000D in molecular weight, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 300,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 100,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 70,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 40,000 D to at or about 50,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at least about 30,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 300,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 100,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 70,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 50,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 40,000 D.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment

selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and at least 30,000D in molecular weight, and the PEG molecule is attached to the hinge region of the antibody fragment.

5 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 300,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

10 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 100,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

15 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 70,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

20 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 50,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

25 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 30,000 D to at or about 40,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

30 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at least about 30,000 D.

35 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 300,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D

to at or about 100,000 D.

5 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 70,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 50,000 D.

10 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 40,000 D.

15 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and at least 30,000D in molecular weight, and the PEG molecule is attached to the hinge region of the antibody fragment.

20 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 300,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

25 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 100,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

30 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 70,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

35 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 50,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no

more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 30,000 D to at or about 40,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

5 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at least about 20,000 D.

10 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 300,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 100,000 D.

15 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 70,000 D.

20 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 50,000 D.

25 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 40,000 D.

30 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 30,000 D.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and at least 20,000D in molecular weight, and the PEG molecule is attached to the hinge region of the antibody fragment.

35 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 300,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 100,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 70,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 50,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 40,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is linear and has a molecular weight that is at or about 20,000 D to at or about 30,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at least about 20,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 300,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 100,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG

molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 70,000 D.

5 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 50,000 D.

10 In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 40,000 D.

In another preferred embodiment, the conjugate contains an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, and wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 30,000 D.

15 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and at least 20,000D in molecular weight, and the PEG molecule is attached to the hinge region of the antibody fragment.

20 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 300,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

25 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 100,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

30 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 70,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

35 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 50,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

In another preferred embodiment, the invention provides a conjugate containing an antibody fragment

selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 40,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

5 In another preferred embodiment, the invention provides a conjugate containing an antibody fragment selected from the group consisting of Fab, Fab', and Fab'-SH, wherein the antibody fragment is attached to no more than 1 PEG molecule, wherein the PEG molecule is branched and has a molecular weight that is at or about 20,000 D to at or about 30,000 D, and the PEG molecule is attached to the hinge region of the antibody fragment.

10 In one aspect, the invention provides any of the above-described conjugates wherein the conjugate contains no more than one antibody fragment. Additionally provided herein is any of the above-described conjugates wherein the conjugate contains one or more antibody fragment(s) covalently linked to one or more polymer molecule(s), such as conjugates containing two or more antibody fragments covalently linked together by polymer molecule(s). In one embodiment, a polymer molecule is used to link together two antibody
15 fragments to form a dumbbell-shaped structure. Also encompassed herein are conjugates formed by more than two antibody fragments joined by polymer molecule(s) to form a rosette or other shapes. The antibody fragments in such structures can be of the same or different fragment type and can have the same antigen specificity or have different antigen specificities. Such structures can be made by using a polymer molecule derivatized with multiple functional groups permitting the direct attachment, or the attachment by means of bi-
20 or multi-functional linkers, of two or more antibody fragments to the polymer backbone.

In another aspect, the invention encompasses any of the above-described conjugates utilizing an antibody fragment comprising an antigen recognition site that binds to rabbit IL-8 and/or human IL-8. In yet another aspect, the invention encompasses any of the above-described conjugates utilizing an antibody fragment comprising 6G4.2.5LV/L1N35A or 6G4.2.5LV/L1N35E as defined below. In still another aspect,
25 the invention encompasses any of the above-described conjugates utilizing an antibody fragment comprising 6G4.5.2.SHV11 as defined below. In a further aspect, the invention encompasses any of the above-described conjugates utilizing an antibody fragment comprising hu6G4.2.5LV/L1N35A or hu6G4.2.5LV/L1N35E as defined below. In an additional aspect, the invention encompasses any of the above-described conjugates utilizing an antibody fragment comprising hu6G4.2.5HV. Further encompassed herein are any of the above-
30 described conjugates utilizing an antibody fragment comprising 6G4.2.5LV/L1N35A or 6G4.2.5LV/L1N35E and further comprising the CDRs of 6G4.2.5HV as defined below. Also encompassed herein are any of the above described conjugates utilizing an antibody fragment comprising hu6G4.2.5LV/L1N35A or hu6G4.2.5LV/L1N35E and further comprising hu6G4.2.5HV as defined below. Additionally encompassed herein are any of the above-described conjugates utilizing an antibody fragment comprising
35 6G4.2.5LV11N35A or 6G4.2.5LV11N35E as defined below. Further provided herein are any of the above-described conjugates utilizing an antibody fragment comprising 6G4.2.5LV11N35A or 6G4.2.5LV11N35E and further comprising 6G4.2.5HV11 as defined below.

a. Production of Antibody Fragments

Antibody fragments can be produced by any method known in the art. Generally, an antibody

fragment is derived from a parental intact antibody. The parental antibody can be generated by raising polyclonal sera against the desired antigen by multiple subcutaneous (sc) or intraperitoneal (ip) injections of antigen and an adjuvant, such as monophosphoryl lipid A (MPL)/trehalose dicrynomycolate (TDM) (Ribi Immunochem. Research, Inc., Hamilton, MT), at multiple sites. Two weeks later the animals are boosted. 7 to 14 days later animals are bled and the serum is assayed for anti-antigen titer. Animals are boosted until titer plateaus. Sera are harvested from animals, and polyclonal antibodies are isolated from sera by conventional immunoglobulin purification procedures, such as protein A-Sepharose chromatography, hydroxylapatite chromatography, gel filtration, dialysis, or antigen affinity chromatography. The desired antibody fragments can be generated from purified polyclonal antibody preparations by conventional enzymatic methods, e.g. F(ab')₂ fragments are produced by pepsin cleavage of intact antibody, and Fab fragments are produced by briefly digesting intact antibody with papain.

Alternatively, antibody fragments are derived from monoclonal antibodies generated against the desired antigen. Monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster or macaque monkey, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOP-21 and M.C.-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of antibody-encoding DNA include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5: 256 (1993) and Pluckthun, *Immunol. Revs.*, 130: 151 (1992).

In a preferred embodiment, the antibody fragment is derived from a humanized antibody. Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. It will be appreciated that variable domain sequences obtained from any non-human animal phage display library-derived Fv clone or from any non-human animal hybridoma-derived antibody clone provided as described herein can serve as the "import" variable domain used in the construction of the humanized antibodies of the invention. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, 321: 522 (1986); Riechmann *et al.*, *Nature*, 332: 323 (1988); Verhoeven *et al.*, *Science*, 239: 1534 (1988)), by substituting non-human animal, *e.g.* rodent, CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (Cabilly *et al.*, *supra*), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in non-human animal, *e.g.* rodent, antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a non-human animal, *e.g.* rodent, antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the non-human animal is then accepted as the human framework (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*,

151: 2296 (1993); Chothia and Lesk, *J. Mol. Biol.*, 196: 901 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup light or heavy chains. The same framework can be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci USA*, 89: 4285 (1992); Presta *et al.*, *J. Immunol.*, 151: 2623 (1993)).

5 It is also important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer
10 programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind to its antigen. In this way, FR residues can be selected and combined from the consensus and import sequences so that the desired antibody characteristic,
15 such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

In addition, antibody fragments for use herein can be derived from human monoclonal antibodies. Human monoclonal antibodies against the antigen of interest can be made by the hybridoma method. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies
20 have been described, for example, by Kozbor *J. Immunol.*, 133: 3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner *et al.*, *J. Immunol.*, 147: 86 (1991).

It is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production.
25 For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits *et al.*, *Proc. Natl. Acad. Sci USA*, 90: 2551 (1993); Jakobovits *et al.*, *Nature*, 362: 255 (1993); Bruggermann *et al.*, *Year in*
30 *Immunol.*, 7: 33 (1993).

Alternatively, phage display technology (McCafferty *et al.*, *Nature* 348:552 (1990)) can be used to produce human antibodies and antibody fragments in vitro, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and
35 displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties.

Thus, the phage mimics some of the properties of the B-cell. Phage display can be performed in a variety of formats; for their review see, e.g., Johnson *et al.*, *Current Opinion in Structural Biology* 3:564 (1993). Several

sources of V-gene segments can be used for phage display. Clackson et al., Nature 352:624 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks et al., J. Mol. Biol. 222:581 (1991), or Griffith et al., EMBO J. 12:725 (1993). In a natural immune response, antibody genes accumulate mutations at a high rate (somatic hypermutation). Some of the changes introduced will confer higher affinity, and B cells displaying high-affinity surface immunoglobulin are preferentially replicated and differentiated during subsequent antigen challenge. This natural process can be mimicked by employing the technique known as "chain shuffling" (Marks et al., Bio/Technol. 10:779 (1992)). In this method, the affinity of "primary" human antibodies obtained by phage display can be improved by sequentially replacing the heavy and light chain V region genes with repertoires of naturally occurring variants (repertoires) of V domain genes obtained from unimmunized donors. This technique allows the production of antibodies and antibody fragments with affinities in the nM range. A strategy for making very large phage antibody repertoires has been described by Waterhouse et al., Nucl. Acids Res. 21:2265 (1993).

Gene shuffling can also be used to derive human antibodies from non-human, e.g. rodent, antibodies, where the human antibody has similar affinities and specificities to the starting non-human antibody. According to this method, which is also called "epitope imprinting", either the heavy or light chain variable region of a non-human antibody fragment obtained by phage display techniques as described above is replaced with a repertoire of human V domain genes, creating a population of non-human chain/human chain scFv or Fab chimeras. Selection with antigen results in isolation of a non-human chain/human chain chimeric scFv or Fab wherein the human chain restores the antigen binding site destroyed upon removal of the corresponding non-human chain in the primary phage display clone, i.e. the epitope governs (imprints) the choice of the human chain partner. When the process is repeated in order to replace the remaining non-human chain, a human antibody is obtained (see PCT WO 93/06213 published April 1, 1993). Unlike traditional humanization of non-human antibodies by CDR grafting, this technique provides completely human antibodies, which have no FR or CDR residues of non-human origin.

The invention also encompasses the use of bispecific and heteroconjugate antibody fragments having specificities for at least two different antigens. Bispecific and heteroconjugate antibodies can be prepared as full length antibodies or as antibody fragments (e.g. $F(ab')_2$ bispecific antibody fragments). Antibody fragments having more than two valencies (e.g. trivalent or higher valency antibody fragments) are also contemplated for use herein. Bispecific antibodies, heteroconjugate antibodies, and multi-valent antibodies can be prepared as described in Section (II)(3)(C) below.

As described above, DNA encoding the monoclonal antibody or antibody fragment of interest can be isolated from its hybridoma or phage display clone of origin, and then manipulated to create humanized and/or affinity matured constructs. In addition, known techniques can be employed to introduce an amino acid residue or residues into any desired location on the polypeptide backbone of the antibody fragment, e.g. a cysteine residue placed in the hinge region of the heavy chain, thereby providing a site for specific attachment

of polymer molecule(s). In one embodiment, the native cysteine residue in either the light or heavy chain of the antibody fragment that would ordinarily form the disulfide bridge linking the light and heavy chains is substituted with another amino acid, such as serine, in order to leave the partner cysteine residue in the opposite chain with a free sulfhydryl for specific attachment of polymer molecule.

5 Upon construction of the desired antibody or antibody fragment-encoding clone, the clone can be used for recombinant production of the antibody fragment as described in Section (II)(4) below. Finally, the antibody or antibody fragment product can be recovered from host cell culture and purified as described in Section (II)(4)(F) below. In the case of embodiments utilizing an antibody fragment engineered to lack a cysteine residue that ordinarily forms the disulfide bridge between the light and heavy chains as described
10 above, preferred recombinant production systems include bacterial expression and product recovery procedures utilizing the low pH osmotic shock method described in the "Alternative Fab'-SH Purification" section of Example T below. If a full length antibody is produced, the desired antibody fragment can be obtained therefrom by subjecting the intact antibody to enzymatic digestion according to known methods, e.g. as described in Section (II)(4)(G) below.

15 **b. Construction of Antibody Fragment-Polymer Conjugates**

The antibody fragment-polymer conjugates of the invention can be made by derivatizing the desired antibody fragment with an inert polymer. It will be appreciated that any inert polymer which provides the conjugate with the desired apparent size or which has the selected actual MW as taught herein is suitable for use in constructing the antibody fragment-polymer conjugates of the invention.

20 Many inert polymers are suitable for use in pharmaceuticals. See, e.g., Davis et al., Biomedical Polymers: Polymeric Materials and Pharmaceuticals for Biomedical Use, pp.441-451 (1980). In all embodiments of the invention, a non-proteinaceous polymer is used. The nonproteinaceous polymer ordinarily is a hydrophilic synthetic polymer, i.e., a polymer not otherwise found in nature. However, polymers which exist in nature and are produced by recombinant or *in vitro* methods are also useful, as are polymers which are
25 isolated from native sources. Hydrophilic polyvinyl polymers fall within the scope of this invention, e.g. polyvinylalcohol and polyvinylpyrrolidone. Particularly useful are polyalkylene ethers such as polyethylene glycol (PEG); polyoxyalkylenes such as polyoxyethylene, polyoxypropylene, and block copolymers of polyoxyethylene and polyoxypropylene (Pluronics); polymethacrylates; carbomers; branched or unbranched polysaccharides which comprise the saccharide monomers D-mannose, D- and L-galactose, fucose, fructose,
30 D-xylose, L-arabinose, D-glucuronic acid, sialic acid, D-galacturonic acid, D-mannuronic acid (e.g. polymannuronic acid, or alginic acid), D-glucosamine, D-galactosamine, D-glucose and neuraminic acid including homopolysaccharides and heteropolysaccharides such as lactose, amylopectin, starch, hydroxyethyl starch, amylose, dextrane sulfate, dextran, dextrans, glycogen, or the polysaccharide subunit of acid mucopolysaccharides, e.g. hyaluronic acid; polymers of sugar alcohols such as polysorbitol and polymannitol;
35 heparin or heparan. The polymer prior to cross-linking need not be, but preferably is, water soluble, but the final conjugate must be water-soluble. Preferably, the conjugate exhibits a water solubility of at least about 0.01 mg/ml, and more preferably at least about 0.1 mg/ml, and still more preferably at least about 1 mg/ml. In addition, the polymer should not be highly immunogenic in the conjugate form, nor should it possess viscosity that is incompatible with intravenous infusion or injection if the conjugate is intended to be administered by

such routes.

In one embodiment, the polymer contains only a single group which is reactive. This helps to avoid cross-linking of protein molecules. However, it is within the scope herein to maximize reaction conditions to reduce cross-linking, or to purify the reaction products through gel filtration or ion exchange chromatography to recover substantially homogenous derivatives. In other embodiments, the polymer contains two or more reactive groups for the purpose of linking multiple antibody fragments to the polymer backbone. Again, gel filtration or ion exchange chromatography can be used to recover the desired derivative in substantially homogeneous form.

The molecular weight of the polymer can range up to about 500,000 D; and preferably is at least about 20,000 D, or at least about 30,000 D, or at least about 40,000 D. The molecular weight chosen can depend upon the effective size of the conjugate to be achieved, the nature (e.g. structure, such as linear or branched) of the polymer, and the degree of derivatization, i.e. the number of polymer molecules per antibody fragment, and the polymer attachment site or sites on the antibody fragment.

The polymer can be covalently linked to the antibody fragment through a multifunctional crosslinking agent which reacts with the polymer and one or more amino acid residues of the antibody fragment to be linked. However, it is also within the scope of the invention to directly crosslink the polymer by reacting a derivatized polymer with the antibody fragment, or vice versa.

The covalent crosslinking site on the antibody fragment includes the N-terminal amino group and epsilon amino groups found on lysine residues, as well as other amino, imino, carboxyl, sulfhydryl, hydroxyl or other hydrophilic groups. The polymer may be covalently bonded directly to the antibody fragment without the use of a multifunctional (ordinarily bifunctional) crosslinking agent. Covalent binding to amino groups is accomplished by known chemistries based upon cyanuric chloride, carbonyl diimidazole, aldehyde reactive groups (PEG alkoxide plus diethyl acetal of bromoacetaldehyde; PEG plus DMSO and acetic anhydride, or PEG chloride plus the phenoxide of 4-hydroxybenzaldehyde, activated succinimidyl esters, activated dithiocarbonate PEG, 2,4,5-trichlorophenylchloroformate or P-nitrophenylchloroformate activated PEG.) Carboxyl groups are derivatized by coupling PEG-amine using carbodiimide. Sulfhydryl groups are derivatized by coupling to maleimido-substituted PEG (e.g. alkoxy-PEG amine plus sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate) as described in WO 97/10847 published March 27, 1997, or PEG-maleimide commercially available from Shearwater Polymers, Inc., Huntsville, AL). Alternatively, free amino groups on the antibody fragment (e.g. epsilon amino groups on lysine residues) can be thiolated with 2-imino-thiolane (Traut's reagent) and then coupled to maleimide-containing derivatives of PEG as described in Pedley et al., *Br. J. Cancer*, 70: 1126-1130 (1994).

The polymer will bear a group which is directly reactive with an amino acid side chain, or the N- or C-terminus of the polypeptide linked, or which is reactive with the multifunctional cross-linking agent. In general, polymers bearing such reactive groups are known for the preparation of immobilized proteins. In order to use such chemistries here, one should employ a water soluble polymer otherwise derivatized in the same fashion as insoluble polymers heretofore employed for protein immobilization. Cyanogen bromide activation is a particularly useful procedure to employ in crosslinking polysaccharides.

"Water soluble" in reference to the starting polymer means that the polymer or its reactive

intermediate used for conjugation is sufficiently water soluble to participate in a derivatization reaction.

The degree of substitution with such a polymer will vary depending upon the number of reactive sites on the antibody fragment, the molecular weight, hydrophilicity and other characteristics of the polymer, and the particular antibody fragment derivatization sites chosen. In general, the conjugate contains from 1 to about 10 polymer molecules, but greater numbers of polymer molecules attached to the antibody fragments of the invention are also contemplated. The desired amount of derivatization is easily achieved by using an experimental matrix in which the time, temperature and other reaction conditions are varied to change the degree of substitution, after which the level of polymer substitution of the conjugates is determined by size exclusion chromatography or other means known in the art.

The polymer, e.g. PEG, is cross-linked to the antibody fragment by a wide variety of methods known *per se* for the covalent modification of proteins with nonproteinaceous polymers such as PEG. Certain of these methods, however, are not preferred for the purposes herein. Cyanuronic chloride chemistry leads to many side reactions, including protein cross-linking. In addition, it may be particularly likely to lead to inactivation of proteins containing sulfhydryl groups. Carbonyl diimidazole chemistry (Beauchamp *et al.*, Anal Biochem. **131**, 25-33 [1983]) requires high pH (>8.5), which can inactivate proteins. Moreover, since the "activated PEG" intermediate can react with water, a very large molar excess of "activated PEG" over protein is required. The high concentrations of PEG required for the carbonyl diimidazole chemistry also led to problems in purification, as both gel filtration chromatography and hydrophilic interaction chromatography are adversely affected. In addition, the high concentrations of "activated PEG" may precipitate protein, a problem that *per se* has been noted previously (Davis, U.S. Patent No. 4,179,337). On the other hand, aldehyde chemistry (Royer, U.S. Patent No. 4,002,531) is more efficient since it requires only a 40-fold molar excess of PEG and a 1-2 hr incubation. However, the manganese dioxide suggested by Royer for preparation of the PEG aldehyde is problematic "because of the pronounced tendency of PEG to form complexes with metal-based oxidizing agents" (Harris *et al.*, J. Polym. Sci. Polym. Chem. Ed. **22**, 341-52 [1984]). The use of a Moffatt oxidation, utilizing DMSO and acetic anhydride, obviates this problem. In addition, the sodium borohydride suggested by Royer must be used at high pH and has a significant tendency to reduce disulfide bonds. In contrast, sodium cyanoborohydride, which is effective at neutral pH and has very little tendency to reduce disulfide bonds is preferred. In another preferred embodiment, maleimido-activated PEG is used for coupling to free thiols on the antibody fragment.

Functionalized PEG polymers to modify the antibody fragments of the invention are available from Shearwater Polymers, Inc. (Huntsville, AL). Such commercially available PEG derivatives include, but are not limited to, amino-PEG, PEG amino acid esters, PEG-hydrazide, PEG-thiol, PEG-succinate, carboxymethylated PEG, PEG-propionic acid, PEG amino acids, PEG succinimidyl succinate, PEG succinimidyl propionate, succinimidyl ester of carboxymethylated PEG, succinimidyl carbonate of PEG, succinimidyl esters of amino acid PEGs, PEG-oxycarbonylimidazole, PEG-nitrophenyl carbonate, PEG tresylate, PEG-glycidyl ether, PEG-aldehyde, PEG-vinylsulfone, PEG-maleimide, PEG-orthopyridyl-disulfide, heterofunctional PEGs, PEG vinyl derivatives, PEG silanes, and PEG phospholides. The reaction conditions for coupling these PEG derivatives will vary depending on the protein, the desired degree of PEGylation, and the PEG derivative utilized. Some factors involved in the choice of PEG derivatives include: the desired point of attachment (such as lysine or

cysteine R-groups), hydrolytic stability and reactivity of the derivatives, stability, toxicity and antigenicity of the linkage, suitability for analysis, etc. Specific instructions for the use of any particular derivative are available from the manufacturer.

The conjugates of this invention are separated from the unreacted starting materials by gel filtration or ion exchange HPLC. Heterologous species of the conjugates are purified from one another in the same fashion.

The conjugates may also be purified by ion-exchange chromatography. The chemistry of many of the electrophilically activated PEG's results in a reduction of amino group charge of the PEGylated product. Thus, high resolution ion exchange chromatography can be used to separate the free and conjugated proteins, and to resolve species with different levels of PEGylation. In fact, the resolution of different species (e.g. containing one or two PEG residues) is also possible due to the difference in the ionic properties of the unreacted amino acids. In one embodiment, species with difference levels of PEGylation are resolved according to the methods described in WO 96/34015 (International Application No. PCT/US96/05550 published October 31, 1996).

In a preferred embodiment, the conjugate is generated by utilizing the derivatization and purification methods described in Section (T) of the Examples below.

In one aspect, the invention provides any of the above-described conjugates formed by its component parts, i.e. one or more antibody fragment(s) covalently attached to one or more polymer molecule(s), without any extraneous matter in the covalent molecular structure of the conjugate.

c. Other Derivatives of Large Effective Size Conjugates

In another aspect, any of the above-described conjugates can be modified to contain one or more component(s) in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate, namely, the substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived. In one embodiment, the invention provides any of the above-described conjugates modified to incorporate one or more nonproteinaceous functional group(s). For example, the conjugate can be modified to incorporate nonproteinaceous labels or reporter molecules, such as radiolabels, including any radioactive substance used in medical treatment or imaging or used as an effector function or tracer in an animal model, such as radioisotopic labels ^{99}Tc , ^{90}Y , ^{111}In , ^{32}P , ^{14}C , ^{125}I , ^3H , ^{131}I , ^{11}C , ^{15}O , ^{13}N , ^{18}F , ^{35}S , ^{51}Cr , ^{57}Co , ^{226}Ra , ^{60}Co , ^{59}Fe , ^{75}Se , ^{152}Eu , ^{67}Cu , ^{217}Bi , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , ^{234}Th , ^{40}K , and the like, non-radioisotopic labels such as ^{157}Gd , ^{55}Mn , ^{52}Tr , ^{56}Fe , etc., fluorescent or chemiluminescent labels, including fluorophores such as rare earth chelates, fluorescein and its derivatives, rhodamine and its derivatives, isothiocyanate, phycoerythrin, phycocyanin, allophycocyanin, o-phthalaldehyde, fluorescamine, ^{152}Eu , dansyl, umbelliferone, luciferin, luminal label, isoluminal label, an aromatic acridinium ester label, an imidazole label, an acridinium salt label, an oxalate ester label, an aequorin label, 2,3-dihydrophthalazinediones, biotin/avidin, spin labels, stable free radicals, and the like.

Conventional methods are available to bind these labels covalently to the polypeptide antibody fragment or polymer component of the conjugate. In one aspect, any conjugate of the invention is modified by

derivatizing the antibody fragment component with any of the above-described non-proteinaceous labels, wherein the label is directly or indirectly (through a coupling agent) attached to the antibody fragment, and wherein such derivatization of the antibody fragment does not contribute or introduce any polymer moiety into the molecular structure of the conjugate. For instance, coupling agents such as dialdehydes, carbodiimides, dimaleimides, bis-imidates, bis-diazotized benzidine, and the like can be used to tag the antibody fragment with the above-described fluorescent or chemiluminescent labels. See, for example, U.S. Pat. No. 3,940,475 (fluorimetry), Morrison, Meth. Enzymol., **32b**, 103 (1974), Svyanen et al., J. Biol. Chem., **284**, 3762 (1973), and Bolton and Hunter, Biochem. J., **133**, 529 (1973).

In the case of embodiments utilizing radiolabels, both direct and indirect labeling can be used to incorporate the selected radionuclide into the conjugate. As used herein in the context of radiolabeling, the phrases "indirect labeling" and "indirect labeling approach" both mean that a chelating agent is covalently attached to the antibody fragment moiety or polymer moiety of the conjugate and at least one radionuclide is inserted into the chelating agent. Preferred chelating agents and radionuclides are set forth in Srivastava, S.C. and Mease, R.C., "Progress in Research on Ligands, Nuclides and Techniques for Labeling Monoclonal Antibodies," Nucl. Med. Bio., **18(6)**: 589-603 (1991). A particularly preferred chelating agent is 1-isothiocyanatobenzyl-3-methyldiethylenetriaminepentaacetic acid ("MX-DTPA"). As used herein in the context of radiolabeling, the phrases "direct labeling" and "direct labeling approach" both mean that a radionuclide is covalently attached directly to the antibody fragment moiety (typically via an amino acid residue) or to the polymer moiety of the conjugate. Preferred radionuclides for use in direct labeling of conjugate are provided in Srivastava and Mease, *supra*. In one embodiment, the conjugate is directly labeled with ¹³¹I covalently attached to tyrosine residues. In another embodiment, the antibody fragment component of the conjugate is directly or indirectly labeled with any of the above-described radiolabels, wherein such labeling of the antibody fragment does not contribute or introduce any polymer moiety into the molecular structure of the conjugate.

d. Therapeutic Compositions and Administration of Large Effective Size Conjugates

The conjugate of the invention is useful for treating the disease indications that are treated with the parent intact antibody. For example, a conjugate derived from an anti-IL-8 antibody or fragment is useful in the treatment of inflammatory disorders as described in Section (II)(5)(B) below. Such conjugates have prophylactic and therapeutic applications in a broad spectrum of IL-8 mediated diseases, such as inflammatory diseases and asthma, in a manner similar to the widespread efficacy of anti-IL-8 antibodies in the treatment of such disease indications that is known in the art, which treatment indications include: (1) ischemic reperfusion injury of the lung (Sekido et al., Nature, **365**: 654 (1993)); (2) acute lung injury and ARDS (WO 96/22785 published August 1, 1996; Folkesson et al., J. Clin. Invest., **96**: 107-116 (1995); Mulligan et al., J. Immunol., **150**: 5585-5595 (1993)); (3) hypovolemic shock (Hebert, C., "Humanized Anti-IL-8: Potential Therapy for Shock and ARDS", seminar presented at Keystone Conference on The Role of Cytokines in Leukocyte Trafficking and Disease, held at Copper Mountain Resort, CO, March 31-April 5, 1997; Sharar, S.A., Harlan, J.H., Patterson, C.A., Hebert, C.A., and Winn, R.K., "Reperfusion Injury After Hemorrhagic Shock in Rabbits is Reduced Similarly by IL-8 or CD-18 Monoclonal Antibodies", manuscript submitted 1998); (4) myocardial

infarction (WO 97/40215 published October 30, 1997); (5) cerebral reperfusion injury (Matsumoto et al., Laboratory Invest., 77: 119-125 (1997)); (6) bacterial pneumonia (U.S. Pat. Nos. 5,702,946, 5,677,426, 5,707,622, and 5,686,070); (7) ulcerative colitis (U.S. Pat. Nos. 5,702,946, 5,677,426, 5,707,622, and 5,686,070); and asthma (WO 97/01354 published January 16, 1997).

5 As shown in the Examples below, the conjugates of the invention mimic the in vitro activities of full-length anti-IL-8 monoclonal antibody (e.g. inhibition of IL-8 binding and activation of human neutrophils as shown in Figs. 54A-54C, 55A-55C and 56A-56C and in Example V below), approximate the in vivo pharmacokinetics (e.g. serum half-life, clearance rate and mean residence time as shown in Fig. 65 and in Example X below) and the in vivo therapeutic efficacy (e.g. the treatment of acute lung injury and ARDS as shown in Figs. 70A-70E and in Example Z below and the treatment of ischemic reperfusion injury as shown in Fig. 71 and in Example AA below) of full length anti-IL-8 monoclonal antibody. Since conjugates of the invention derived from anti-IL-8 antibodies and fragments display the same or substantially similar in vitro and in vivo activities as full length anti-IL-8 monoclonal antibody across a range of different parameters, including pharmacokinetic characteristics and therapeutic endpoints in various animal models, the data support the efficacy of the conjugates in the same broad spectrum of disease indications that responds to full length anti-IL-8 antibody treatment.

10 As noted above, any conjugate of the invention derived from an anti-IL-8 antibody or fragment can be advantageously utilized in a method of treating an IL-8 mediated disease or disorder, such as inflammatory diseases. In one embodiment, the invention provides a method of treating an inflammatory disorder in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

25 In another aspect, the invention encompasses the foregoing method of treating inflammatory disorders wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an antibody fragment comprising 6G4.2.5LV/L1N35A as defined below; (2) an antibody fragment comprising 6G4.2.5LV/L1N35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below; (4) an antibody fragment comprising hu6G4.2.5LV/L1N35A as defined below; (5) an antibody fragment comprising hu6G4.2.5LV/L1N35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as defined below; (7) an antibody fragment comprising 6G4.2.5LV/L1N35A and further comprising the CDRs of 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.5LV/L1N35E and further

comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising hu6G4.2.5LV/L1N35A and further comprising hu6G4.2.5HV as defined below; (10) an antibody fragment comprising hu6G4.2.5LV/L1N35E and further comprising hu6G4.2.5HV as defined below; (11) an antibody fragment comprising 6G4.2.5LV11N35A as defined below; (12) an antibody fragment comprising
 5 6G4.2.5LV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.5LV11N35A and further comprising 6G4.2.5HV11 as defined below; and (14) an antibody fragment comprising 6G4.2.5LV11N35E and further comprising 6G4.2.5HV11 as defined below.

In yet another aspect, the invention encompasses any of the foregoing methods of treating an inflammatory disorder wherein the conjugate contains no more than one antibody fragment, wherein the
 10 antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH; wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

In another embodiment, the invention provides a method of treating ischemic reperfusion injury in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every
 20 conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate
 25 one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

In another aspect, the invention encompasses the foregoing method of treating ischemic reperfusion injury wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an
 30 antibody fragment comprising 6G4.2.5LV/L1N35A as defined below; (2) an antibody fragment comprising 6G4.2.5LV/L1N35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below; (4) an antibody fragment comprising hu6G4.2.5LV/L1N35A as defined below; (5) an antibody fragment comprising hu6G4.2.5LV/L1N35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as defined below; (7) an antibody fragment comprising 6G4.2.5LV/L1N35A and further comprising the CDRs of
 35 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.5LV/L1N35E and further comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising hu6G4.2.5LV/L1N35A and further comprising hu6G4.2.5HV as defined below; (10) an antibody fragment comprising hu6G4.2.5LV/L1N35E and further comprising hu6G4.2.5HV as defined below; (11) an antibody fragment comprising 6G4.2.5LV11N35A as defined below; (12) an antibody fragment comprising

6G4.2.SLV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.SLV11N35A and further comprising 6G4.2.5HV11 as defined below; and (14) an antibody fragment comprising 6G4.2.SLV11N35E and further comprising 6G4.2.5HV11 as defined below.

In yet another aspect, the invention encompasses the foregoing methods of treating ischemic reperfusion injury wherein the ischemic reperfusion injury is induced by or incident to a surgical procedure, i.e. a surgical tissue reperfusion injury.

In still another aspect, the invention encompasses the foregoing methods of treating ischemic reperfusion injury wherein the ischemic reperfusion injury is a myocardial ischemic reperfusion injury, such as myocardial infarction, reperfusion after cardiac surgery, cardiac arrest, and constriction after percutaneous transluminal coronary angioplasty.

In yet another aspect, the invention encompasses any of the foregoing methods of treating ischemic reperfusion injury wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

In another embodiment, the invention provides a method of treating acute lung injury in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

In another aspect, the invention encompasses the foregoing method of treating acute lung injury wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an antibody fragment comprising 6G4.2.SLV/L1N35A as defined below; (2) an antibody fragment comprising 6G4.2.SLV/L1N35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below; (4) an antibody fragment comprising hu6G4.2.SLV/L1N35A as defined below; (5) an antibody fragment comprising hu6G4.2.SLV/L1N35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as defined below; (7) an antibody fragment comprising 6G4.2.SLV/L1N35A and further comprising the CDRs of 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.SLV/L1N35E and further comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising

hu6G4.2.SLV/LIN35A and further comprising hu6G4.2.SHV as defined below; (10) an antibody fragment comprising hu6G4.2.SLV/LIN35E and further comprising hu6G4.2.SHV as defined below; (11) an antibody fragment comprising 6G4.2.SLV11N35A as defined below; (12) an antibody fragment comprising 6G4.2.SLV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.SLV11N35A and further comprising 6G4.2.SHV11 as defined below; and (14) an antibody fragment comprising 6G4.2.SLV11N35E and further comprising 6G4.2.SHV11 as defined below.

In yet another aspect, the invention encompasses the foregoing methods of treating acute lung injury wherein the acute lung injury includes adult respiratory distress syndrome (ARDS).

In a further aspect, the invention encompasses any of the foregoing methods of treating acute lung injury wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD.

In a further aspect, the invention encompasses any of the foregoing methods of treating acute lung injury, wherein the patient is selected for prophylactic treatment prior to onset of acute lung injury (with or without progression to ARDS), such as at least 2 hours prior to onset, or at least 90 minutes prior to onset, or at least 60 minutes prior to onset, or at least 30 minutes prior to onset, by the assessment of biological parameters displayed in the patient's condition that indicate likely progression of disease to acute lung injury which may include ARDS, e.g. by using any of the prognostic methods described in Section (II)(5)(B) below, wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

In another embodiment, the invention provides a method of treating hypovolemic shock in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

In another aspect, the invention encompasses the foregoing method of treating hypovolemic shock wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an antibody fragment comprising 6G4.2.5LV/LIN35A as defined below; (2) an antibody fragment comprising 6G4.2.5LV/LIN35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below; (4) an antibody fragment comprising hu6G4.2.5LV/LIN35A as defined below; (5) an antibody fragment comprising hu6G4.2.5LV/LIN35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as defined below; (7) an antibody fragment comprising 6G4.2.5LV/LIN35A and further comprising the CDRs of 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.5LV/LIN35E and further comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising hu6G4.2.5LV/LIN35A and further comprising hu6G4.2.5HV as defined below; (10) an antibody fragment comprising hu6G4.2.5LV/LIN35E and further comprising hu6G4.2.5HV as defined below; (11) an antibody fragment comprising 6G4.2.5LV11N35A as defined below; (12) an antibody fragment comprising 6G4.2.5LV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.5LV11N35A and further comprising 6G4.2.5HV11 as defined below; and (14) an antibody fragment comprising 6G4.2.5LV11N35E and further comprising 6G4.2.5HV11 as defined below.

In yet another aspect, the invention encompasses any of the foregoing methods of treating hypovolemic shock wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

In another embodiment, the invention provides a method of treating an inflammatory bowel disease in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

In another aspect, the invention encompasses the foregoing method of treating an inflammatory bowel disease wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an antibody fragment comprising 6G4.2.5LV/LIN35A as defined below; (2) an antibody fragment comprising 6G4.2.5LV/LIN35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below;

(4) an antibody fragment comprising hu6G4.2.SLV/LIN35A as defined below; (5) an antibody fragment comprising hu6G4.2.SLV/LIN35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as defined below; (7) an antibody fragment comprising 6G4.2.SLV/LIN35A and further comprising the CDRs of 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.SLV/LIN35E and further comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising hu6G4.2.SLV/LIN35A and further comprising hu6G4.2.5HV as defined below; (10) an antibody fragment comprising hu6G4.2.SLV/LIN35E and further comprising hu6G4.2.5HV as defined below; (11) an antibody fragment comprising 6G4.2.SLV11N35A as defined below; (12) an antibody fragment comprising 6G4.2.SLV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.SLV11N35A and further comprising 6G4.2.5HV11 as defined below; and (14) an antibody fragment comprising 6G4.2.SLV11N35E and further comprising 6G4.2.5HV11 as defined below.

In still another aspect, the invention encompasses the foregoing methods of treating an inflammatory bowel disease wherein the inflammatory bowel disease is ulcerative colitis.

In yet another aspect, the invention encompasses any of the foregoing methods of treating inflammatory bowel disease wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

In another embodiment, the invention provides a method of treating a bacterial pneumonia in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MRT and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

In another aspect, the invention encompasses the foregoing method of treating bacterial pneumonia wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an antibody fragment comprising 6G4.2.SLV/LIN35A as defined below; (2) an antibody fragment comprising 6G4.2.SLV/LIN35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below; (4) an antibody fragment comprising hu6G4.2.SLV/LIN35A as defined below; (5) an antibody fragment comprising hu6G4.2.SLV/LIN35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as

defined below; (7) an antibody fragment comprising 6G4.2.5LV/LIN35A and further comprising the CDRs of 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.5LV/LIN35E and further comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising hu6G4.2.5LV/LIN35A and further comprising hu6G4.2.5HV as defined below; (10) an antibody fragment comprising hu6G4.2.5LV/LIN35E and further comprising hu6G4.2.5HV as defined below; (11) an antibody fragment comprising 6G4.2.5LV11N35A as defined below; (12) an antibody fragment comprising 6G4.2.5LV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.5LV11N35A and further comprising 6G4.2.5HV11 as defined below; and (14) an antibody fragment comprising 6G4.2.5LV11N35E and further comprising 6G4.2.5HV11 as defined below.

10 In yet another aspect, the invention encompasses any of the foregoing methods of treating bacterial pneumonia wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

20 In another embodiment, the invention provides a method of treating an asthmatic disease in a mammal comprising administering to the mammal an effective amount of a conjugate selected from the group consisting of: (1) every conjugate described in Section (II)(1) above formed by its component parts, i.e. the antibody fragment or fragments and the nonproteinaceous polymer or polymer molecules that form the conjugate, without any extraneous matter in the covalent molecular structure of the conjugate, (2) every conjugate described in Section (II)(1) above modified to contain one or more additional components, in addition to the antibody fragment component(s) and polymer component(s) that form the conjugate, wherein the modification does not alter the essential functional property of the conjugate of substantially improved serum half-life, MR1 and/or serum clearance rate as compared to that of the parental antibody fragment from which the conjugate is derived, (3) every conjugate described in Section (II)(1) above modified to incorporate one or more nonproteinaceous labels or reporter molecules, and (4) every conjugate described in Section (II)(1) above modified to incorporate one or more radiolabels; wherein at least one antibody fragment in the conjugate comprises an antigen binding site that binds to human IL-8.

30 In another aspect, the invention encompasses the foregoing method of treating an asthmatic disease wherein at least one antibody fragment in the conjugate is selected from the group consisting of: (1) an antibody fragment comprising 6G4.2.5LV/LIN35A as defined below; (2) an antibody fragment comprising 6G4.2.5LV/LIN35E as defined below; (3) an antibody fragment comprising 6G4.2.5HV11 as defined below; (4) an antibody fragment comprising hu6G4.2.5LV/LIN35A as defined below; (5) an antibody fragment comprising hu6G4.2.5LV/LIN35E as defined below; (6) an antibody fragment comprising hu6G4.2.5HV as defined below; (7) an antibody fragment comprising 6G4.2.5LV/LIN35A and further comprising the CDRs of 6G4.2.5HV11 as defined below; (8) an antibody fragment comprising 6G4.2.5LV/LIN35E and further comprising the CDRs of 6G4.2.5HV11 as defined below; (9) an antibody fragment comprising hu6G4.2.5LV/LIN35A and further comprising hu6G4.2.5HV as defined below; (10) an antibody fragment

comprising hu6G4.2.SLV/L1N35E and further comprising hu6G4.2.SHV as defined below; (11) an antibody fragment comprising 6G4.2.SLV11N35A as defined below; (12) an antibody fragment comprising 6G4.2.SLV11N35E as defined below; (13) an antibody fragment comprising 6G4.2.SLV11N35A and further comprising 6G4.2.SHV11 as defined below; and (14) an antibody fragment comprising 6G4.2.SLV11N35E and further comprising 6G4.2.SHV11 as defined below.

In yet another aspect, the invention encompasses the foregoing methods of treating asthmatic disease wherein the asthmatic disease is allergic asthma.

In yet another aspect, the invention encompasses any of the foregoing methods of treating an asthmatic disease wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a linear polyethylene glycol having a molecular weight of at least at or about 20 kD, or at least at or about 30 kD or at least at or about 40 kD, or is a branched polyethylene glycol having a molecular weight of at least at or about 40 kD.

In a preferred embodiment, the invention encompasses any of the foregoing methods of treating inflammatory diseases or asthmatic diseases wherein the mammal is a human.

Therapeutic formulations of the conjugate of the invention can be prepared by utilizing the same procedures described for the formulation of the anti-IL-8 antibodies and fragments of the invention in Section (II)(X)(B) below. The conjugate of the invention can be administered in place of the parent antibody for a given disease indication by modifying the formulation, dosage, administration protocol, and other aspects of a therapeutic regimen as required by the different pharmacodynamic characteristics of the conjugate and as dictated by common medical knowledge and practice.

e. Reagent Uses for Large Effective Size Conjugates

The conjugate of the invention also finds application as a reagent in an animal model system for in vivo study of the biological functions of the antigen recognized by the conjugate. The conjugate would enable the practitioner to inactivate or detect the cognate antigen in circulation or in tissue for a far greater period of time than would be possible with art-known constructs while removing any Fc interaction (which could attend the use of an intact antibody) from the system. In addition, the increased half-life of the conjugate of the invention can be applied advantageously to the induction of tolerance for the underivatized antibody fragment in a test animal by employing the Wie *et al.*, Int. Archs. Allergy Appl. Immunol., 64: 84-99 (1981) method for allergen tolerization, which would permit the practitioner to repeatedly challenge the tolerized animal with the underivatized parental antibody fragment without generating an immune response against the parental fragment.

2. HUMANIZED 6G4.2.5 MONOCLONAL ANTIBODIES AND ANTIBODY FRAGMENTS

In one embodiment, the invention provides an antibody fragment or full length antibody comprising a heavy chain comprising the amino acid sequence of amino acids 1-230 (herein referred to as "6G4.2.SHV11") of the humanized anti-IL-8 6G4.2.Sv11 heavy chain polypeptide amino acid sequence of Figs. 37A-37B (SEQ ID NO: 60).

The invention encompasses a single chain antibody fragment comprising the 6G4.2.SHV11, with or

without any additional amino acid sequence. In one embodiment, the invention provides a single chain antibody fragment comprising the 6G4.2.5HV11 without any associated light chain amino acid sequence, i.e. a single chain species that makes up one half of a Fab fragment.

Further provided herein are an antibody or antibody fragment comprising the 6G4.2.5HV11, and
5 further comprising a light chain comprising the amino acid sequence of amino acids 1-219 (herein referred to as "6G4.2.5LV11") of the humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51).

In one embodiment, the invention provides a single chain antibody fragment wherein the 6G4.2.5HV11 and the 6G4.2.5LV11 are contained in a single chain polypeptide species. In a preferred
10 embodiment, the single chain antibody fragment comprises the 6G4.2.5HV11 joined to the 6G4.2.5LV11 by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In another embodiment, the single chain antibody fragment is a species comprising the 6G4.2.5HV11 joined to the 6G4.2.5LV11 by a linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide
15 monomer that forms a diabody upon dimerization with another monomer.

In yet another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the 6G4.2.5HV11 and a second polypeptide chain comprises the 6G4.2.5LV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the foregoing two-chain antibody fragment is selected
20 from the group consisting of Fab, Fab', Fab'-SH, and F(ab')₂.

The invention also provides an antibody or antibody fragment comprising a heavy chain containing the 6G4.2.5HV11 and optionally further comprising a light chain containing the 6G4.2.5LV11, wherein the heavy chain, and optionally the light chain, is (are) fused to an additional moiety, such as additional immunoglobulin constant domain sequence. Constant domain sequence can be added to the heavy chain
25 and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.* (*supra*).

In a preferred embodiment, the antibody or antibody fragment comprises the 6G4.2.5HV11 in a heavy
30 chain that is fused to or contains a leucine zipper sequence. The leucine zipper can increase the affinity and/or production efficiency of the antibody or antibody fragment of interest. Suitable leucine zipper sequences include the jun and fos leucine zippers taught by Kostelny *et al.*, *J. Immunol.*, 148: 1547-1553 (1992) and the GCN4 leucine zipper described in the Examples below. In a preferred embodiment, the antibody or antibody
35 fragment comprises the 6G4.2.5HV11 fused at its C-terminus to the GCN4 leucine zipper to yield the amino acid sequence of amino acids 1-275 (herein referred to as "6G4.2.5HV11GCN4") of the heavy chain polypeptide amino acid sequence of Figs. 37A-37B (SEQ ID NO: 60).

3. VARIANTS OF HUMANIZED 6G4.2.5 MONOCLONAL ANTIBODIES AND ANTIBODY FRAGMENTS

The invention additionally encompasses humanized anti-IL-8 monoclonal antibody and antibody fragments comprising variants of the 6G4.2.5 complementarity determining regions (CDRs) or variants of the 6G4.2.5v11 variable domains which exhibit higher affinity for human IL-8 and/or possess properties that yield greater efficiency in recombinant production processes.

A. 6G4.2.5LV VARIANTS

In one aspect, the invention provides humanized anti-IL-8 monoclonal antibodies and antibody fragments comprising the complementarity determining regions (referred to herein as the "CDRs of 6G4.2.5LV") L1, L2, and L3 of the 6G4.2.5 light chain variable domain amino acid sequence of Fig. 24, wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35).

In addition, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a humanized light chain variable domain comprising a variant (hereinafter referred to as "6G4.2.5LV CDRs variant") of the complementarity determining regions L1, L2, and L3 of the 6G4.2.5 variable light chain domain amino acid sequence of Fig. 24 (SEQ ID NO: 35). In one embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1N35X₃₅") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35). In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1N35A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Ala is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35). In another preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1N35E") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Glu is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35).

In a second aspect, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1S26X₂₆") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26, L2

corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35). In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1S26A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Ala is substituted for Ser at amino acid position 26, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35).

In a third aspect, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L3H98X₉₈") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98. In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L3H98A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Ala is substituted for His at amino acid position 98.

In a fourth aspect, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1S26X₂₆,N35X₃₅") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26 and any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35). In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1S26A,N35A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Ala is substituted for Ser at amino acid position 26 and Ala is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35).

In a fifth aspect, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1N35X₃₅/L3H98X₉₈") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds

to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98. In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1N35A/L3H98A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that Ala is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that Ala is substituted for His at amino acid position 98.

In a sixth aspect, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1S26X₂₆/L3H98X₉₈") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98. In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (herein referred to as "6G4.2.5LV/L1S26A/L3H98A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that Ala is substituted for Ser at amino acid position 26, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO 35) with the proviso that Ala is substituted for His at amino acid position 98.

In a seventh aspect, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (here referred to as "6G4.2.5LV/L1S26X₂₆,N35X₃₅/L3H98X₉₈") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26 and any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98. In a preferred embodiment, the invention provides a variant 6G4.2.5 humanized antibody or antibody fragment comprising a 6G4.2.5LV CDRs variant (here referred to as "6G4.2.5LV/L1S26A,N35A/L3H98A") wherein L1 corresponds to amino acids 24-39 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Ala is substituted for Ser at amino acid position 26 and Ala is substituted for Asn at amino acid position 35, L2 corresponds to amino acids 55-61 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35), and L3 corresponds to amino acids 94-102 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35) with the proviso that Ala is substituted for His at amino acid position 98.

The humanized light chain variable domains of the invention can be constructed by using any of the techniques for antibody humanization known in the art. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, Nature 321:522 (1986); Riechmann *et al.*, Nature 332:323 (1988); Verhoeven *et al.*, Science 239:1534 (1988)), by substituting the CDRs of 6G4.2.5LV or the CDRs of a 6G4.2.5LV CDRs variant for the corresponding sequences of a human antibody light chain variable domain. Accordingly, such "humanized" derivatives containing the CDRs of 6G4.2.5LV or the CDRs of a 6G4.2.5LV CDRs variant are chimeric (Cabilly *et al.*, supra). The humanized light chain variable domain comprising the CDRs of 6G4.2.5LV or the CDRs of a 6G4.2.5LV CDRs variant can also contain some FR residues that are substituted by residues from analogous sites in the murine 6G4.2.5 antibody light chain variable domain ("6G4.2.5LV"). The complete amino acid sequence of 6G4.2.5LV is set out as amino acids 1-114 of the amino acid sequence of Fig. 24 (SEQ ID NO: 35).

The invention further provides a humanized antibody or antibody fragment comprising a humanized light chain variable domain comprising the CDRs of 6G4.2.5LV or the CDRs of a 6G4.2.5LV CDRs variant as described above, and further comprising a humanized heavy chain variable domain comprising the complementarity determining regions (CDRs) H1, H2, and H3 of the 6G4.2.5 (murine monoclonal antibody) variable heavy chain domain amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37). The above-described H1, H2, and H3 CDRs of the 6G4.2.5 heavy chain variable domain ("6G4.2.5HV") are collectively referred to as the "CDRs of 6G4.2.5HV".

In another embodiment, the invention provides a humanized antibody or antibody fragment comprising a humanized light chain variable domain comprising the CDRs of 6G4.2.5LV or the CDRs of a 6G4.2.5LV CDRs variant as described above, and further comprising a humanized heavy chain variable domain comprising a variant (herein referred to as a "6G4.2.5HV CDRs variant") of the H1, H2, and H3 CDRs of the 6G4.2.5 (murine monoclonal antibody) variable heavy chain domain amino acid sequence of Fig. 25 (SEQ ID NO: 37). In one 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31Z₃₁"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37). In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31A"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37).

In a second 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino

acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37). In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37).

In a third 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3D100E"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100.

In a fourth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3R102K"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102.

In a fifth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3D106E"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106.

In a seventh 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3D100E,R102K"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102.

In an eighth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3R102K,D106E"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

In a ninth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3D100E,D106E"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position

106.

In a tenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H3D100E,R102K,D106E"), wherein H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), wherein H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and
 5 wherein H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102, and Glu is substituted for Asp at amino acid position 106.

In an eleventh 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the provis
 10 that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37). In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A"), H1 correspond to amino
 15 acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37).

~~In a twelfth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H3D100E"), H1~~
 20 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100. In a preferred 6G4.2.5HV CDRs variant (referred to herein as
 25 "6G4.2.5HV/H1S31A/H3D100E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100.

30 In a thirteenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H3R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is
 35 substituted for Arg at amino acid position 102. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H3R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25

(SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102.

5 A fourteenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31Z₃₁/H3D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31A/H3D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106.

15 A fifteenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31Z₃₁/H3D100E,R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31A/H3D100E,R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102.

20 In a sixteenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31Z₃₁/H3R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/HIS31A/H3R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of

Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

In a seventeenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H3D100E,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H3D100E,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position 106.

In an eighteenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H3D100E,R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H3D100E,R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

In a nineteenth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3D100E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3

corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A/H3D100E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100.

In a twentieth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3R102K"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A/H3R102K"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102.

In a twenty-first 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A/H3D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106.

In a twenty-second 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3D100E,R102K"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102. In a preferred 6G4.2.5HV CDRs variant (referred to herein as

"6G4.2.5HV/H2S54A/H3D100E,R102K"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102.

In a twenty-third 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3R102K,D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A/H3R102K,D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

In a twenty-fourth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3D100E,D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A/H3D100E,D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position 106.

In a twenty-fifth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54Z₅₄/H3D100E,R102K,D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted

for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H2S54A/H3D100E,R102K,D106E"), H1 corresponds to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37), H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

In a twenty-sixth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3D100E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100.

In a twenty-seventh 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the

amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102.

In a twenty-eighth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 106.

In a twenty-ninth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3D100E,R102K"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Lys is substituted for Arg at amino acid position 102.

In a thirtieth 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄")

is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

In a thirty-first 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3D100E,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100 and Glu is substituted for Asp at amino acid position 106.

In a thirty-second 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₃₁") is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that any amino acid other than Ser (denoted as "Z₅₄") is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106. In a preferred 6G4.2.5HV CDRs variant (referred to herein as "6G4.2.5HV/H1S31A/H2S54A/H3D100E,R102K,D106E"), H1 correspond to amino acids 26-35 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 31, H2 corresponds to amino acids 50-66 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Ala is substituted for Ser at amino acid position 54, and H3 corresponds to amino acids 99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

99-111 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37) with the proviso that Glu is substituted for Asp at amino acid position 100, Lys is substituted for Arg at amino acid position 102 and Glu is substituted for Asp at amino acid position 106.

As in the humanization of the light chain variable domain described above, a humanized heavy chain variable domain is constructed by substituting the CDRs of 6G4.2.5HV or the CDRs of a 6G4.2.5HV CDRs variant for the corresponding sequences in a human heavy chain variable domain. The humanized heavy chain variable domain comprising the CDRs of 6G4.2.5HV or the CDRs of a 6G4.2.5HV CDRs variant can also contain some FR residues that are substituted by residues from analogous sites in the murine 6G4.2.5 antibody heavy chain variable domain. The complete amino acid sequence of 6G4.2.5HV is set out as amino acids 1-122 of the amino acid sequence of Fig. 25 (SEQ ID NO: 37).

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies and antibody fragments is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody (Sims *et al.*, J. Immunol. 151: 2296 (1993); Chothia and Lesk, J. Mol. Biol. 196:901 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework can be used for several different humanized antibodies (Carter *et al.*, Proc. Natl. Acad. Sci. U.S.A. 89:4285 (1992); Presta *et al.*, J. Immunol. 151:2623 (1993)).

It is also important that the antibodies and antibody fragments of the invention be humanized with retention of high affinity for human IL-8 and other favorable biological properties. To achieve this goal, according to a preferred method, the humanized antibodies and antibody fragments of the invention are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the consensus and parental sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved.

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV are collectively referred to herein as "hu6G4.2.5LV".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIN35X₃₅ are collectively referred to herein as "hu 6G4.2.5LV/LIN35X₃₅".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIN35A are collectively referred to herein as "hu6G4.2.5LV/LIN35A".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs

of 6G4.2.5LV/LIN35E are collectively referred to herein as "hu6G4.2.5LV/LIN35E".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26X₂₆ are collectively referred to herein as "hu6G4.2.5LV/LIS26X₂₆".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26A are collectively referred to herein as "hu6G4.2.5LV/LIS26A".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/L3H98X₉₈ are collectively referred to herein as "hu6G4.2.5LV/L3H98X₉₈".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/L3H98A are collectively referred to herein as "hu6G4.2.5LV/L3H98A".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26X₂₆,N35X₃₅ are collectively referred to herein as "hu6G4.2.5LV/LIS26X₂₆,N35X₃₅".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26A,N35A are collectively referred to herein as "hu6G4.2.5LV/LIS26A,N35A".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIN35X₃₅/L3H98X₉₈ are collectively referred to herein as "hu6G4.2.5LV/LIN35X₃₅/L3H98X₉₈".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIN35A/L3H98A are collectively referred to herein as "hu6G4.2.5LV/LIN35A/L3H98A".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26X₂₆/L3H98X₉₈ are collectively referred to herein as "hu6G4.2.5LV/LIS26X₂₆/L3H98X₉₈".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26A/L3H98A are collectively referred to herein as "hu6G4.2.5LV/LIS26A/L3H98A".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26X₂₆,N35X₃₅/L3H98X₉₈ are collectively referred to herein as "hu6G4.2.5LV/LIS26X₂₆,N35X₃₅/L3H98X₉₈".

Any and all humanized light chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5LV/LIS26A,N35A/L3H98A are collectively referred to herein as "hu6G4.2.5LV/LIS26A,N35A/L3H98A".

The humanized light chain variable domain amino acid sequences of hu6G4.2.5LV/LIN35X₃₅, hu6G4.2.5LV/LIS26X₂₆, hu6G4.2.5LV/LIS26X₂₆/L3H98X₉₈, hu6G4.2.5LV/LIS26X₂₆,N35X₃₅, hu6G4.2.5LV/LIN35X₃₅/L3H98X₉₈, hu6G4.2.5LV/LIS26X₂₆/L3H98X₉₈, and hu6G4.2.5LV/LIS26X₂₆,N35X₃₅/L3H98X₉₈ are collectively referred to herein as "hu6G4.2.5LV/vL1-3X".

The humanized light chain variable domain amino acid sequences of hu6G4.2.5LV/LIN35A,

hu6G4.2.5LV/L1S26A, hu6G4.2.5LV/L1S26A/L3H98A, hu6G4.2.5LV/L1S26A,N35A, hu6G4.2.5LV/L1N35A/L3H98A, hu6G4.2.5LV/L1S26A/L3H98A, hu6G4.2.5LV/L1S26A,N35A/L3H98A are collectively referred to herein as "hu6G4.2.5LV/vL1-3A".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV are collectively referred to herein as "hu6G4.2.5HV".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁ are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A are collectively referred to herein as "hu6G4.2.5HV/H1S31A".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54Z₅₄ are collectively referred to herein as "hu6G4.2.5HV/H2S54Z₅₄".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A are collectively referred to herein as "hu6G4.2.5HV/H2S54A".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3D100E are collectively referred to herein as "hu6G4.2.5HV/H3D100E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3R102K are collectively referred to herein as "hu6G4.2.5HV/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3D106E are collectively referred to herein as "hu6G4.2.5HV/H3D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3D100E,R102K are collectively referred to herein as "hu6G4.2.5HV/H3D100E,R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3D100E,D106E are collectively referred to herein as "hu6G4.2.5HV/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H3D100E,R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄ are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H3D100E are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H3D100E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H3R102K are collectively referred to herein as

"hu6G4.2.5HV/HIS31Z₃₁/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31Z₃₁/H3D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31Z₃₁/H3D106E".

5 Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31Z₃₁/H3D100E,R102K are collectively referred to herein as "hu6G4.2.5HV/HIS31Z₃₁/H3D100E,R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31Z₃₁/H3R102K,D106E are collectively referred to herein as 10 "hu6G4.2.5HV/HIS31Z₃₁/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31Z₃₁/H3D100E,D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31Z₃₁/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs 15 of 6G4.2.5HV/HIS31Z₃₁/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31Z₃₁/H3D100E,R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54Z₅₄/H3D100E are collectively referred to herein as "hu6G4.2.5HV/H2S54Z₅₄/H3D100E".

20 Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54Z₅₄/H3R102K are collectively referred to herein as "hu6G4.2.5HV/H2S54Z₅₄/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54Z₅₄/H3D106E are collectively referred to herein as 25 "hu6G4.2.5HV/H2S54Z₅₄/H3D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54Z₅₄/H3R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54Z₅₄/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs 30 of 6G4.2.5HV/H2S54Z₅₄/H3D100E,D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54Z₅₄/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54Z₅₄/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54Z₅₄/H3D100E,R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A/H2S54A are collectively referred to herein as "hu6G4.2.5HV/H1S31A/H2S54A".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A/H3D100E are collectively referred to herein as "hu6G4.2.5HV/H1S31A/H3D100E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A/H3R102K are collectively referred to herein as "hu6G4.2.5HV/H1S31A/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A/H3D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31A/H3D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs

of 6G4.2.5HV/HIS31A/H3D100E,R102K are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H3D100E,R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H3R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H3D100E,D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H3D100E,R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A/H3D100E are collectively referred to herein as "hu6G4.2.5HV/H2S54A/H3D100E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A/H3R102K are collectively referred to herein as "hu6G4.2.5HV/H2S54A/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A/H3D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54A/H3D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A/H3R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54A/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A/H3D100E,D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54A/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H2S54A/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H2S54A/H3D100E,R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H2S54A/H3D100E are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H2S54A/H3D100E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H2S54A/H3R102K are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H2S54A/H3R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H2S54A/H3D106E are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H2S54A/H3D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/HIS31A/H2S54A/H3D100E,R102K are collectively referred to herein as "hu6G4.2.5HV/HIS31A/H2S54A/H3D100E,R102K".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs

of 6G4.2.5HV/H1S31A/H2S54A/H3R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31A/H2S54A/H3R102K,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A/H2S54A/H3D100E,D106E are collectively referred to herein as
5 "hu6G4.2.5HV/H1S31A/H2S54A/H3D100E,D106E".

Any and all humanized heavy chain variable domain amino acid sequences which comprise the CDRs of 6G4.2.5HV/H1S31A/H2S54A/H3D100E,R102K,D106E are collectively referred to herein as "hu6G4.2.5HV/H1S31A/H2S54A/H3D100E,R102K,D106E".

The humanized heavy chain variable domain amino acid sequences of
10 hu6G4.2.5HV/H1S31Z₃₁, hu6G4.2.5HV/H2S54Z₅₄, hu6G4.2.5HV/H3D100E, hu6G4.2.5HV/H3R102K, hu6G4.2.5HV/H3D106E, hu6G4.2.5HV/H3D100E,R102K, hu6G4.2.5HV/H3R102K,D106E, hu6G4.2.5HV/H3D100E,D106E, hu6G4.2.5HV/H3D100E,R102K,D106E, hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄, hu6G4.2.5HV/H1S31Z₃₁/H3D100E, hu6G4.2.5HV/H1S31Z₃₁/H3R102K, hu6G4.2.5HV/H1S31Z₃₁/H3D106E,
15 hu6G4.2.5HV/H1S31Z₃₁/H3D100E,R102K, hu6G4.2.5HV/H1S31Z₃₁/H3R102K,D106E, hu6G4.2.5HV/H1S31Z₃₁/H3D100E,D106E, hu6G4.2.5HV/H1S31Z₃₁/H3D100E,R102K,D106E, hu6G4.2.5HV/H2S54Z₅₄/H3D100E, hu6G4.2.5HV/H2S54Z₅₄/H3R102K, hu6G4.2.5HV/H2S54Z₅₄/H3D106E, hu6G4.2.5HV/H2S54Z₅₄/H3R102K,D106E, hu6G4.2.5HV/H2S54Z₅₄/H3D100E,D106E, hu6G4.2.5HV/H2S54Z₅₄/H3D100E,R102K,D106E,
20 hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E, hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K, hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D106E, hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K, hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3R102K,D106E, hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,D106E, and
hu6G4.2.5HV/H1S31Z₃₁/H2S54Z₅₄/H3D100E,R102K,D106E are collectively referred to herein as
25 "hu6G4.2.5HV/vH1-3Z".

The humanized heavy chain variable domain amino acid sequences of
hu6G4.2.5HV/H1S31A, hu6G4.2.5HV/H2S54A, hu6G4.2.5HV/H3D100E, hu6G4.2.5HV/H3R102K, hu6G4.2.5HV/H3D106E, hu6G4.2.5HV/H3D100E,R102K, hu6G4.2.5HV/H3R102K,D106E, hu6G4.2.5HV/H3D100E,D106E, hu6G4.2.5HV/H3D100E,R102K,D106E, hu6G4.2.5HV/H1S31A/H2S54A,
30 hu6G4.2.5HV/H1S31A/H3D100E, hu6G4.2.5HV/H1S31A/H3R102K, hu6G4.2.5HV/H1S31A/H3D106E, hu6G4.2.5HV/H1S31A/H3D100E,R102K, hu6G4.2.5HV/H1S31A/H3R102K,D106E, hu6G4.2.5HV/H1S31A/H3D100E,D106E, hu6G4.2.5HV/H1S31A/H3D100E,R102K,D106E, hu6G4.2.5HV/H2S54A/H3D100E, hu6G4.2.5HV/H2S54A/H3R102K, hu6G4.2.5HV/H2S54A/H3D106E, hu6G4.2.5HV/H2S54A/H3R102K,D106E, hu6G4.2.5HV/H2S54A/H3D100E,D106E,

hu6G4.2.5HV/H2S54A/H3D100E,R102K,D106E, hu6G4.2.5HV/H1S31A/H2S54A/H3D100E,
 hu6G4.2.5HV/H1S31A/H2S54A/H3R102K, hu6G4.2.5HV/H1S31A/H2S54A/H3D106E,
 hu6G4.2.5HV/H1S31A/H2S54A/H3D100E,R102K, hu6G4.2.5HV/H1S31A/H2S54A/H3R102K,D106E,
 hu6G4.2.5HV/H1S31A/H2S54A/H3D100E,D106E, and
 5 hu6G4.2.5HV/H1S31A/H2S54A/H3D100E,R102K,D106E are collectively referred to herein as
 "hu6G4.2.5HV/vH1-3A".

The invention provides a humanized antibody or antibody fragment that comprises a light chain
 variable domain comprising the hu6G4.2.5LV/vL1-3X. In another embodiment, the invention provides a
 humanized antibody or antibody fragment that comprises a light chain variable domain comprising the
 10 hu6G4.2.5LV/vL1-3A. In yet another embodiment, the invention provides a humanized antibody or antibody
 fragment that comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35X₃₅. In still
 another embodiment, the invention provides a humanized antibody or antibody fragment that comprises a light
 chain variable domain comprising the hu6G4.2.5LV/L1N35A. In a further embodiment, the invention
 provides a humanized antibody or antibody fragment that comprises a light chain variable domain comprising
 15 the hu6G4.2.5LV/L1N35E.

The invention additionally provides a humanized antibody or antibody fragment that comprises a light
 chain variable domain comprising the hu6G4.2.5LV/vL1-3X, and further comprises a heavy chain variable
 domain comprising the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z. In another embodiment, the inventor
 provides a humanized antibody or antibody fragment that comprises a light chain variable domain comprising
 20 the hu6G4.2.5LV/vL1-3A, and further comprises a heavy chain variable domain comprising the hu6G4.2.5HV
 or hu6G4.2.5HV/vH1-3Z. In yet another embodiment, the invention provides a humanized antibody or
 antibody fragment that comprises a light chain variable domain comprising the hu6G4.2.5LV/vL1-3A, and
 further comprises a heavy chain variable domain comprising the hu6G4.2.5HV/vH1-3A.

In a further embodiment, the invention provides a humanized antibody or antibody fragment that
 25 comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35X₃₅, and further comprises a
 heavy chain variable domain comprising the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z. In another embodiment,
 the invention provides a humanized antibody or antibody fragment that comprises a light chain variable
 domain comprising the hu6G4.2.5LV/N35X₃₅, and further comprises a heavy chain variable domain
 comprising the hu6G4.2.5HV/vH1-3A. In a preferred embodiment, the antibody or antibody fragment
 30 comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35X₃₅ and further comprises a
 humanized heavy chain comprising the amino acid sequence of 6G4.2.5HV11.

In an additional embodiment, the invention provides a humanized antibody or antibody fragment that
 comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35A, and further comprises a heavy
 chain variable domain comprising the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z. In another embodiment, the
 35 invention provides a humanized antibody or antibody fragment that comprises a light chain variable domain
 comprising the hu6G4.2.5LV/N35A, and further comprises a heavy chain variable domain comprising the
 hu6G4.2.5HV/vH1-3A. In still another embodiment, the humanized antibody or antibody fragment comprises

a light chain variable domain comprising the hu6G4.2.5LV/L1N35A, and further comprises a heavy chain variable domain comprising the hu6G4.2.5HV. In a further embodiment, the humanized antibody or antibody fragment comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35E, and further comprises a heavy chain variable domain comprising the hu6G4.2.5HV. In a preferred embodiment, the antibody or antibody fragment comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35A and further comprises a humanized heavy chain comprising the amino acid sequence of 6G4.2.5HV11. In another preferred embodiment, the antibody or antibody fragment comprises a light chain variable domain comprising the hu6G4.2.5LV/L1N35E and further comprises a humanized heavy chain comprising the amino acid sequence of 6G4.2.5HV11.

The invention encompasses a single chain antibody fragment comprising the hu6G4.2.5LV/vL1-3X, with or without any additional amino acid sequence. In one embodiment, the invention provides a single chain antibody fragment comprising the hu6G4.2.5LV/vL1-3X without any associated heavy chain variable domain amino acid sequence, i.e. a single chain species that makes up one half of an Fv fragment. In another embodiment, the invention provides a single chain antibody fragment comprising the hu6G4.2.5LV/vL1-3A without any associated heavy chain variable domain amino acid sequence. In still another embodiment, the invention provides a single chain antibody fragment comprising the hu6G4.2.5LV/L1N35X₃₅ without any associated heavy chain variable domain amino acid sequence. In a preferred embodiment, the invention provides a single chain antibody fragment comprising the hu6G4.2.5LV/L1N35A without any associated heavy chain variable domain amino acid sequence. In another preferred embodiment, the invention provides a single chain antibody fragment comprising the hu6G4.2.5LV/L1N35E without any associated heavy chain variable domain amino acid sequence.

In one embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/vL1-3X and the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/vL1-3X joined to the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/vL1-3X joined to the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In another embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/vL1-3A and the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/vL1-3A joined to the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/vL1-3A joined to the hu6G4.2.5HV or

hu6G4.2.5HV/vH1-3Z by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

5 In yet another embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/vL1-3A and the hu6G4.2.5HV/vH1-3A are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/vL1-3A joined to the hu6G4.2.5HV/vH1-3A by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the
10 hu6G4.2.5LV/vL1-3A joined to the hu6G4.2.5HV/vH1-3A by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In still another embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/L1N35X₃₅ and the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z are contained in a single chain
15 polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/L1N35X₃₅ joined to the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/L1N35X₃₅ joined to the hu6G4.2.5HV or
20 hu6G4.2.5HV/vH1-3Z by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In a further embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/L1N35X₃₅ and the hu6G4.2.5HV/vH1-3A are contained in a single chain polypeptide species.
25 In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/L1N35X₃₅ joined to the hu6G4.2.5HV/vH1-3A by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/L1N35X₃₅ joined to the hu6G4.2.5HV/vH1-3A by a linker that is too short to
30 permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In an additional embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/L1N35A and the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species
35 comprising the hu6G4.2.5LV/L1N35A joined to the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a

"dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/L1N35A joined to the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

Also provided herein is a single chain antibody fragment wherein the hu6G4.2.5LV/L1N35E and the hu6G4.2.5HV are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/L1N35E joined to the hu6G4.2.5HV by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/L1N35E joined to the hu6G4.2.5HV by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In still another embodiment, the invention provides a single chain antibody fragment wherein the hu6G4.2.5LV/L1N35A and the hu6G4.2.5HV/vH1-3A are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5LV/L1N35A joined to the hu6G4.2.5HV/vH1-3A by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5LV/L1N35A joined to the hu6G4.2.5HV/vH1-3A by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In yet another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/vL1-3X and a second polypeptide chain comprises the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

In still another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/vL1-3X and a second polypeptide chain comprises the hu6G4.2.5HV/vH1-3A and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/vL1-3X and a second polypeptide chain comprises the amino acid sequence of 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

In a further embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/vL1-3A and a second polypeptide chain comprises the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

In still another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/vL1-3A and a second

polypeptide chain comprises the hu6G4.2.5HV/vH1-3A and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/vL1-3A and a second polypeptide chain comprises the amino acid sequence of 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

The invention also encompasses an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35X₃₅ and a second polypeptide chain comprises the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

In still another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35X₃₅ and a second polypeptide chain comprises the hu6G4.2.5HV/vH1-3A and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35X₃₅ and a second polypeptide chain comprises the amino acid sequence of 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

The invention further encompasses an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35A and a second polypeptide chain comprises the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

The invention also encompasses an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35E and a second polypeptide chain comprises the hu6G4.2.5HV and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

In still another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35A and a second polypeptide chain comprises the hu6G4.2.5HV/vH1-3A and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35A and a second polypeptide chain comprises the amino acid sequence of 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In another preferred embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5LV/L1N35E and a second polypeptide chain comprises the amino acid sequence of 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds.

In a preferred embodiment, any of the foregoing two-chain antibody fragments are selected from the group consisting of Fab, Fab', Fab'-SH, Fv, and F(ab')₂. In another preferred embodiment, the antibody

fragment is selected from the group consisting of Fab, Fab', Fab'-SH, Fv, and F(ab')₂, wherein the antibody fragment comprises one polypeptide chain comprising the hu6G4.2.5LV/L1N35X₃₅ and a second polypeptide chain comprising the hu6G4.2.5HV. In yet another preferred embodiment, the antibody fragment is selected from the group consisting of Fab, Fab', Fab'-SH, Fv, and F(ab')₂, wherein the antibody fragment comprises one polypeptide chain comprising the hu6G4.2.5LV/L1N35A and a second polypeptide chain comprising the hu6G4.2.5HV. In a further preferred embodiment, the antibody fragment is selected from the group consisting of Fab, Fab', Fab'-SH, Fv, and F(ab')₂, wherein the antibody fragment comprises one polypeptide chain comprising the hu6G4.2.5LV/L1N35E and a second polypeptide chain comprising the hu6G4.2.5HV. In still another preferred embodiment, the antibody fragment is a F(ab')₂ that comprises one polypeptide chain comprising the hu6G4.2.5LV/L1N35A and a second polypeptide chain comprising the amino acid sequence of 6G4.2.5HV11. In an additional preferred embodiment, the antibody fragment is a F(ab')₂ that comprises one polypeptide chain comprising the hu6G4.2.5LV/L1N35E and a second polypeptide chain comprising the amino acid sequence of 6G4.2.5HV11.

The invention also provides an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/vL1-3X and optionally further comprising a heavy chain variable domain containing the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z, wherein the light chain variable domain, and optionally the heavy chain variable domain, is (are) fused to an additional moiety, such as a immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

The invention additionally provides an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/vL1-3X and optionally further comprising a heavy chain variable domain containing the hu6G4.2.5HV/vH1-3A, wherein the light chain variable domain, and optionally the heavy chain variable domain, is (are) fused to an additional moiety, such as a immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

The invention further provides an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/L1N35X₃₅ and optionally further comprising a heavy chain variable domain containing the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z, wherein the light chain variable domain, and optionally the heavy chain variable domain, is (are) fused to an additional moiety, such as a immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to

form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

5 The invention additionally provides an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/L1N35X₃₅ and optionally further comprising a heavy chain variable domain containing the hu6G4.2.5HV/vH1-3A, wherein the light chain variable domain, and optionally the heavy chain variable domain, is (are) fused to an additional moiety, such as a immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form
10 species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

15 The invention also encompasses an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/L1N35A and optionally further comprising a heavy chain variable domain containing the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z, wherein the light chain variable domain, and optionally the heavy chain variable domain, is (are) fused to an additional moiety, such as a immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions
20 of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

25 The invention additionally provides an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/L1N35A and optionally further comprising a heavy chain variable domain containing the hu6G4.2.5HV/vH1-3A, wherein the light chain variable domain, and optionally the heavy chain variable domain, is (are) fused to an additional moiety, such as a immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that
30 such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

35 The invention additionally encompasses an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/L1N35A and optionally further comprising a heavy chain containing the amino acid sequence of 6G4.2.5HV11, wherein the light chain variable domain, and optionally the heavy chain, is (are) fused to an additional moiety, such as immunoglobulin constant domain sequences. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant

regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

The invention further encompasses an antibody or antibody fragment comprising a light chain variable domain containing the hu6G4.2.5LV/L1N35E and optionally further comprising a heavy chain containing the amino acid sequence of 6G4.2.5HV11, wherein the light chain variable domain, and optionally the heavy chain, is (are) fused to an additional moiety, such as immunoglobulin constant domain sequences. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

In a preferred embodiment, the antibody or antibody fragment comprises a light chain variable domain containing the hu6G4.2.5LV/vL1-3X, and further comprises the hu6G4.2.5HV or hu6G4.2.5HV/vH1-3Z in a heavy chain that is fused to or contains a leucine zipper sequence. The leucine zipper can increase the affinity or production efficiency of the antibody or antibody fragment of interest. Suitable leucine zipper sequences include the jun and fos leucine zippers taught by Kostelny *et al.*, J. Immunol., 148: 1547-1553 (1992) and the GCN4 leucine zipper described in the Examples below.

In particular, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35 (herein referred to as "6G4.2.5LV11N35X₃₅").

In another embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26 (herein referred to as "6G4.2.5LV11S26X₂₆").

In yet another embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11H98X₉₈").

In still another embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26 and any

amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35 (herein referred to as "6G4.2.5LV11S26X₂₆/N35X₃₅").

5 In a further embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35 and any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11N35X₃₅/H98X₉₈").

10 In an additional embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26 and any amino acid other than His (denoted as "X₉₈") is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11S26X₂₆/H98X₉₈").

15 The invention also encompasses an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that any amino acid other than Ser (denoted as "X₂₆") is substituted for Ser at amino acid position 26, any amino acid other than Asn (denoted as "X₃₅") is substituted for Asn at amino acid position 35 and any amino acid other than His
20 (denoted as "X₉₈") is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11S26X₂₆/N35X₃₅/H98X₉₈").

25 Additionally, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence (SEQ ID NO: 56) of Fig. 36 (herein referred to as "6G4.2.5LV11N35A").

Further provided herein is an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence (SEQ ID NO: 62) of Fig. 45 (herein referred to as "6G4.2.5LV11N35E").

30 In another embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that Ala is substituted for Ser at amino acid position 26 (herein referred to as "6G4.2.5LV11S26A").

In yet another embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8

6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that Ala is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11H98A").

In still another embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8
5 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that Ala is substituted for Ser at amino acid position 26 and Ala is substituted for Asn at amino acid position 35 (herein referred to as "6G4.2.5LV11S26A/N35A").

In a further embodiment, the invention provides an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8
10 6G4.2.5v11 light chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that Ala is substituted for Ser at amino acid position 26 and Ala is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11S26A/H98A").

The invention also encompasses an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light
15 chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that Ala is substituted for Asn at amino acid position 35 and Ala is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11N35A/H98A").

The invention further encompasses an antibody or antibody fragment comprising a light chain comprising the amino acid sequence of amino acids 1-219 of the variant humanized anti-IL-8 6G4.2.5v11 light
20 chain polypeptide amino acid sequence of Fig. 31B (SEQ ID NO: 51) with the proviso that Ala is substituted for Ser at amino acid position 26, Ala is substituted for Asn at amino acid position 35, and Ala is substituted for His at amino acid position 98 (herein referred to as "6G4.2.5LV11S26A/N35A/H98A").

The invention provides a single chain antibody fragment comprising a variant light chain selected from the group consisting of 6G4.2.5LV11N35X₃₅, 6G4.2.5LV11S26X₂₆, 6G4.2.5LV11H98X₉₈,
25 6G4.2.5LV11S26X₂₆/ N35X₃₅, 6G4.2.5LV11N35X₃₅/ H98X₉₈, 6G4.2.5LV11S26X₂₆/H98X₉₈, and 6G4.2.5LV11S26X₂₆/ N35X₃₅/H98X₉₈, with or without any additional amino acid sequence. It will be understood that the group consisting of 6G4.2.5LV11N35X₃₅, 6G4.2.5LV11S26X₂₆, 6G4.2.5LV11H98X₉₈,
6G4.2.5LV11S26X₂₆/ N35X₃₅, 6G4.2.5LV11N35X₃₅/ H98X₉₈, 6G4.2.5LV11S26X₂₆/H98X₉₈, and 6G4.2.5LV11S26X₂₆/ N35X₃₅/H98X₉₈, is collectively referred to herein as the "group of 6G4.2.5LV11X
30 variants", and that individual members of this group are generically referred to herein as a "6G4.2.5LV11X variant." In one embodiment, the invention provides a single chain antibody fragment comprising a 6G4.2.5LV11X variant without any associated heavy chain amino acid sequence, i.e. a single chain species that makes up one half of a Fab fragment. In a preferred embodiment, the invention provides a 6G4.2.5LV11N35X₃₅ variant without any associated heavy chain amino acid sequence.

35 The invention encompasses a single chain antibody fragment comprising a variant light chain selected from the group consisting of 6G4.2.5LV11N35A, 6G4.2.5LV11S26A, 6G4.2.5LV11H98A,

6G4.2.SLV11S26A/ N35A, 6G4.2.SLV11N35A/ H98A, 6G4.2.SLV11S26A/H98A, and 6G4.2.SLV11S26A/ N35A/H98A, with or without any additional amino acid sequence. It will be understood that the group consisting of 6G4.2.SLV11N35A, 6G4.2.SLV11S26A, 6G4.2.SLV11H98A, 6G4.2.SLV11S26A/ N35A, 6G4.2.SLV11N35A/ H98A, 6G4.2.SLV11S26A/H98A, and 6G4.2.SLV11S26A/ N35A/H98A is collectively referred to herein as the "group of 6G4.2.SLV11A variants", and that individual members of this group are generically referred to herein as a "6G4.2.SLV11A variant." In one embodiment, the invention provides a single chain antibody fragment comprising a 6G4.2.SLV11A variant without any associated heavy chain amino acid sequence, i.e. a single chain species that makes up one half of a Fab fragment. In a preferred embodiment, the invention provides the 6G4.2.SLV11N35A without any associated heavy chain amino acid sequence.

Further provided herein are an antibody or antibody fragment comprising a light chain comprising a 6G4.2.SLV11X variant, and further comprising a heavy chain comprising the 6G4.2.5HV11. In a preferred embodiment, the invention provides an antibody or antibody fragment comprising a 6G4.2.SLV11N35X₃₅ variant and further comprising the 6G4.2.5HV11. In a preferred embodiment, the invention provides an antibody or antibody fragment comprising the 6G4.2.SLV11N35A and further comprising the 6G4.2.5HV11. In another preferred embodiment, the invention provides an antibody or antibody fragment comprising the 6G4.2.SLV11N35E and further comprising the 6G4.2.5HV11.

In one embodiment, the invention provides a single chain antibody fragment wherein a 6G4.2.SLV11X variant and the 6G4.2.5HV11 are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment comprises a 6G4.2.SLV11X variant joined to the 6G4.2.5HV11 by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In another embodiment, the single chain antibody fragment is a species comprising a 6G4.2.SLV11X variant joined to the 6G4.2.5HV11 by a linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In still another embodiment, the invention provides a single chain antibody fragment wherein a 6G4.2.SLV11N35X₃₅ variant and the 6G4.2.5HV11 are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment comprises a 6G4.2.SLV11N35X₃₅ variant joined to the 6G4.2.5HV11 by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In another embodiment, the single chain antibody fragment is a species comprising a 6G4.2.SLV11N35X₃₅ variant joined to the 6G4.2.5HV11 by a linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In a further embodiment, the invention provides a single chain antibody fragment wherein the 6G4.2.SLV11N35A and the 6G4.2.5HV11 are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment comprises the 6G4.2.SLV11N35A joined to the 6G4.2.5HV11 by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In another embodiment, the

single chain antibody fragment is a species comprising the 6G4.2.5LV11N35A joined to the 6G4.2.5HV11 by a linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

5 In an additional embodiment, the invention provides a single chain antibody fragment wherein the 6G4.2.5LV11N35E and the 6G4.2.5HV11 are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment comprises the 6G4.2.5LV11N35E joined to the 6G4.2.5HV11 by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In another embodiment, the single chain antibody fragment is a species comprising the 6G4.2.5LV11N35E joined to the 6G4.2.5HV11 by a
10 linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In yet another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises a 6G4.2.5LV11X variant and a second polypeptide chain comprises the 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or
15 more interchain disulfide bonds. In still another embodiment, the invention provides an antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises a 6G4.2.5LV11N35X₃₅ variant and a second polypeptide chain comprises the 6G4.2.5HV11 and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, any of the foregoing two-chain antibody fragments is selected from the group consisting of Fab, Fab', Fab'-SH, and F(ab')₂. In still
20 another preferred embodiment, the two-chain antibody fragment is a F(ab')₂ wherein one polypeptide chain comprises the 6G4.2.5LV11N35A and the second polypeptide chain comprises the 6G4.2.5HV11. In a further preferred embodiment, the antibody fragment is a Fab, Fab', Fab'-SH, or F(ab')₂ wherein one polypeptide chain comprises the 6G4.2.5LV11N35E and the second polypeptide chain comprises the 6G4.2.5HV11. A particularly preferred embodiment, the antibody fragment is the 6G4V11N35A F(ab')₂ GCN4 leucine zipper
25 species described in the Examples below. In another particularly preferred embodiment, the antibody fragment is the 6G4V11N35E F(ab')₂ GCN4 leucine zipper species described in the Examples below. In yet another particularly preferred embodiment, the antibody fragment is the 6G4V11N35E Fab described in the Examples below.

The invention also provides an antibody or antibody fragment comprising a light chain containing a
30 6G4.2.5LV11X variant and optionally further comprising a heavy chain containing the 6G4.2.5HV11, wherein the light chain, and optionally the heavy chain, is (are) fused to an additional moiety, such as additional immunoglobulin constant domain sequence. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD,
35 and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can

be obtained from Kabat *et al.*

The invention additionally provides an antibody or antibody fragment comprising a light chain containing a 6G4.2.SLV11N35X₃₅ variant and optionally further comprising a heavy chain containing the 6G4.2.SHV11, wherein the light chain, and optionally the heavy chain, is (are) fused to an additional moiety, such as additional immunoglobulin constant domain sequence. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

The invention further provides an antibody or antibody fragment comprising a light chain containing the 6G4.2.SLV11N35A and optionally further comprising a heavy chain containing the 6G4.2.SHV11, wherein the light chain, and optionally the heavy chain, is (are) fused to an additional moiety, such as additional immunoglobulin constant domain sequence. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

The invention further provides an antibody or antibody fragment comprising a light chain containing the 6G4.2.SLV11N35E and optionally further comprising a heavy chain containing the 6G4.2.SHV11, wherein the light chain, and optionally the heavy chain, is (are) fused to an additional moiety, such as additional immunoglobulin constant domain sequence. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

In a preferred embodiment, the antibody or antibody fragment comprises a light chain containing a 6G4.2.SLV11X variant, and further comprises the 6G4.2.SHV11 in a heavy chain that is fused to or contains a leucine zipper sequence. The leucine zipper can increase the affinity or production efficiency of the antibody or antibody fragment of interest. Suitable leucine zipper sequences include the jun and fos leucine zippers taught by Kostelney *et al.*, *J. Immunol.*, 148: 1547-1553 (1992) and the GCN4 leucine zipper described in the Examples below. In another preferred embodiment, the antibody or antibody fragment comprises a light chain containing the 6G4.2.SLV11N35A, and further comprises a heavy chain containing the 6G4.2.SHV11 fused to the GCN4-leucine zipper. In yet another preferred embodiment, the antibody or antibody fragment comprises a light chain containing the 6G4.2.SLV11N35E, and further comprises a heavy chain containing the 6G4.2.SHV11 fused to the GCN4 leucine zipper.

B. 6G4.2.5HV VARIANTS

The invention provides humanized antibodies and antibody fragments comprising the CDRs of a 6G4.2.5HV CDR variant. The use of a 6G4.2.5HV CDRs variant in the humanized antibodies and antibody fragments of the invention confer the advantages of higher affinity for human IL-8 and/or improved recombinant manufacturing economy.

A heavy chain variable domain comprising the CDRs of a 6G4.2.5HV CDRs variant can be humanized in conjunction with a light chain comprising the CDRs of 6G4.2.5LV or the CDRs of a 6G4.2.5LV CDRs variant, essentially as described in Section (II)(2)(A) above. In one embodiment, the invention provides a humanized antibody or antibody fragment comprising a 6G4.2.5HV CDRs variant selected from the group consisting of 6G4.2.5HV/HIS31Z₃₁, 6G4.2.5HV/H2S54Z₅₄, and 6G4.2.5HV/HIS31Z₃₁/H2S54Z₅₄. In addition, the invention provides a humanized antibody or antibody fragment comprising a 6G4.2.5HV CDRs variant selected from the group consisting of 6G4.2.5HV/HIS31A, 6G4.2.5HV/H2S54A, and 6G4.2.5HV/HIS31A/H2S54A. In particular, the 6G4.2.5HV CDRs variants can be used to construct a humanized antibody or antibody comprising the hu6G4.2.5HV/vH1-3Z as described in Section (II)(2)(A) above.

The invention additionally provides a humanized antibody or antibody fragment that comprises a heavy chain variable domain comprising the hu6G4.2.5HV/vH1-3Z, and further comprises a light chain variable domain comprising the hu6G4.2.5LV or hu6G4.2.5LV/vL1-3X.

The invention further encompasses a single chain humanized antibody fragment comprising the hu6G4.2.5HV/vH1-3Z, with or without any additional amino acid sequence. In one embodiment, the invention provides a single chain antibody fragment comprising the hu6G4.2.5HV/vH1-3Z without any associated heavy chain variable domain amino acid sequence, i.e. a single chain species that makes up one half of an Fv fragment.

In one embodiment, the invention provides a single chain humanized antibody fragment wherein the hu6G4.2.5HV/vH1-3Z and the hu6G4.2.5LV or hu6G4.2.5LV/vL1-3X are contained in a single chain polypeptide species. In a preferred embodiment, the single chain antibody fragment is a scFv species comprising the hu6G4.2.5HV/vH1-3Z joined to the hu6G4.2.5LV or hu6G4.2.5LV/vL1-3X by means of a flexible peptide linker sequence, wherein the heavy chain and light chain variable domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fv species. In another embodiment, the single chain antibody fragment is a species comprising the hu6G4.2.5HV/vH1-3Z joined to the hu6G4.2.5LV or hu6G4.2.5LV/vL1-3X by a linker that is too short to permit intramolecular pairing of the two variable domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In yet another embodiment, the invention provides a humanized antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises the hu6G4.2.5HV/vH1-3Z and a second polypeptide chain comprises the hu6G4.2.5LV or hu6G4.2.5LV/vL1-3X and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the foregoing two-chain antibody fragment is selected from the group consisting of Fab, Fab', Fab'-SH, Fv, and

F(ab')₂.

The invention also provides a humanized antibody or antibody fragment comprising a heavy chain variable domain containing the hu6G4.2.5HV/vH1-3Z and optionally further comprising a light chain variable domain containing the hu6G4.2.5LV or hu6G4.2.5LV/vL1-3X, wherein the heavy chain variable domain, and optionally the light chain variable domain, is (are) fused to an additional moiety, such as an immunoglobulin constant domain. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.*

In a preferred embodiment, the humanized antibody or antibody fragment comprises the hu6G4.2.5HV/vH1-3Z in a heavy chain that is fused to or contains a leucine zipper sequence. The leucine zipper can increase the affinity or production efficiency of the antibody or antibody fragment of interest. Suitable leucine zipper sequences include the jun and fos leucine zippers taught by Kostelney *et al.*, J. Immunol., 148: 1547-1553 (1992) and the GCN4 leucine zipper described in the Examples below.

In addition, the invention provides a humanized antibody or antibody fragment comprising a heavy chain comprising the amino acid sequence of amino acids 1-230 of the 6G4.2.5HV11 polypeptide amino acid sequence of Figs. 37A-37B (SEQ ID NO: 60) with the proviso that Ala is substituted for Ser at amino acid position 31 (hereinafter referred to as "6G4.2.5HV11S31A").

In another embodiment, the invention provides a humanized antibody or antibody fragment comprising a heavy chain comprising the amino acid sequence of amino acids 1-230 of the 6G4.2.5HV11 polypeptide amino acid sequence of Figs. 37A-37B (SEQ ID NO: 60) with the proviso that Ala is substituted for Ser at amino acid position 54 (hereinafter referred to as "6G4.2.5HV11S54A").

In yet another embodiment, the invention provides a humanized antibody or antibody fragment comprising a heavy chain comprising the amino acid sequence of amino acids 1-230 of the 6G4.2.5HV11 polypeptide amino acid sequence of Figs. 37A-37B (SEQ ID NO: 60) with the proviso that Ala is substituted for Ser at amino acid position 31 and Ala is substituted for Ser at amino acid position 54 (hereinafter referred to as "6G4.2.5HV11S31A/S54A").

Further provided herein is a humanized antibody or antibody fragment that comprises any of the light and heavy chain combinations listed in Table 1 below.

Table 1

	Heavy Chain	Light Chain
35	6G4.2.5HV11S31A	6G4.2.5LV11
	6G4.2.5HV11S31A	6G4.2.5LV11N35A
	6G4.2.5HV11S31A	6G4.2.5LV11S26A
	6G4.2.5HV11S31A	6G4.2.5LV11H98A
	6G4.2.5HV11S31A	6G4.2.5LV11S26A/N35A
40	6G4.2.5HV11S31A	6G4.2.5LV11S26A/H98A
	6G4.2.5HV11S31A	6G4.2.5LV11N35A/H98A
	6G4.2.5HV11S31A	6G4.2.5LV11S26A/N35A/H98A

	Heavy Chain	Light Chain
	6G4.2.5HV11S54A	6G4.2.5LV11
5	6G4.2.5HV11S54A	6G4.2.5LV11N35A
	6G4.2.5HV11S54A	6G4.2.5LV11S26A
	6G4.2.5HV11S54A	6G4.2.5LV11H98A
	6G4.2.5HV11S54A	6G4.2.5LV11S26A/N35A
	6G4.2.5HV11S54A	6G4.2.5LV11S26A/H98A
10	6G4.2.5HV11S54A	6G4.2.5LV11N35A/H98A
	6G4.2.5HV11S54A	6G4.2.5LV11S26A/N35A/H98A
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11N35A
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26A
15	6G4.2.5HV11S31A/S54A	6G4.2.5LV11H98A
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26A/N35A
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26A/H98A
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11N35A/H98A
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26A/N35A/H98A
20	6G4.2.5HV11S31A	6G4.2.5LV11
	6G4.2.5HV11S31A	6G4.2.5LV11N35X ₃₅
	6G4.2.5HV11S31A	6G4.2.5LV11S26X ₂₆
	6G4.2.5HV11S31A	6G4.2.5LV11H98X ₉₈
	6G4.2.5HV11S31A	6G4.2.5LV11S26X ₂₆ /N35X ₃₅
25	6G4.2.5HV11S31A	6G4.2.5LV11S26X ₂₆ /H98X ₉₈
	6G4.2.5HV11S31A	6G4.2.5LV11N35X ₃₅ /H98X ₉₈
	6G4.2.5HV11S31A	6G4.2.5LV11S26X ₂₆ /N35X ₃₅ /H98X ₉₈
	6G4.2.5HV11S54A	6G4.2.5LV11
	6G4.2.5HV11S54A	6G4.2.5LV11N35X ₃₅
30	6G4.2.5HV11S54A	6G4.2.5LV11S26X ₂₆
	6G4.2.5HV11S54A	6G4.2.5LV11H98X ₉₈
	6G4.2.5HV11S54A	6G4.2.5LV11S26X ₂₆ /N35X ₃₅
	6G4.2.5HV11S54A	6G4.2.5LV11S26X ₂₆ /H98X ₉₈
	6G4.2.5HV11S54A	6G4.2.5LV11N35X ₃₅ /H98X ₉₈
35	6G4.2.5HV11S54A	6G4.2.5LV11S26X ₂₆ /N35X ₃₅ /H98X ₉₈
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11N35X ₃₅
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26X ₂₆
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11H98X ₉₈
40	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26X ₂₆ /N35X ₃₅
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26X ₂₆ /H98X ₉₈
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11N35X ₃₅ /H98X ₉₈
	6G4.2.5HV11S31A/S54A	6G4.2.5LV11S26X ₂₆ /N35X ₃₅ /H98X ₉₈
45		

The invention encompasses a single chain humanized antibody fragment comprising a variant heavy chain selected from the group consisting of 6G4.2.5HV11S31A, 6G4.2.5HV11S54A, and 6G4.2.5HV11S31A/S54A, with or without any additional amino acid sequence. It will be understood that the group consisting of

6G4.2.5HV11S31A, 6G4.2.5HV11S54A, and 6G4.2.5HV11S31A/ S54A is collectively referred to herein as the "group of 6G4.2.5HV11A variants", and that individual members of this group are generically referred to herein as a "6G4.2.5HV11A variant." In one embodiment, the invention provides a single chain humanized antibody fragment comprising a 6G4.2.5HV11A variant without any associated light chain amino acid sequence, i.e. a single chain species that makes up one half of a Fab fragment.

Further provided herein are a humanized antibody or antibody fragment comprising a heavy chain comprising a 6G4.2.5HV11A variant, and further comprising a light chain comprising a 6G4.2.5LV11A variant or a 6G4.2.5LV11X variant. In another embodiment, the humanized antibody or antibody fragment comprises any combination of light and heavy chains listed in Table 1 above. In one embodiment, the invention provides a humanized antibody or antibody fragment comprising a 6G4.2.5HV11A variant and further comprising the 6G4.2.5LV11N35X₃₅. In a preferred embodiment, the invention provides a humanized antibody or antibody fragment comprising a 6G4.2.5HV11A variant and further comprising the 6G4.2.5LV11N35A.

In yet another embodiment, the invention provides a single chain humanized antibody fragment wherein a 6G4.2.5HV11A variant and the 6G4.2.5LV11 are contained in a single chain polypeptide species. In another embodiment, the invention provides a single chain humanized antibody fragment wherein any pair of light and heavy chains listed in Table 1 above is contained in a single chain polypeptide species. In yet another embodiment, the invention provides a single chain humanized antibody fragment wherein a 6G4.2.5HV11A variant and a 6G4.2.5LV11X variant are contained in a single chain polypeptide species. In still another embodiment, the invention provides a single chain humanized antibody fragment wherein a 6G4.2.5HV11A variant and a 6G4.2.5LV11N35X₃₅ variant are contained in a single chain polypeptide species. In an additional embodiment, the invention provides a single chain humanized antibody fragment wherein a 6G4.2.5HV11A variant and the 6G4.2.5LV11N35A variant are contained in a single chain polypeptide species.

In a preferred embodiment, the single chain humanized antibody fragment comprises a 6G4.2.5HV11A variant joined to a 6G4.2.5LV11X variant, 6G4.2.5LV11N35X₃₅ variant, 6G4.2.5LV11N35A variant, or 6G4.2.5LV11 by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In a further embodiment, the single chain humanized antibody fragment is a species comprising a 6G4.2.5HV11A variant joined to a 6G4.2.5LV11X variant, 6G4.2.5LV11N35X₃₅ variant, 6G4.2.5LV11N35A variant, or 6G4.2.5LV11 by a linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide monomer that forms a diabody upon dimerization with another monomer.

In still another embodiment, the single chain humanized antibody fragment comprises any pair of light and heavy chains listed in Table 1 above joined by means of a flexible peptide linker sequence, wherein the heavy chain and light chain domains can associate in a "dimeric" structure analogous to that formed in a two-chain Fab species. In an additional embodiment, the single chain humanized antibody fragment comprises any pair of light and heavy chains listed in Table 1 above joined by a linker that is too short to permit intramolecular pairing of complementary domains, i.e. a single chain polypeptide monomer that forms a

diabody upon dimerization with another monomer.

In yet another embodiment, the invention provides a humanized antibody fragment comprising a plurality of polypeptide chains, wherein one polypeptide chain comprises a 6G4.2.5HV11A variant and a second polypeptide chain comprises a 6G4.2.5LV11X variant, 6G4.2.5LV11N35X₃₅ variant, 6G4.2.5LV11N35A variant, or 6G4.2.5LV11, and the two polypeptide chains are covalently linked by one or more interchain disulfide bonds. In a preferred embodiment, the foregoing two-chain antibody fragment is selected from the group consisting of Fab, Fab', Fab'-SH, and F(ab')₂.

In an additional embodiment, the invention provides a two-chain humanized antibody fragment comprising any pair of heavy and light chains listed in Table 1 above, wherein each chain is contained on a separate molecule. In another embodiment, the two-chain antibody fragment comprising any pair of heavy and light chains listed in Table 1 above is selected from the group consisting of Fab, Fab', Fab'-SH, and F(ab')₂. In a preferred embodiment, the two-chain humanized antibody fragment is a F(ab')₂ comprising any pair of heavy and light chains listed in Table 1 above. In another preferred embodiment, the two-chain humanized antibody fragment is a F(ab')₂ wherein one polypeptide chain comprises a 6G4.2.5HV11A variant and the second polypeptide chain comprises the 6G4.2.5LV11N35A.

The invention also provides a humanized antibody or antibody fragment comprising a heavy chain containing a 6G4.2.5HV11A variant and optionally further comprising a light chain containing a 6G4.2.5LV11X variant, 6G4.2.5LV11N35X₃₅ variant, 6G4.2.5LV11N35A, or 6G4.2.5HV11, wherein the heavy chain, and optionally the light chain, is (are) fused to an additional moiety, such as additional immunoglobulin constant domain sequence. Constant domain sequence can be added to the heavy chain and/or light chain sequence(s) to form species with full or partial length heavy and/or light chain(s). It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. Preferably, the constant domain sequence is human in origin. Suitable human constant domain sequences can be obtained from Kabat *et al.* (supra).

In a preferred embodiment, the humanized antibody or antibody fragment comprises a 6G4.2.5HV11A variant in a heavy chain that is fused to or contains a leucine zipper sequence. The leucine zipper can increase the affinity or production efficiency of the antibody or antibody fragment of interest. Suitable leucine zipper sequences include the jun and fos leucine zippers taught by Kostelney *et al.*, J. Immunol., 148: 1547-1553 (1992) and the GCN4 leucine zipper described in the Examples below.

C. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for IL-8, the other one is for any other antigen. For example, bispecific antibodies specifically binding a IL-8 and neurotrophic factor, or two different types of IL-8 polypeptides are within the scope of the present invention.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature 305:537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829 published 13 May 1993, and in Traunecker *et al.*, EMBO J. 10:3655 (1991).

According to a different and more preferred approach, antibody-variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant-domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1), containing the site necessary for light-chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the maximum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the production of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance. In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. This asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. For further details of generating bispecific antibodies, see, for example, Suresh *et al.*, Methods in Enzymology 121:210 (1986).

According to another approach, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/00373, and EP 03089). Heteroconjugate antibodies may be

made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the HER2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147: 60 (1991).

4. Production of Humanized Anti-IL-8 6G4.2.5 Monoclonal Antibody, Antibody Fragments, and Variants

The antibodies and antibody fragments of the invention can be produced using any convenient antibody manufacturing process known in the art. Typically, the antibody or antibody fragment is made using recombinant expression systems. A multiple polypeptide chain antibody or antibody fragment species can be made in a single host cell expression system wherein the host cell produces each chain of the

antibody or antibody fragment and assembles the polypeptide chains into a multimeric structure to form the antibody or antibody fragment in vivo, followed by recovery of the antibody or antibody fragment from the host cell. For example, suitable recombinant expression systems for the production of complete antibody or antibody fragment are described in Lucas *et al.*, Nucleic Acids Res., **24**: 1774-1779 (1996). Alternatively, the separate polypeptide chains of the desired antibody or antibody fragment can be made in separate expression host cells, separately recovered from the respective host cells, and then mixed in vitro under conditions permitting the formation of the multi-subunit antibody or antibody fragment of interest. For example, U.S. Pat. No. 4,816,567 to Cabilly *et al.* and Carter *et al.*, Bio/Technology, **10**: 163-167 (1992) provide methods for recombinant production of antibody heavy and light chains in separate expression hosts followed by assembly of antibody from separate heavy and light chains in vitro.

The following discussion of recombinant expression methods applies equally to the production of single chain antibody polypeptide species and multi-subunit antibody and antibody fragment species. All recombinant procedures for the production of antibody or antibody fragment provided below shall be understood to describe: (1) manufacture of single chain antibody species as the desired end-product; (2) manufacture of multi-subunit antibody or antibody fragment species by production of all subunits in a single host cell, subunit assembly in the host cell, optionally followed by host cell secretion of the multi-subunit end-product into the culture medium, and recovery of the multi-subunit end-product from the host cell and/or culture medium; and (3) manufacture of multi-subunit antibody or antibody fragment by production of subunits in separate host cells (optionally followed by host cell secretion of subunits into the culture medium), recovery of subunits from the respective host cells and/or culture media, followed by in vitro subunit assembly to form the multi-subunit end-product. In the case of a multi-subunit antibody or antibody fragment produced in a single host cell, it will be appreciated that production of the various subunits can be effected by expression of multiple polypeptide-encoding nucleic acid sequences carried on a single vector or by expression of polypeptide-encoding nucleic acid sequences carried on multiple vectors contained in the host cell.

A. Construction of DNA Encoding Humanized 6G4.2.5 Monoclonal Antibodies, Antibody Fragments, and Variants

Following the selection of the humanized antibody or antibody fragment of the invention according to the methods described above, the practitioner can use the genetic code to design DNAs encoding the desired antibody or antibody fragment. In one embodiment, codons preferred by the expression host cell are used in the design of a DNA encoding the antibody or antibody fragment of interest. DNA encoding the desired antibody or antibody fragment can be prepared by a variety of methods known in the art. These methods include, but are not limited to, chemical synthesis by any of the methods described in Engels *et al.*, Agnew. Chem. Int. Ed. Engl., **28**: 716-734 (1989), the entire disclosure of which is incorporated herein by reference, such as the triester, phosphite, phosphoramidite and H-phosphonate methods.

A variation on the above procedures contemplates the use of gene fusions, wherein the gene(s) encoding the antibody or antibody fragment is associated, in the vector, with a gene encoding another protein or a fragment of another protein. This results in the antibody or antibody fragment being produced by the host cell as a fusion with another protein. The "other" protein is often a protein or peptide which can be secreted by the cell, making it possible to isolate and purify the desired protein from the culture medium and eliminating

the necessity of destroying the host cells which arises when the desired protein remains inside the cell. Alternatively, the fusion protein can be expressed intracellularly. It is advantageous to use fusion proteins that are highly expressed.

5 The use of gene fusions, though not essential, can facilitate the expression of heterologous proteins in *E. coli* as well as the subsequent purification of those gene products (Harris, T. J. R. in *Genetic Engineering*, Williamson, R., Ed., Academic, London, Vol. 4, p. 127(1983); Uhlen, M. & Moks, T., *Methods Enzymol.* 185:129-143 (1990)). Protein A fusions are often used because the binding of protein A, or more specifically the Z domain of protein A, to IgG provides an "affinity handle" for the purification of the fused protein (Nilsson, B. & Abrahmsen, L. *Methods Enzymol.* 185:144-161 (1990)). It has also been shown that many
10 heterologous proteins are degraded when expressed directly in *E. coli*, but are stable when expressed as fusion proteins (Marston, F. A. O., *Biochem J.* 240: 1 (1986)).

Fusion proteins can be cleaved using chemicals, such as cyanogen bromide, which cleaves at a methionine, or hydroxylamine, which cleaves between an Asn and Gly. Using standard recombinant DNA methodology, the nucleotide base pairs encoding these amino acids may be inserted just prior to the 5' end of
15 the antibody or antibody fragment gene(s).

Alternatively, one can employ proteolytic cleavage of fusion proteins, which has been recently reviewed (Carter, P. (1990) in *Protein Purification: From Molecular Mechanisms to Large-Scale Processes*, Ladisch, M. R., Willson, R. C., Painton, C. C., and Builder, S. E., eds., American Chemical Society Symposium Series No. 427, Ch 13, 181-193).

20 Proteases such as Factor Xa, thrombin, subtilisin and mutants thereof, have been successfully used to cleave fusion proteins. Typically, a peptide linker that is amenable to cleavage by the protease used is inserted between the "other" protein (e.g., the Z domain of protein A) and the protein of interest, such as humanized anti-IL-8 antibody or antibody fragment. Using recombinant DNA methodology, the nucleotide base pairs encoding the linker are inserted between the genes or gene fragments coding for the other proteins. Proteolytic
25 cleavage of the partially purified fusion protein containing the correct linker can then be carried out on either the native fusion protein, or the reduced or denatured fusion protein.

Various techniques are also available which may now be employed to produce variant humanized antibodies or antibody fragments, which encodes for additions, deletions, or changes in amino acid sequence of the resultant protein(s) relative to the parent humanized antibody or antibody fragment.

30 By way of illustration, with expression vectors encoding humanized antibody or antibody fragment in hand, site specific mutagenesis (Kunkel *et al.*, *Methods Enzymol.* 204:125-139 (1991); Carter, P., *et al.*, *Nucl. Acids. Res.* 13:4331 (1986); Zoller, M. J. *et al.*, *Nucl. Acids Res.* 10:6487 (1982)), cassette mutagenesis (Wells, J. A., *et al.*, *Gene* 34:315 (1985)), restriction selection mutagenesis (Wells, J. A., *et al.*, *Philos. Trans, R. Soc. London SerA* 317, 415 (1986)) or other known techniques may be performed on the antibody or antibody
35 fragment DNA. The variant DNA can then be used in place of the parent DNA by insertion into the aforementioned expression vectors. Growth of host bacteria containing the expression vectors with the mutant DNA allows the production of variant humanized antibodies or antibody fragments, which can be isolated as described herein.

B. Insertion of DNA into a Cloning Vehicle

The DNA encoding the antibody or antibody fragment is inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. Many vectors are available, and selection of the appropriate vector will depend on (1) whether it is to be used for DNA amplification or for DNA expression, (2) the size of the DNA to be inserted into the vector, and (3) the host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the host cell for which it is compatible. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

(i) Signal Sequence Component

In general, a signal sequence may be a component of the vector, or it may be a part of the antibody or antibody fragment DNA that is inserted into the vector. Preferably, a heterologous signal sequence selected and fused to the antibody or antibody fragment DNA such that the signal sequence in the corresponding fusion protein is recognized, transported and processed (*i.e.*, cleaved by a signal peptidase) in the host cell's protein secretion system. In the case of prokaryotic host cells, the signal sequence is selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. In a preferred embodiment, the STII signal sequence is used as described in the Examples below. For yeast secretion the native signal sequence may be substituted by, *e.g.*, the yeast invertase leader, α factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders), or acid phosphatase leader, the *C. albicans* glucoamylase leader, or the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

(ii) Origin of Replication Component

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

Most expression vectors are "shuttle" vectors, *i.e.* they are capable of replication in at least one class of organisms but can be transfected into another organism for expression. For example, a vector is cloned in *E. coli* and then the same vector is transfected into yeast or mammalian cells for expression even though it is not capable of replicating independently of the host cell chromosome.

DNA may also be amplified by insertion into the host genome. This is readily accomplished using *Bacillus* species as hosts, for example, by including in the vector a DNA sequence that is homologous to a sequence found in *Bacillus* genomic DNA. Transfection of *Bacillus* with this vector results in homologous recombination with the genome and insertion of the antibody or antibody fragment DNA.

(iii) Selection Gene Component

Expression and cloning vectors should contain a selection gene, also termed a selectable marker. This gene encodes a protein necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g. ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g. the gene encoding D-alanine racemase for *Bacilli*.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene express a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin (Southern *et al.*, J. Molec. Appl. Genet., 1: 327 (1982)), mycophenolic acid (Mulligan *et al.*, Science, 209: 1422 (1980)) or hygromycin (Sugden *et al.*, Mol. Cell. Biol., 5: 410-413 (1985)). The three examples given above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug (G418 or neomycin (geneticin), xgpt (mycophenolic acid), and hygromycin, respectively.)

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the antibody or antibody fragment nucleic acid, such as dihydrofolate reductase (DHFR) or thymidine kinase. The mammalian cell transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of having taken up the marker. Selection pressure is imposed by culturing the transformants under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes the antibody or antibody fragment. Amplification is the process by which genes in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Increased quantities of the antibody or antibody fragment are synthesized from the amplified DNA.

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity, prepared and propagated as described by Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77: 4216 (1980). The transformed cells are then exposed to increased levels of methotrexate. This leads to the synthesis of multiple copies of the DHFR gene, and, concomitantly, multiple copies of other DNA comprising the expression vectors, such as the DNA encoding the antibody or antibody fragment. This amplification technique can be used with any otherwise suitable host, e.g., ATCC No. CCL61 CHO-K1, notwithstanding the presence of endogenous DHFR if, for example, a mutant DHFR gene that is highly resistant to Mtx is employed (EP 117,060). Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding the antibody or antibody fragment, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3' phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g., kanamycin, neomycin, or G418. See U.S. Pat.

No. 4,965,199.

A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7. Stinchcomb *et al.*, Nature, **282**: 39 (1979); Kingsman *et al.*, Gene, **7**: 141 (1979); or Tschemper *et al.*, Gene, **10**: 157 (1980). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, Genetics, **85**: 12 (1977). The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* gene.

(iv) Promoter Component

Expression vectors usually contain a promoter that is recognized by the host organism and is operably linked to the antibody or antibody fragment nucleic acid. Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of a particular nucleic acid sequence, such as the antibody or antibody fragment encoding sequence, to which they are operably linked. Such promoters typically fall into two classes, inducible and constitutive. Inducible promoters are promoters that initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g. the presence or absence of a nutrient or a change in temperature. At this time a large number of promoters recognized by a variety of potential host cells are well known.

Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems (Chang *et al.*, Nature, **275**: 615 (1978); and Goeddel *et al.*, Nature, **281**: 544 (1979)), alkaline phosphatase, a tryptophan (*trp*) promoter system (Goeddel, Nucleic Acids Res., **8**: 4057 (1980) and EP 36,776) and hybrid promoters such as the *tac* promoter (deBoer *et al.*, Proc. Natl. Acad. Sci. USA, **80**: 21-25 (1983)). However, other known bacterial promoters are suitable. Their nucleotide sequences have been published, thereby enabling a skilled worker to operably ligate them to DNA encoding the antibody or antibody fragment (Siebenlist *et al.*, Cell, **20**: 269 (1980)) using linkers or adaptors to supply any required restriction sites. Promoters for use in bacterial systems also generally will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the antibody or antibody fragment.

Suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem., **255**: 2073 (1980)) or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg., **7**: 149 (1968); and Holland, Biochemistry, **17**: 4900 (1978)), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in Hitzeman *et al.*, EP 73,657A. Yeast enhancers also are advantageously used with yeast promoters.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CXCAAT region where X may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into mammalian expression vectors.

Vector driven transcription of antibody or antibody fragment encoding DNA in mammalian host cells can be controlled by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g. the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. Fiers *et al.*, Nature, 273: 113 (1978); Mulligan and Berg, Science, 209: 1422-1427 (1980); Pavlakis *et al.*, Proc. Natl. Acad. Sci. USA, 78: 7398-7402 (1981). The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. Greenaway *et al.*, Gene, 18: 355-360 (1982). A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. 4,419,446. A modification of this system is described in U.S. 4,601,978. See also Gray *et al.*, Nature, 295: 503-508 (1982) on expressing cDNA encoding immune interferon in monkey cells, Reyes *et al.*, Nature, 297: 598-601 (1982) on expression of human γ -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus, Canaani and Berg, Proc. Natl. Acad. Sci. USA, 79: 5166-5170 (1982) on expression of the human interferon β gene in cultured mouse and rabbit cells, and Gorman *et al.*, Proc. Natl. Acad. Sci. USA, 79: 6777-6781 (1982) on expression of bacterial CAT sequences in CV-1 monkey kidney cells, chicken embryo fibroblasts, Chinese hamster ovary cells, HeLa cells, and mouse NIH-3T3 cells using the Rous sarcoma virus long terminal repeat as a promoter.

(v) Enhancer Element Component

Transcription of a DNA encoding antibody or antibody fragment by higher eukaryotic host cells is often increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10-300 bp, that act on a promoter to increase its transcription. Enhancers are relatively orientation and position independent having been found 5' (Laimins *et al.*, Proc. Natl. Acad. Sci. USA, 78: 993 (1981)) and 3' (Lusky *et al.*, Mol. Cell Bio., 3: 1108 (1983)) to the transcription unit, within an intron (Banerji *et al.*, Cell, 33: 729 (1983)) as well as within the coding sequence itself (Osborne *et al.*, Mol. Cell Bio., 4: 1293 (1984)). Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, Nature, 297: 17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the

antibody or antibody fragment DNA, but is preferably located at a site 5' from the promoter.

(vi) Transcription Termination Component

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) can also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3' untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the antibody or antibody fragment. The 3' untranslated regions also include transcription termination sites.

Suitable vectors containing one or more of the above listed components and the desired coding and control sequences are constructed by standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required.

For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced by the method of Messing *et al.*, Nucleic Acids Res., **9**: 309 (1981) or by the method of Maxam *et al.*, Methods in Enzymology, **65**: 499 (1980).

Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA encoding the antibody or antibody fragment. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector.

Other methods, vectors, and host cells suitable for adaptation to the synthesis of the antibody or antibody fragment in recombinant vertebrate cell culture are described in Gething *et al.*, Nature, **293**: 620-625 (1981); Mantei *et al.*, Nature, **281**: 40-46 (1979); Levinson *et al.*, EP 117,060; and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of the IgE peptide antagonist is pRK5 (EP pub. no. 307,247) or pSV16B (PCT pub. no. WO 91/08291 published 13 June 1991).

C. Selection and Transformation of Host Cells

Suitable host cells for cloning or expressing the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes include eubacteria, such as Gram-negative or Gram-positive organisms, for example, *E. coli*, *Bacilli* such as *B. subtilis*, *Pseudomonas* species such as *P. aeruginosa*, *Salmonella typhimurium*, or *Serratia marcescens*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* 1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting. Preferably the host cell should secrete minimal amounts of proteolytic enzymes. In a preferred embodiment, the *E. coli* strain 49D6 is used as the expression host as described in the Examples below. Review articles describing the recombinant production of antibodies in bacterial host cells include Skerra *et al.*, Curr. Opinion in Immunol., **5**: 256 (1993) and Pluckthun, Immunol. Revs., **130**: 151 (1992).

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable hosts

for vectors containing antibody or antibody fragment DNA. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *S. pombe* (Beach and Nurse, Nature, 290: 140 (1981)), *Kluyveromyces lactis* (Louvencourt *et al.*, J. Bacteriol., 737 (1983)), *Yarrowia* (EP 402,226), *Pichia pastoris* (EP 183,070), *Trichoderma reesia* (EP 244,234), *Neurospora crassa* (Case *et al.*, Proc. Natl. Acad. Sci. USA, 76: 5259-5263 (1979)), and *Aspergillus* hosts such as *A. nidulans* (Ballance *et al.*, Biochem. Biophys. Res. Commun., 112: 284-289 (1983); Tilburn *et al.*, Gene, 26: 205-221 (1983); Yelton *et al.*, Proc. Natl. Acad. Sci. USA, 81: 1470-1474 (1984)) and *A. niger* (Kelly and Hynes, EMBO J., 4: 475-479 (1985)).

Host cells derived from multicellular organisms can also be used in the recombinant production of antibody or antibody fragment. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher eukaryotic cell culture is workable, whether from vertebrate or invertebrate culture. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* host cells have been identified. See, e.g., Luckow *et al.*, Bio/Technology, 6: 47-55 (1988); Miller *et al.*, in Genetic Engineering, Setlow, J.K. *et al.*, 8: 277-279 (Plenum Publishing, 1986), and Maeda *et al.*, Nature, 315: 592-594 (1985). A variety of such viral strains are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can be utilized as hosts. Typically, plant cells are transfected by incubation with certain strains of the bacterium *Agrobacterium tumefaciens*, which has been previously manipulated to contain the antibody or antibody fragment DNA. During incubation of the plant cell culture with *A. tumefaciens*, the DNA encoding antibody or antibody fragment is transferred to the plant cell host such that it is transfected, and will, under appropriate conditions, express the antibody or antibody fragment DNA. In addition, regulatory and signal sequences compatible with plant cells are available, such as the nopaline synthase promoter and polyadenylation signal sequences. Depicker *et al.*, J. Mol. Appl. Gen., 1: 561 (1982). In addition, DNA segments isolated from the upstream region of the T-DNA 780 gene are capable of activating or increasing transcription levels of plant-expressible genes in recombinant DNA-containing plant tissue. See EP 321,196 published 21 June 1989.

Vertebrate cell culture is preferred for the recombinant production of full length antibodies. The propagation of vertebrate cells in culture (tissue culture) has become a routine procedure in recent years (Tissue Culture, Academic Press, Kruse and Patterson, editors (1973)). Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen Virol., 36: 59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77: 4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23: 243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK,

ATCC CCL 34); buffal rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, Annals N.Y. Acad. Sci., 383: 44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma cell line (Hep G2). Preferred host cells are human embryonic kidney 293 and Chinese hamster ovary cells.

5 Myeloma cells that do not otherwise produce immunoglobulin protein are also useful host cells for the recombinant production of full length antibodies.

Host cells are transfected and preferably transformed with the above-described expression or cloning vectors of this invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

10 Transfection refers to the taking up of an expression vector by a host cell whether or not any coding sequences are in fact expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, CaPO₄ precipitation and electroporation. Successful transfection is generally recognized when any indication of the operation of this vector occurs within the host cell.

15 Transformation means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook *et al.*, *supra*, is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw *et al.*, Gene, 23: 315 (1983) and WO 89/05859 published 29 June 20 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method described in sections 16.30-16.37 of Sambrook *et al.*, *supra*, is preferred. General aspects of mammalian cell host system transformations have been described by Axel in U.S. 4,399,216 issued 16 August 1983. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, J. Bact., 130: 946 (1977) and Hsiao *et al.*, Proc. Natl. Acad. Sci. (USA), 76: 3829 (1979). However, other methods for introducing DNA 25 into cells such as by nuclear injection, electroporation, or by protoplast fusion may also be used.

D. Culturing the Host Cells

Prokaryotic cells used to produce the antibody or antibody fragment are cultured in suitable media as described generally in Sambrook *et al.*, *supra*.

30 The mammalian host cells used to produce the antibody or antibody fragment can be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham and Wallace, Meth. Enz., 58: 44 (1979), Barnes and Sato, Anal. Biochem., 102: 255 (1980), U.S. 4,767,704; 4,657,866; 4,927,762; or 4,560,655; WO 90/03430; WO 87/00195; U.S. Pat. Re. 30,985; or U.S. 5,122,469, the disclosures of all of 35 which are incorporated herein by reference, may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as GentamycinTM drug), trace elements (defined as inorganic compounds usually present at final concentrations in the

micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

5 The host cells referred to in this disclosure encompass cells in *in vitro* culture as well as cells that are within a host animal.

E. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77: 5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an
10 appropriately labeled probe, based on the sequences provided herein. Various labels may be employed, most commonly radioisotopes, particularly ³²P. However, other techniques may also be employed, such as using biotin-modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides,
15 fluorescers, enzymes, or the like. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

~~20 Gene expression, alternatively, may be measured by immunoological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. With immunohistochemical staining techniques, a cell sample is prepared, typically by dehydration and fixation, followed by reaction with labeled antibodies specific for the gene product, where the labels are usually visually detectable, such as enzymatic labels, fluorescent labels,~~
25 luminescent labels, and the like. A particularly sensitive staining technique suitable for use in the present invention is described by Hsu *et al.*, Am. J. Clin. Path., 75: 734-738 (1980).

F. Purification of the Antibody or Antibody Fragment

In the case of a host cell secretion system, the antibody or antibody fragment is recovered from the culture medium. Alternatively, the antibody can be produced intracellularly, or produced in
30 the periplasmic space of a bacterial host cell. If the antibody is produced intracellularly, as a first step, the host cells are lysed, and the resulting particulate debris is removed, for example, by centrifugation or ultrafiltration. Carter *et al.*, BioTechnology 10:163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min. Cell debris can be removed by
35 centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious

contaminants.

The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark *et al.*, *J. Immunol. Meth.* 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss *et al.*, *EMBO J.* 5:1567-1575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_{H3} domain, the Bakerbond ABXTM resin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SepharoseTM chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (e.g. from about 0-0.25M salt).

G. Production of Antibody Fragments

Various techniques have been developed for the production of the humanized antibody fragments of the invention, including Fab, Fab', Fab'-SH, or F(ab')₂ fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology*, 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner.

5. Uses of Anti-IL-8 Antibodies

A. Diagnostic Uses

For diagnostic applications requiring the detection or quantitation of IL-8, the antibodies or antibody fragments of the invention typically will be labeled with a detectable moiety. The detectable moiety can be any one which is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety can be a radioisotope, such as ³H, ¹⁴C, ³²P, ³⁵S, or ¹²⁵I; a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; radioactive isotopic labels, such as, e.g., ¹²⁵I, ³²P, ¹⁴C, or ³H; or an enzyme, such as alkaline phosphatase, beta-galactosidase, or

horseradish peroxidase.

Any method known in the art for separately conjugating the antibody or antibody fragment to the detectable moiety can be employed, including those methods described by Hunter *et al.*, Nature 144:945 (1962); David *et al.*, Biochemistry 13:1014 (1974); Pain *et al.*, J. Immunol. Meth. 40:219 (1981); and Nygren, J. Histochem. and Cytochem. 30:407 (1982).

The antibodies and antibody fragments of the present invention can be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. For example, see Zola, Monoclonal Antibodies: A Manual of Techniques, pp. 147-158 (CRC Press, Inc., 1987).

Competitive binding assays rely on the ability of a labeled standard (which can be a IL-8 or an immunologically reactive portion thereof) to compete with the test sample analyte (IL-8) for binding with a limited amount of antibody or antibody fragment. The amount of IL-8 in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies or antibody fragments generally are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies can conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different antigenic portion, or epitope, of the protein (IL-8) to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex (U.S. Patent No. 4,376,110). The second antibody can itself be labeled with a detectable moiety (direct sandwich assays) or can be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme (e.g., horseradish peroxidase).

IL-8 antibodies and antibody fragments also are useful for the affinity purification of IL-8 from recombinant cell culture or natural sources. For example, these antibodies can be fixed to a solid support by techniques well known in the art so as to purify IL-8 from a source such as culture supernatant or tissue.

B. Therapeutic Compositions and Administration of Anti-IL-8 Antibody

The humanized anti-IL-8 antibodies and antibody fragments of the invention are useful in the treatment of inflammatory disorders, including inflammations of the lung, such as adult respiratory distress syndrome (ARDS) and any stage of acute lung injury in the pathogenesis of ARDS described in Bernard *et al.*, Am. J. Respir. Crit. Care Med., 149: 818-824 (1994), bacterial pneumonia, hypovolemic shock, ischemic reperfusion disorders such as surgical tissue reperfusion injury, myocardial ischemic conditions such as myocardial infarction, reperfusion after cardiac surgery, cardiac arrest, and constriction after percutaneous transluminal coronary angioplasty, inflammatory bowel disorders such as ulcerative colitis, and autoimmune diseases such as rheumatoid arthritis. In addition, the humanized anti-IL-8 antibodies and antibody fragments of the invention are useful in the treatment of asthmatic diseases, such as allergic asthma.

Therapeutic formulations of the humanized anti-IL-8 antibodies and antibody fragments are prepared for storage by mixing the antibody or antibody fragment having the desired degree of purity with optional

physiologically acceptable carriers, excipients, or stabilizers (Remington's Pharmaceutical Sciences, supra), in the form of lyophilized cake or aqueous solutions. Acceptable carriers, excipients or stabilizers are n ntotoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and ther organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Pluronic or polyethylene glycol (PEG).

10 The humanized anti-IL-8 mAb or antibody fragment to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. The humanized anti-IL-8 mAb or antibody fragment ordinarily will be stored in lyophilized form or in solution.

15 Therapeutic humanized anti-IL-8 mAb or antibody fragment compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

20 The route of humanized anti-IL-8 mAb or antibody fragment administration is in accord with known methods, e.g., inhalation, injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial, or intralesional routes, by enema or suppository, or by sustained release systems as noted below. Preferably the antibody is given systemically or at a site of inflammation.

25 In one embodiment, the invention provides for the treatment of asthmatic diseases by administration of humanized anti-IL-8 mAb or antibody fragment to the respiratory tract. The invention contemplates formulations comprising humanized anti-IL-8 mAb or antibody fragment for use in a wide variety of devices that are designed for the delivery of pharmaceutical compositions and therapeutic formulations to the respiratory tract. In one aspect, humanized anti-IL-8 mAb or antibody fragment is administered in aerosolized or inhaled form. The humanized anti-IL-8 mAb or antibody fragment, combined with a dispersing agent, or dispersant, can be administered in an aerosol formulation as a dry powder or in a solution or suspension with a diluent.

30 Suitable dispersing agents are well known in the art, and include but are not limited to surfactants and the like. Surfactants are generally used in the art to reduce surface induced aggregation of protein caused by atomization of the solution forming the liquid aerosol. Examples of such surfactants include polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitan fatty acid esters. Amounts of surfactants used will vary, being generally within the range of about 0.001 to 4% by weight of the formulation. In a specific aspect, the surfactant is polyoxyethylene sorbitan monooleate or sorbitan trioleate.

35 Liquid aerosol formulations contain the humanized anti-IL-8 mAb or antibody fragment and a dispersing agent in a physiologically acceptable diluent. The dry powder formulations of the invention consist of a finely divided solid form of the humanized anti-IL-8 mAb or antibody fragment and a dispersing agent, and optionally a bulking agent, such as lactose, sorbitol, sucrose, r mannitol, and the like, to facilitate dispersal of the powder. With either the liquid or dry powder aerosol formulation, the formulations must be

aerosolized. It must be broken down into liquid or solid particles in order to ensure that the aerosolized dose actually reaches the bronchii and/or alveoli, as desired. For example, in the methods for treatment of asthma provided herein, it is preferable to deliver aerosolized humanized anti-IL-8 mAb or antibody fragment to the bronchii. In other embodiments, such as the present methods for treating ARDS and any stage of acute lung injury in the pathogenesis of ARDS, it is preferable to deliver aerosolized humanized anti-IL-8 mAb or antibody fragment to the alveoli. In general, the mass median dynamic diameter will be 5 micrometers (μm) or less to ensure that the drug particles reach the lung bronchii or alveoli (Wearly, L.L., 1991, Crit. Rev. in Ther. Drug Carrier Systems, 8:333).

With regard to construction of the delivery device, any form of aerosolization known in the art, including but not limited to nebulization, atomization or pump aerosolization of a liquid formulation, and aerosolization of a dry powder formulation, can be used in the practice of the invention. A delivery device that is uniquely designed for administration of solid formulations is envisioned. Often, the aerosolization of a liquid or a dry powder formulation will require a propellant. The propellant can be any propellant generally used in the art. Examples of useful propellants include chlorofluorocarbons, hydrofluorocarbons, hydrochlorofluorocarbons, and hydrocarbons, including trifluoromethane, dichlorofluoromethane, dichlorotetrafluoroethanol, and 1,1,1,2-tetrafluoroethane, and combinations thereof.

In a preferred aspect of the invention, the device for aerosolization is a metered dose inhaler. A metered dose inhaler provides a specific dosage when administered, rather than a variable dose depending on administration. Such a metered dose inhaler can be used with either a liquid or a dry powder aerosol formulation.

Systems of aerosol delivery, such as the pressurized metered dose inhaler and the dry powder inhaler are disclosed in Newman, S.P., *Aerosols and the Lung*, Clarke, S.W. and Davia, D. editors, pp.197-22 and can be used in connection with the present invention.

Sustained release systems can be used in the practice of the methods of the invention. Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, Biopolymers 22:547 (1983)), poly (2-hydroxyethyl-methacrylate) (Langer *et al.*, J. Biomed. Mater. Res. 15:167 (1981) and Langer, Chem. Tech. 12:98 (1982)), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release humanized anti-IL-8 antibody or antibody fragment compositions also include liposomally entrapped antibody or antibody fragment. Liposomes containing an antibody or antibody fragment are prepared by methods known per se: DE 3,218,121; Epstein *et al.*, Proc. Natl. Acad. Sci. U.S.A. 82:3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. U.S.A. 77:4030 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mole percent cholesterol, the selected proportion being adjusted for the most efficacious antibody or antibody fragment therapy.

An "effective amount" of the humanized anti-IL-8 antibody or antibody fragment to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, and the

condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. Typically, the clinician will administer the humanized anti-IL-8 antibody or antibody fragment until a dosage is reached that achieves the desired effect. The progress of this therapy is easily monitored by conventional assays.

5 In the treatment and prevention of an inflammatory disorder or asthmatic disorder with a humanized anti-IL-8 antibody or antibody fragment of the invention, the antibody composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the antibody, the particular type of antibody,
10 the method of administration, the scheduling of administration, and other factors known to medical practitioners. The "therapeutically effective amount" of antibody to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the inflammatory disorder, including treating acute or chronic respiratory diseases and reducing inflammatory responses. Such amount is preferably below the amount that is toxic to the host or renders the host significantly more
15 susceptible to infections.

As a general proposition, the initial pharmaceutically effective amount of the antibody or antibody fragment administered parenterally per dose will be in the range of about 0.1 to 50 mg/kg of patient body weight per day, with the typical initial range of antibody used being 0.3 to 20 mg/kg/day, more preferably 0.3 to 15 mg/kg/day.

20 In one embodiment, using systemic administration, the initial pharmaceutically effective amount will be in the range of about 2 to 5 mg/kg/day.

For methods of the invention using administration by inhalation, the initial pharmaceutically effective amount will be in the range of about 1 microgram (μ g)/kg/day to 100 mg/kg/day.

The invention provides for both prophylactic and therapeutic treatment of inflammatory disorders.
25 Without intending to limit the methods of the invention to a particular mechanism of action or a particular disease intervention strategy, it is noted that in some indications it is beneficial to treat the disease in question prior to or early on in the stage of the underlying disease that involves neutrophil activation, recruitment and infiltration at sites of inflammation. Accordingly, it may be advantageous to utilize the humanized anti-IL-8 mAb or antibody fragment in a prophylactic treatment regimen for an inflammatory disease indication in order
30 to attenuate or eliminate a pathogenic neutrophil response that may or will arise during the course of the disease.

In patients at risk of developing acute lung injury with possible or likely progression to ARDS, it is desirable to employ a prophylactic course of treatment in order to ameliorate or prevent the deterioration of lung function and the pathogenesis of associated disease sequelae (which may greatly increase patient
35 morbidity and mortality) prior to the onset of such conditions. Certain biological parameters, such as IL-8 levels in bronchial alveolar lavage (BAL) fluid and ferritin levels in serum, can be used for prognosis of acute lung injury and ARDS in patients who are predisposed to such disease progression, i.e. patients suffering from diseases or other insults that commonly precipitate acute lung injury and ARDS, such as aspiration, diffuse pulmonary infection, near-drowning, toxic inhalation, lung contusion, multiple trauma, pancreatitis, perforated

bowel, sepsis, and the like. In one embodiment, acute lung injury and ARDS at-risk patients presenting BAL fluid IL-8 concentrations of at or above 0.2 ng/ml are selected for prophylactic treatment according to the methods of the invention. Any suitable method for assay of IL-8 in patient BAL fluid may be employed, such as the method described in Donnelly et al., Lancet, 341: 643-647 (1993).

5 In another embodiment, acute lung injury/ARDS at-risk female and male patients presenting ferritin serum concentrations of at or above 270 ng/ml and 680 ng/ml, respectively, are selected for prophylactic treatment according to the methods of the invention. Any suitable method for assay of ferritin in patient serum may be employed, such as the method described in U.S. Pat. No. 5,679,532 for "Serum Ferritin as a Predictor of the Acute Respiratory Distress Syndrome" to Repine issued on October 21, 1997.

10 In patients presenting ischemic conditions or undergoing surgical procedures that generate ischemic conditions in tissue and concomitant risk of tissue injury upon reperfusion, it is desirable to employ a course of treatment wherein the humanized anti-IL-8 mAb or antibody fragment is administered to the patient prior to the reperfusion of ischemic tissue, or prior to or as soon as possible after the onset of an inflammatory response following reperfusion of ischemic tissue. In the patients presenting acute myocardial infarction, for example, it
15 is advantageous to employ a course of treatment wherein the humanized anti-IL-8 mAb or antibody fragment is administered to the patient prior to or concomitant with recanalization therapy, including pharmaceutical recanalization therapies such as the administration of tissue plasminogen activators, streptokinase, or other thrombolytic drugs with or without anti-clotting agents such as platelet-fibrin binding antagonists (e.g. anti-IIbIIIa integrin antibody), blood thinning agents such as heparin, or other anti-reocclusion agents such as
20 aspirin, and the like, and including mechanical recanalization therapies such as percutaneous transluminal coronary angioplasty, or wherein the humanized anti-IL-8 mAb or antibody fragment is administered to the patient prior to or as soon as possible after the onset of an inflammatory response following reperfusion of ischemic myocardium. In yet another embodiment, the humanized anti-IL-8 mAb or antibody fragment of the invention can be employed in the methods of treating acute myocardial infarction with anti-IL-8 antibody
25 described in WO 97/40215 published October 30, 1997.

The invention provides for both prophylactic and therapeutic treatment of asthma with humanized anti-IL-8 mAb and antibody fragment. In the case of prophylactic treatment for allergic asthma with the antibodies or antibody fragments of the invention, it is desirable to administer about 0.1 to 10 mg/kg of the antibody agent to the patient up to about 24 hours prior to anticipated exposure to allergen or prior to onset of
30 allergic asthma. In the case of therapeutic treatment for acute asthma, including allergic asthma, it is desirable to treat the asthmatic patient as early as possible following onset of an asthma attack. In one embodiment, an episode of acute asthma is treated within 24 hours of the onset of symptoms by administration of about 0.1 to 10 mg/kg of an anti-IL-8 antibody agent. However, it will be appreciated that the methods of the invention can be used to ameliorate symptoms at any point in the pathogenesis of asthmatic disease. Additionally, the
35 methods of the invention can be used to alleviate symptoms of chronic asthmatic conditions.

The antibody or antibody fragment need not be, but is optionally formulated with one or more agents currently used to prevent or treat the inflammatory disorder or asthmatic disease in question. For example, in rheumatoid arthritis, the antibody can be given in conjunction with a glucocorticosteroid. In the case of treating asthmatic diseases with anti-IL-8 antibody or antibody fragment, the invention contemplates the

coadministration of antibody or antibody fragment and one or more additional agents useful in treating asthma, such as bronchodilators, antihistamines, epinephrine, and the like. The effective amount of such other agents depends on the amount of antibody or antibody fragment present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

The following examples are offered by way of illustration and not by way of limitation. The disclosures of all references cited in the specification, and the disclosures of all citations in such references, are expressly incorporated herein by reference.

EXAMPLES

A. GENERATION AND CHARACTERIZATION OF MONOCLONAL ANTIBODIES AGAINST HUMAN IL-8

Balb/c mice were immunized in each hind footpad or intraperitoneally with 10 µg of recombinant human IL-8 (produced as a fusion of (ser-IL-8)₇₂ with ubiquitin (Hebert *et al.* J. Immunology 145:3033-3040 (1990)); IL-8 is available commercially from PeproTech, Inc., Rocky Hill, NJ) resuspended in MPL/TDM (Ribi Immunochem. Research Inc., Hamilton, MT) and boosted twice with the same amount of IL-8. In these experiments, "IL-8" is intended to mean (ser-IL-8)₇₂ unless otherwise specified. A final boost of 10 µg of IL-8 was given 3 days before the fusion. Spleen cells or popliteal lymph node cells were fused with mouse myeloma P3X63Ag8U.1 (ATCC CRL1597), a non-secreting clone of the myeloma P3X63Ag8, using 35% polyethylene glycol as described before. Ten days after the fusion, culture supernatant was screened for the presence of monoclonal antibodies to IL-8 by ELISA.

The ELISA was performed as follows. Nunc 96-well immunoplates (Flow Lab, McLean, VA) were coated with 50 µl/well of 2 µg/ml IL-8 in phosphate-buffered saline (PBS) overnight at 4°C. The remaining steps were carried out at room temperature. Non-specific binding sites were blocked with 0.5% bovine serum albumin (BSA) for 1 hour (hr). Plates were then incubated with 50 µl/well of hybridoma culture supernatants from 672 growing parental fusion wells for 1 hr, followed by the incubation with 50 µl/well of 1:1000 dilution of a 1 mg/ml stock solution of alkaline phosphatase-conjugated goat anti-mouse Ig (Tago Co., Foster City, CA) for 1 hr. The level of enzyme-linked antibody bound to the plate was determined by the addition of 100 µl/well of 0.5 mg/ml of p-nitrophenyl phosphate in sodium bicarbonate buffer, pH 9.6. The color reaction was measured at 405 nm with an ELISA plate reader (Titertrek Multiscan, Flow Lab, McLean, VA). Between each step, plates were washed three times in PBS containing 0.05% Tween 20.

Culture supernatants which promoted 4-fold more binding of IL-8 than did control medium were selected as positives. According to this criterion, 16 of 672 growing parental fusion wells (2%) were positive. These positive hybridoma cell lines were cloned at least twice by using the limiting dilution technique.

Seven of the positive hybridomas were further characterized as follows. The isotypes of the monoclonal antibodies were determined by coating Nunc 96-well immunoplates (Flow Lab, McLean, VA) with IL-8 overnight, blocking with BSA, incubating with culture supernatants followed by the addition of predetermined amount of isotype-specific alkaline phosphatase-conjugated goat anti-mouse Ig (Fisher Biotech, Pittsburgh, PA). The level of conjugated antibodies bound to the plate was determined by the addition of r-

nitrophenyl phosphate as described above.

All the monoclonal antibodies tested belonged to either IgG₁ or IgG₂ immunoglobulin isotype. Ascites fluid containing these monoclonal antibodies had antibody titers in the range of 10,000 to 100,000 as determined by the reciprocal of the dilution factor which gave 50% of the maximum binding in the ELISA.

- 5 To assess whether these monoclonal antibodies bound to the same epitopes, a competitive binding ELISA was performed. At a ratio of biotinylated mAb to unlabeled mAb of 1:100, the binding of biotinylated mAb 5.12.14 was significantly inhibited by its homologous mAb but not by mAb 4.1.3, while the binding of biotinylated mAb 4.1.3 was inhibited by mAb 4.1.3 but not by mAb 5.12.14. Monoclonal antibody 5.2.3 behaved similarly to mAb 4.1.3, while monoclonal antibodies 4.8 and 12.3.9 were similar to mAb 5.12.14.
- 10 Thus, mAb 4.1.3 and mAb 5.2.3 bind to a different epitope(s) than the epitope recognized by monoclonal antibodies 12.3.9, 4.8 and 5.12.14.

Immunodot blot analysis was performed to assess antibody reactivity to IL-8 immobilized on nitrocellulose paper. All seven antibodies recognized IL-8 immobilized on paper, whereas a control mouse IgG antibody did not.

- 15 The ability of these monoclonal antibodies to capture soluble ¹²⁵I-IL-8 was assessed by a radioimmune precipitation test (RIP). Briefly, tracer ¹²⁵I-IL-8 (4 x 10⁴ cpm) was incubated with various dilutions of the monoclonal anti-IL-8 antibodies in 0.2 ml of PBS containing 0.5% BSA and 0.05% Tween 20 (assay buffer) for 1 hr at room temperature. One hundred microliters of a predetermined concentration of goat anti-mouse Ig antisera (Pel-Freez, Rogers, AR) were added and the mixture was incubated at room temperature
- 20 for 1 hr. Immune complexes were precipitated by the addition of 0.5 ml of 6% polyethylene glycol (M.W. 8000) kept at 4°C. After centrifugation at 2,000 x g for 20 min at 4°C, the supernatant was removed by aspiration and the radioactivity remaining in the pellet was counted in a gamma counter. Percent specific binding was calculated as (precipitated cpm - background cpm) / (total cpm - background cpm). Monoclonal antibodies 4.1.3, 5.2.3, 4.8, 5.12.14 and 12.3.9 captured ¹²⁵I-IL-8 very efficiently, while antibodies 9.2.4 and
- 25 8.9.1 were not able to capture soluble ¹²⁵I-IL-8 in the RIP even though they could bind to IL-8 coated onto ELISA plates (Table I).

- The dissociation constants of these monoclonal antibodies were determined using a competitive binding RIP assay. Briefly, competitive inhibition of the binding each antibody to ¹²⁵I-IL-8 (20,000-40,000 cpm per assay) by various amounts of unlabeled IL-8 was determined by the RIP described above. The
- 30 dissociation constant (affinity) of each mAb was determined by using Scatchard plot analysis (Munson, *et al.*, Anal. Biochem. 107:220 (1980)) as provided in the VersaTerm-PRO computer program (Synergy Software, Reading, PA). The K_d's of these monoclonal antibodies (with the exception of 9.2.4. and 8.9.1) were in the range from 2 x 10⁻⁸ to 3 x 10⁻¹⁰ M. Monoclonal antibody 5.12.14 with a K_d of 3 x 10⁻¹⁰ M showed the highest affinity among all the monoclonal antibodies tested (Table 3).

Table 3. Characterization of Anti-IL-8 Monoclonal Antibodies

Antibody	%Specific Binding to IL-8	$K_d(M)$	isotype	pi
4.1.3	58	2×10^{-9}	IgG ₁	4.3-6.1
5.2.3	34	2×10^{-8}	IgG ₁	5.2-5.6
9.2.4	1	-	IgG ₁	7.0-7.5
8.9.1	2	-	IgG ₁	6.8-7.6
4.8	62	3×10^{-8}	IgG _{2a}	6.1-7.1
5.12.14	98	3×10^{-10}	IgG _{2a}	6.2-7.4
12.3.9	86	2×10^{-9}	IgG _{2a}	6.5-7.1

To assess the ability of these monoclonal antibodies to neutralize IL-8 activity, the amount of ^{125}I -IL-8 bound to human neutrophils in the presence of various amounts of culture supernatants and purified monoclonal antibodies was measured. Neutrophils were prepared by using Mono-Poly Resolving Medium (M-PRM) (Flow Lab. Inc., McLean, VA). Briefly fresh, heparinized human blood was loaded onto M-PRM at a ratio of blood to medium, 3.5:3.0, and centrifuged at 300 x g for 30 min at room temperature. Neutrophils enriched at the middle layer were collected and washed once in PBS. Such a preparation routinely contained greater than 95% neutrophils according to the Wright's Giemsa staining. The receptor binding assay was done as follows. 50 μ l of ^{125}I -IL-8 (5 ng/ml) was incubated with 50 μ l of unlabeled IL-8 (100 μ g/ml) or monoclonal antibodies in PBS containing 0.1% BSA for 30 min at room temperature. The mixture was then incubated with 100 μ l of neutrophils (10⁷ cells/ml) for 15 min at 37°C. The ^{125}I -IL-8 bound was separated from the unbound material by loading mixtures onto 0.4 ml of PBS containing 20% sucrose and 0.1% BSA and by centrifugation at 300 x g for 15 min. The supernatant was removed by aspiration and the radioactivity associated with the

pellet was counted in a gamma counter.

Monoclonal antibodies 4.1.3, 5.2.3, 4.8, 5.12.14, and 12.3.9 inhibited greater than 85% of the binding of IL-8 to human neutrophils at a 1:25 molar ratio of IL-8 to mAb. On the other hand, monoclonal antibodies 9.2.4 and 8.9.1 appeared to enhance the binding of IL-8 to its receptors on human neutrophils. Since a control mouse IgG also enhanced the binding of IL-8 on neutrophils, the enhancement of IL-8 binding to its receptors by mAb 9.2.4 and 8.9.1 appears to be nonspecific. Thus, monoclonal antibodies, 4.1.3, 5.1.3, 4.8, 5.12.14, and 12.3.9 are potential neutralizing monoclonal antibodies while monoclonal antibodies 8.9.1 and 9.2.4 are non-neutralizing monoclonal antibodies.

The ability of the anti-IL-8 antibodies to block neutrophil chemotaxis induced by IL-8 was tested as follows. Neutrophil chemotaxis induced by IL-8 was determined using a Boyden chamber method (Larsen, *et al. Science* 243:1464 (1989)). One hundred μ l of human neutrophils (10^6 cells/ml) resuspended in RPMI containing 0.1% BSA were placed in the upper chamber and 29 μ l of the IL-8 (20 nM) with or without monoclonal antibodies were placed in the lower chamber. Cells were incubated for 1 hr at 37°C. Neutrophils migrated into the lower chamber were stained with Wright's Giemsa stain and counted under the microscope (100x magnification). Approximately 10 different fields per experimental group were examined. Neutralizing monoclonal antibodies 5.12.14 and 4.1.3 blocked almost 70% of the neutrophil chemotactic activity of IL-8 at 1:10 ratio of IL-8 to mAb.

The isoelectric focusing (IEF) pattern of each mAb was determined by applying purified antibodies on an IEF polyacrylamide gel (pH 3-9, Pharmacia) using the Fast gel system (Pharmacia, Piscataway, NJ). The IEF gel was pretreated with pharmalyte containing 1% Triton X100 (Sigma, St. Louis, MO) for 10 min before loading the samples. The IEF pattern was visualized by silver staining according to the instructions from the manufacturer. All of the monoclonal antibodies had different IEF patterns, confirming that they originated from different clones. The pI values for the antibodies are listed in Table 3.

All these monoclonal antibodies bound equally well to both (ala-IL-8)₇₇ and (ser-IL-8)₇₂ forms of IL-8. Because IL-8 has greater than 30% sequence homology with certain other members of the platelet factor 4 (PF4) family of inflammatory cytokines such as β -TG (Van Damme *et al.*, Eur. J. Biochem. 181:337(1989); Tanaka *et al.*, FEB 236(2):467 (1988)) and PF4 (Deuel *et al.*, Proc. Natl. Acad. Sci. U.S.A. 74:2256 (1977)), they were tested for possible cross reactivity to β -TG and PF4, as well as to another neutrophil activating factor, C5a. No detectable binding to any of these proteins was observed, with the exception of mAb 4.1.3, which had a slight cross reactivity to β -TG.

One of the antibodies, mAb 5.12.14, was further studied to determine whether it could block the IL-8 mediated release of elastase by neutrophils. Briefly, human neutrophils were resuspended in Hanks balanced salt solution (Gibco, Grand Island, NY) containing 1.0% BSA, Fraction V (Sigma, St. Louis, MO), 2 mg/ml alpha-D-glucose (Sigma), 4.2 mM sodium bicarbonate (Sigma) and 0.01 M HEPES, pH 7.1 (JRH Bioscience, Lenexa, KS). A stock of cytochalasin B (Sigma) was prepared (5 mg/ml in dimethylsulfoxide (Sigma) and stored at 2-8°C. Cytochalasin B was added to the neutrophil preparation to produce a final concentration of 5 μ g/ml, and incubated for 15 min at 37°C. Human IL-8 was incubated with mAb 5.12.14 (20 μ l), or a negative control antibody, in 1 ml polypropylene tubes (DBM Scientific, San Fernando, CA) for 30 min at 37°C. The

final assay concentrations of IL-8 were 50 and 500 nM. The monoclonal antibodies were diluted to produce the following ratios (IL-8:Mab): 1:50, 1:10, 1:2, 1:1, and 1:0.25. Cytochalasin B-treated neutrophils were added (100 μ l/tube) and incubated for 2 hours at 25°C. The tubes were centrifuged (210 X g, 2-8°C) for 10 min, and supernatants were transferred to 96 well tissue culture plates (30 μ l/well). Elastase substrate stock, 10 mM methoxysuccinyl-alanyl-alanyl-propyl-valyl-p-nitroanilide (Calbiochem, La Jolla, CA) in DMSO was prepared and stored at 2-8°C. Elastase substrate solution (1.2 mM substrate, 1.2 M NaCl (Mallinckrodt, Paris, Kentucky), 0.12 M HEPES pH 7.2 in distilled water) was added (170 μ l/well) to the supernatants and incubated for 0.5 to 2 hours at 37°C (until control O.D. of 1.0 was reached). Absorbance was measured at 405 nm (SLT 340 ATTC plate reader, SLT Lab Instruments, Austria).

The results are shown in Figure 1. At a 1:1 ratio of IL-8 to mAb 5.12.14, the antibody was able to effectively block the release of elastase from neutrophils.

The hybridoma producing antibody 5.12.14 was deposited on February 15, 1993 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, U.S.A. (ATCC) and assigned ATCC Accession No. HB 11553.

B. GENERATION AND CHARACTERIZATION OF MONOCLONAL ANTIBODIES AGAINST RABBIT IL-8

Antibodies against rabbit IL-8 were generated in essentially the same process as anti-human IL-8 antibodies using rabbit IL-8 as immunogen (kindly provided by C. Broaddus; see also Yoshimura *et al.*, Immunol. 146:3483 (1991)). The antibody was characterized as described above for binding to other cytokines coated onto ELISA plates; no measurable binding was found to MGSA, fMLP, C5a, b-TG, TNF, PF4, or IL-1.

The hybridoma producing antibody 6G4.2.5 was deposited on September 28, 1994, with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, U.S.A. (ATCC) and assigned ATCC Accession No. HB 11722.

Recombinant human-murine chimeric Fabs for 5.12.14 and 6G4.2.5 were constructed as described below. A chimeric 6G4.2.5 Fab is compared with a chimeric 5.12.14 Fab in detail below.

1. INHIBITION OF IL-8 BINDING TO HUMAN NEUTROPHILS BY 5.12.14-FAB AND 6G4.2.5-FAB

The ability of the two chimeric Fabs, 5.12.14-Fab and 6G4.2.5-Fab, to efficiently bind IL-8 and prevent IL-8 from binding to IL-8 receptors on human neutrophils was determined by performing a competition binding assay which allows the calculation of the IC₅₀ - concentration required to achieve 50% inhibition of IL-8 binding.

Human neutrophils (5×10^5) were incubated for 1 hour at 4°C with 0.5nM ¹²⁵I-IL-8 in the presence of various concentrations (0 to 300 nM) of 5.12.14-Fab, 6G4.2.5-Fab, an isotype control (4D5-Fab) or unlabeled IL-8. After the incubation, the unbound ¹²⁵I-IL-8 was removed by centrifugation through a solution of 20% sucrose and 0.1% bovine serum albumin in phosphate buffered saline and the amount of ¹²⁵I-IL-8 bound to the cells was determined by counting the cell pellets in a gamma counter. Figure 2 demonstrates the inhibition of ¹²⁵I-IL-8 binding to neutrophils by unlabeled IL-8. Figure 3 demonstrates that a negative isotype

matched Fab does not inhibit the binding of ^{125}I -IL-8 to human neutrophils. Both the anti-IL-8 Fabs, 5.12.14 Fab (Figure 4) and 6G4.25 Fab (Figure 5) were able to inhibit the binding of ^{125}I -IL-8 to human neutrophils with an average IC_{50} of 1.6 nM and 7.5 nM, respectively.

2. INHIBITION OF IL-8-MEDIATED NEUTROPHIL CHEMOTAXIS BY 5.12.14-FAB AND 6G4.2.5-FAB

Human neutrophils were isolated, counted and resuspended at 5×10^6 cells/ml in Hank's balanced salt solution (abbreviated HBSS; without calcium and magnesium) with 0.1% bovine serum albumin. The neutrophils were labeled by adding calcein AM (Molecular Probe, Eugene, OR) at a final concentration of 2.0 μM . Following a 30 minute incubation at 37°C , cells were washed twice with HBSS-BSA and resuspended at 5×10^6 cells/ml.

Chemotaxis experiments were carried out in a Neuro Probe (Cabin John, MD) 96-well chamber, model MBB96. Experimental samples (buffer only control, IL-8 alone or IL-8 + Fabs) were loaded in a Polyfiltronics 96-well View plate (Neuro Probe Inc.) placed in the lower chamber. 100 μl of the calcein AM-labeled neutrophils were added to the upper chambers and allowed to migrate through a 5 micrometer porosity PVP free polycarbonate framed filter (Neuro Probe Inc.) toward the bottom chamber sample. The chemotaxis apparatus was then incubated for 40 to 60 minutes at 37°C with 5% CO_2 . At the end of the incubation, neutrophils remaining in the upper chamber were aspirated and upper chambers were washed three times with PBS. Then the polycarbonate filter was removed, non-migrating cells were wiped off with a squeegee wetted with PBS, and the filter was air dried for 15 minutes.

The relative number of neutrophils migrating through the filter (Neutrophil migration index) was determined by measuring fluorescence intensity of the filter and the fluorescence intensity of the contents of the lower chamber and adding the two values together. Fluorescence intensity was measured with a CytoFluor 2300 fluorescent plate reader (Millipore Corp. Bedford, MA) configured to read a Corning 96-well plate using the 485-20 nm excitation filter and a 530-25 emission filter, with the sensitivity set at 3.

The results are shown in Figures 6 and 7. Figure 6 demonstrates the inhibition of human IL-8 mediated neutrophil chemotaxis by chimeric 6G4.2.5 and 5.12.14 Fabs. Figure 7 demonstrates the relative abilities of chimeric 6G4.2.5 and 5.12.14 Fabs to inhibit rabbit IL-8 mediated neutrophil chemotaxis.

3. INHIBITION OF IL-8-MEDIATED NEUTROPHIL ELASTASE RELEASE BY VARIOUS CONCENTRATIONS OF 6G4.2.5 AND 5.12.14 FABS

Blood was drawn from healthy male donors into heparinized syringes. Neutrophils were isolated by dextran sedimentation, centrifugation over Lymphocyte Separation Medium (Organon Teknika, Durham, NC), and hypotonic lysis of contaminating red blood cells as described by Berman *et al.* (*J. Cell Biochem.* 52:183 (1993)). The final neutrophil pellet was suspended at a concentration of 1×10^7 cells/ml in assay buffer, which consisted of Hanks Balanced Salt Solution (GIBCO, Grand Island, NY) supplemented with 1.0% BSA (fraction V, Sigma, St. Louis, MO), 2 mg/ml glucose, 4.2 mM sodium bicarbonate, and 0.01 M HEPES, pH 7.2. The neutrophils were stored at 4°C for not longer than 1 hr.

IL-8 (10 μl) was mixed with anti-IL-8 Fab, an isotype control Fab, or buffer (20 μl) in 1 ml

polypropylene tubes and incubated in a 37°C water bath for 30 min. IL-8 was used at final concentrations ranging from 0.01 to 1000 nM in dose response studies (Figure 8) and at a final concentration of 100 nM in the experiments addressing the effects of the Fabs on elastase release (Figures 9 and 10). Fab concentrations ranged from approximately 20 nM to 300 nM, resulting in Fab:IL-8 molar ratios of 0.2:1 to 3:1. Cytochalasin B (Sigma) was added to the neutrophil suspension at a concentration of 5 µg/ml (using a 5mg/ml stock solution made up in DMSO), and the cells were incubated for 15 min in a 37°C water bath. Cytochalasin B-treated neutrophils (100 µl) were then added to the IL-8/Fab mixtures. After a 3 hr incubation at room temperature, the neutrophils were pelleted by centrifugation (200 x g for 5 min), and aliquots of the cell-free supernatants were transferred to 96 well plates (30 µl/well). The elastase substrate, methoxysuccinyl-alanyl-alanyl-prolyl-valyl-p-nitroanilide (Calbiochem, La Jolla, CA), was prepared as a 10 mM stock solution in DMSO and stored at 4°C. Elastase substrate working solution was prepared just prior to use (1.2 mM elastase substrate, 1.2 M NaCl, 0.12 M HEPES, pH 7.2), and 170 µl was added to each sample-containing well. The plates were placed in a 37°C tissue culture incubator for 30 min or until an optical density reading for the positive controls reached at least 1.0. Absorbance was measured at 405 nm using an SLT 340 plate reader (SLT Lab Instruments, Austria).

Figure 9 demonstrates the ability of the chimeric anti-IL-8 Fabs to inhibit elastase release from human neutrophils stimulated by human IL-8; Figure 10 demonstrates the relative abilities of the chimeric anti-IL-8 Fabs to inhibit elastase release from human neutrophils stimulated by rabbit IL-8.

C. MOLECULAR CLONING OF THE VARIABLE LIGHT AND HEAVY REGIONS OF THE MURINE

20 5.12.14 (ANTI-IL-8) MONOCLONAL ANTIBODY

Total RNA was isolated from 1×10^6 cells (hybridoma cell line ATCC HB-11722) using the procedure described by Chomczynski and Sacchi (*Anal. Biochem.* 162:156 (1987)). First strand cDNA was synthesized by specifically priming the mRNA with synthetic DNA oligonucleotides designed to hybridize with regions of the murine RNA encoding the constant region of the kappa light chain or the IgG2a heavy chain (the DNA sequence of these regions are published in *Sequences of Proteins of Immunological Interest*, Kabat, E. A. *et al.* (1991) NIH Publication 91-3242, V 1-3.). Three primers (SEQ ID NOS: 1-6) were designed for each of the light and heavy chains to increase the chances of primer hybridization and efficiency of first strand cDNA synthesis (Figure 13). Amplification of the first strand cDNA to double-stranded (ds) DNA was accomplished using two sets of synthetic DNA oligonucleotide primers: one forward primer (SEQ ID NOS: 7-9) and one reverse primer (SEQ ID NO: 10) for the light chain variable region amplification (Figure 14) and one forward primer (SEQ ID NOS: 11-14) and one reverse primer (SEQ ID NOS: 11, 15, 14 and 13) for the heavy chain variable region amplification (Figure 15). The N-terminal sequence of the first eight amino acids of either the light or heavy chains of 5.12.14 was used to generate a putative murine DNA sequence corresponding to this region. (A total of 29 amino acids was sequenced from the N-terminus of both the light chain and heavy chain variable regions using the Edman degradation protein sequencing technique.) This information was used to design the forward amplification primers which were made degenerate in the third position for some codons to increase the chances of primer hybridization to the natural murine DNA codons and also included the unique restriction site, MluI, for both the light chain variable region forward primer and the heavy chain variable region forward primer to facilitate ligation to the 3' end of the STII element in the

cloning vector. The reverse amplification primers were designed to anneal with the murine DNA sequence corresponding to a portion of the constant region of the light or heavy chains near the variable/constant junction. The light chain variable region reverse primer contained a unique BstBI restriction site and the heavy chain variable region reverse primer contained a unique ApaI restriction site for ligation to the 5' end of either the human IgG1 constant light or IgG1 constant heavy regions in the vectors, pB13.1 (light chain) and pB14 (heavy chain). The polymerase chain reaction using these primer sets yielded DNA fragments of approximately 400 bp. The cDNA encoding the 5.12.14 light chain variable region was cloned into the vector pB13.1, to form pA51214VL and the 5.12.14 heavy chain variable region was cloned into the vector, pB14, to form pA51214VH. The cDNA inserts were characterized by DNA sequencing and are presented in the DNA sequence (SEQ ID NO: 16) and amino acid sequence (SEQ ID NO: 17) of Figure 16 (murine light chain variable region) and in the DNA sequence (SEQ ID NO: 18) and amino acid (SEQ ID NO: 19) of Figure 17 (murine heavy chain variable region).

D. CONSTRUCTION OF A 5.12.14 FAB VECTOR

In the initial construct, pA51214VL, the amino acids between the end of the 5.12.14 murine light chain variable sequence and the unique cloning site, BstBI, in the human IgG1 constant light sequence were of murine origin corresponding to the first 13 amino acids of the murine IgG1 constant region (Figure 16). Therefore, this plasmid contained a superfluous portion of the murine constant region separating the 5.12.14 murine light chain variable region and the human light chain IgG1 constant region. This intervening sequence would alter the amino acid sequence of the chimera and most likely produce an incorrectly folded Fab. This problem was addressed by immediately truncating the cDNA clone after A109 and re-positioning the BstBI site to the variable/constant junction by the polymerase chain reaction. Figure 18 shows the amplification primers used to make these modifications. The forward primer, VL.front (SEQ ID NO: 20), was designed to match the last five amino acids of the STII signal sequence, including the MluI cloning site, and the first 4 amino acids of the 5.12.14 murine light chain variable sequence. The sequence was altered from the original cDNA in the third position of the first two codons D1 (T to C) and I2 (C to T) to create a unique EcoRV cloning site which was used for later constructions. The reverse primer, VL.rear (SEQ ID NO: 21), was designed to match the first three amino acids of the human IgG1 constant light sequence and the last seven amino acids of the 5.12.14 light chain variable sequence which included a unique BstBI cloning site. In the process of adding the BstBI site, the nucleotide sequence encoding several amino acids were altered: L106 (TTG to CTT), K107 (AAA to CGA) resulting in a conservative amino acid substitution to arginine, and R108 (CGG to AGA). The PCR product encoding the modified 5.12.14 light chain variable sequence was then subcloned into pB13.1 in a two-part ligation. The MluI-BstBI digested 5.12.14 PCR product encoding the light chain variable region was ligated into MluI-BstBI digested vector to form the plasmid, pA51214VL'. The modified cDNA was characterized by DNA sequencing. The coding sequence for the 5.12.14 light chain is shown in Figure 19.

Likewise, the DNA sequence between the end of the heavy chain variable region and the unique cloning site, ApaI, in the human IgG1 heavy chain constant domain of pA51214VH was reconstructed to change the amino acids in this area from murine to human. This was done by the polymerase chain reaction. Amplification of the murine 5.12.14 heavy chain variable sequence was accomplished using the primers shown in Figure 18. The forward PCR primer (SEQ ID NO: 22) was designed to match nucleotides 867-887 in

pA51214VH upstream of the STII signal sequence and the putative cDNA sequence encoding the heavy chain variable region and included the unique cloning site SpeI. The reverse PCR primer (SEQ ID NO: 23) was designed to match the last four amino acids of the 5.12.14 heavy chain variable sequence and the first six amino acids corresponding to the human IgG1 heavy constant sequence which also included the unique cloning site, Apal. The PCR product encoding the modified 5.12.14 heavy chain variable sequence was then subcloned to the expression plasmid, pMHM24.2.28 in a two-part ligation. The vector was digested with SpeI-Apal and the SpeI-Apal digested 5.12.14 PCR product encoding the heavy chain variable region was ligated into it to form the plasmid, pA51214VH'. The modified cDNA was characterized by DNA sequencing. The coding sequence for the 5.12.14 heavy chain is shown in the DNA sequence (SEQ ID NO: 26) and amino acid sequence (SEQ ID NO: 27) of Figures 20A-20B.

The first expression plasmid, pantilL-8.1, encoding the chimeric Fab of 5.12.14 was made by digesting pA51214VH' with EcoRV and Bpu1102I to replace the EcoRV-Bpu1102I fragment with a EcoRV-Bpu1102I fragment encoding the murine 5.12.14 light chain variable region of pA51214VL'. The resultant plasmid thus contained the murine-human variable/constant regions of both the light and heavy chains of 5.12.14.

Preliminary analysis of Fab expression using pantilL-8.1 showed that the light and heavy chains were produced intracellularly but very little was being secreted into the periplasmic space of *E. coli*. To correct this problem, a second expression plasmid was constructed.

The second expression plasmid, pantilL-8.2, was constructed using the plasmid, pmy187, as the vector. Plasmid pantilL-8.2 was made by digesting pmy187 with MluI and SphI and the MluI (partial)-SphI fragment encoding the murine 5.12.14 murine-human chimeric Fab of pantilL-8.1 was ligated into it. The resultant plasmid thus contained the murine-human variable/constant regions of both the light and heavy chains of 5.12.14.

The plasmid pantilL-8.2 was deposited on February 10, 1995 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, U.S.A. (ATCC) and assigned ATCC Accession No. ATCC 97056.

E. MOLECULAR CLONING OF THE VARIABLE LIGHT AND HEAVY REGIONS OF THE MURINE 6G4.2.5 MONOCLONAL ANTIBODY

Total RNA was isolated from 1×10^8 cells (hybridoma cell line 6G4.2.5) using the procedure described by Chomczynski and Sacchi (*Anal. Biochem.* 162:156 (1987)). First strand cDNA was synthesized by specifically priming the mRNA with synthetic DNA oligonucleotides designed to hybridize with regions of the murine RNA encoding the constant region of the kappa light chain or the IgG2a heavy chain (the DNA sequence of these regions are published in *Sequences of Proteins of Immunological Interest*, Kabat *et al.* (1991) NIH Publication 91-3242, V 1-3). Three primers (SEQ ID NOS: 1-6) were designed for each the light and heavy chains to increase the chances of primer hybridization and efficiency of first strand cDNA synthesis (Figure 21). Amplification of the first strand cDNA to double-stranded (ds) DNA was accomplished using two sets of synthetic DNA oligonucleotide primers: one forward primer (SEQ ID NOS: 28-30) and one reverse primer (SEQ ID NO: 31) for the light chain variable region amplification (Figure 22) and one forward primer (SEQ ID NOS: 32-33) and one reverse primer (SEQ ID NOS: 11,15,14 and 13) for the

heavy chain variable region amplification (Figure 23). The N-terminal sequence of the first eight amino acids of either the light or heavy chains of 6G4.2.5 was used to generate a putative murine DNA sequence corresponding to this region. (A total of 29 amino acids were sequenced from the N-terminus of both the light chain and heavy chain variable regions using the Edman degradation protein sequencing technique.) This information was used to design the forward amplification primers which were made degenerate in the third position for some codons to increase the chances of primer hybridization to the natural murine DNA codons and also included the unique restriction site, NsiI, for the light chain variable region forward primer and the unique restriction site, MluI, for the heavy chain variable region forward primer to facilitate ligation to the 3' end of the STII element in the vector, pchimFab. The reverse amplification primers were designed to anneal with the murine DNA sequence corresponding to a portion of the constant region of the light or heavy chains near the variable/constant junction. The light chain variable region reverse primer contained a unique MunI restriction site and the heavy chain variable region reverse primer contained a unique ApaI restriction site for ligation to the 5' end of either the human IgG1 constant light or IgG1 constant heavy regions in the vector, pchimFab. The polymerase chain reaction using these primer sets yielded DNA fragments of approximately 400 bp and were cloned individually into the vector, pchimFab, to form p6G425VL and p6G425VH. The cDNA inserts were characterized by DNA sequencing and are presented in the DNA sequence (SEQ ID NO: 34) and amino acid sequence (SEQ ID NO: 35) of Figure 24 (murine light chain variable region) and the DNA sequence (SEQ ID NO: 36) and amino acid sequence (SEQ ID NO: 37) of Figure 25 (murine heavy chain variable region).

20 F. CONSTRUCTION OF A 6G4.2.5 CHIMERIC FAB VECTOR

~~In the initial construct, p6G425VL, the amino acids between the end of the 6G4.2.5 murine light chain variable sequence and the unique cloning site, MunI, in the human IgG1 constant light sequence were of murine origin. These amino acids must match the human IgG1 amino acid sequence to allow proper folding of the chimeric Fab. Two murine amino acids, D115 and S121, differed dramatically from the amino acids found in the loops of the β -strands of the human IgG1 constant domain and were converted to the proper human amino acid residues, V115 and F121, by site-directed mutagenesis using the primers (SEQ ID NOS: 38,39,40) shown in Figure 26. These specific mutations were confirmed by DNA sequencing and the modified plasmid named p6G425VL'. The coding sequence is shown in the DNA sequence (SEQ ID NO: 41) and amino acid sequence (SEQ ID NO: 42) of Figures 27A-27B.~~

30 Likewise, the DNA sequence between the end of the heavy chain variable region and the unique cloning site, ApaI, in the human IgG1 heavy chain constant domain of p6G425VH was reconstructed to change the amino acids in this area from murine to human. This process was facilitated by the discovery of a BstEII site near the end of the heavy chain variable region. This site and the ApaI site were used for the addition of a synthetic piece of DNA encoding the corresponding IgG human amino acid sequence. The synthetic oligo-nucleotides shown in Figure 26 were designed as complements of one another to allow the formation of a 27 bp piece of ds DNA. The construction was performed as a three-part ligation because the plasmid, p6G425VH, contained an additional BstEII site within the vector sequence. A 5309 bp fragment of p6G425VH digested with MluI-ApaI was ligated to a 388 bp fragment carrying the 6G4.2.5 heavy chain variable region and a 27 bp synthetic DNA fragment encoding the first six amino acids of the human IgG1

constant region to form the plasmid, p6G425VH'. The insertion of the synthetic piece of DNA was confirmed by DNA sequencing. The coding sequence is shown in the DNA sequence (SEQ ID NO: 43) and amino acid sequence (SEQ ID NO: 44) of Figures 28A-28B.

The expression plasmid, p6G425chim2, encoding the chimeric Fab of 6G4.2.5 was made by digesting p6G425chimVL' with MluI and ApaI to remove the-STII-murine HPC4 heavy chain variable region and replacing it with the MluI-ApaI fragment encoding the STII-murine 6G4.2.5 heavy chain variable region of p6G425chimVH'. The resultant plasmid thus contained the murine-human variable/constant regions of both the light and heavy chains of 6G4.2.5.

The plasmid p6G425chim2 was deposited on February 10, 1995 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, U.S.A. (ATCC) and assigned ATCC Accession No. 97055.

G. CONSTRUCTION OF HUMANIZED VERSIONS OF ANTI-IL-8 ANTIBODY 6G4.2.5

The murine cDNA sequence information obtained from the hybridoma cell line, 6G4.2.5, was used to construct recombinant humanized variants of the murine anti-IL-8 antibody. The first humanized variant, F(ab)-1, was made by grafting synthetic DNA oligonucleotide primers encoding the murine CDRs of the heavy and light chains onto a phagemid vector, pEMXI (Werther *et al.*, *J. Immunol.*, **157**: 4986-4995 (1996)), which contains a human 6-subgroup I light chain and a human IgG1 subgroup III heavy chain (Fig. 29). Amino acids comprising the framework of the antibody that were potentially important for maintaining the conformations necessary for high affinity binding to IL-8 by the complementarity-determining regions (CDR) were identified by comparing molecular models of the murine and humanized 6G4.2.5 (F(ab)-1) variable domains using methods described by Carter *et al.*, PNAS 89:4285 (1992) and Eigenbrot, *et al.*, J. Mol. Biol. 229:969 (1993). Additional humanized framework variants (F(ab) 2-9) were constructed from the information obtained from these models and are presented in Table 2 below. In these variants, the site-directed mutagenesis methods of Kunkel, *Proc. Natl. Acad. Sci USA*, **82**:488 (1985) were utilized to exchange specific human framework residues with their corresponding 6G4.2.5 murine counterparts. Subsequently, the entire coding sequence of each variant was confirmed by DNA sequencing. Expression and purification of each F(ab) variant was performed as previously described by Werther *et al.*, *supra*, with the exception that hen egg white lysozyme was omitted from the purification protocol. The variant antibodies were analyzed by SDS-PAGE, electrospray mass spectroscopy and amino acid analysis.

Table 4 - Humanized 6G425 Variants

Variant	Version	Template	Changes ^a	Purpose ^b	IC50 ^c		
					Mean	S.D.	N
F(ab)-1	version 1		CDR Swap		63.0	12.3	4
F(ab)-2	version 2	F(ab)-1	PheH67Ala	packaging w/ CDR H2	106.0	17.0	2

F(ab)-3	version 3	F(ab)-1	ArgH71 <i>Val</i>	packaging w/ CDRs H1, H2	79.8	42.2	4
F(ab)-4	version 6	F(ab)-1	IleH69 <i>Leu</i>	packaging w/ CDR H2	44.7	9.0	3
F(ab)-5	version 7	F(ab)-1	LeuH78 <i>Ala</i>	packaging w/ CDRs H1, H2	52.7	31.0	9
F(ab)-6	version 8	F(ab)-1	IleH69 <i>Leu</i> LeuH78 <i>Ala</i>	combine F(ab)-4 and -5	34.6	6.7	7
F(ab)-7	version 16	F(ab)-6	LeuH80 <i>Val</i>	packaging w/ CDR H1	38.4	9.1	2
F(ab)-8	version 19	F(ab)-6	ArgH38 <i>Lys</i>	packaging w/ CDR H2	14.0	5.7	2
F(ab)-9	version 11	F(ab)-6	GluH6 <i>Gln</i>	packaging w/ CDR H3	19.0	5.1	7
Chimeric ^d F(ab)					11.4	7.0	13
rhu4D5 ^e F(ab)					>200μM		5

- a Amino acid changes made relative to the template used. Murine residues are in bold italics and residue numbering is according to Kabat *et al.*
- b Purpose for making changes based upon interactions observed in molecular models of the humanized and murine variable domains.
- 5 c. nM concentration of variant necessary to inhibit binding of iodinated IL-8 to human neutrophils in the competitive binding assay.
- d Chimeric F(ab) is a F(ab) which carries the murine heavy and light chain variable domains fused to the human light chain κI constant domain and the human heavy chain subgroup III constant domain I respectively.
- 10 e. rhu4D5F(ab) is of the same isotype as the humanized 6G425 F(ab)s and is a humanized anti-HER2 F(ab) and therefore should not bind to IL8.

The first humanized variant, F(ab)-1, was an unaltered CDR swap in which all the murine CDR amino acids defined by both x-ray crystallography and sequence hypervariability were transferred to the human framework. When the purified F(ab) was tested for its ability to inhibit ¹²⁵I-IL-8 binding to human neutrophils according to the methods described in Section (B)(1) above, a 5.5 fold reduction in binding affinity was evident as shown in Table 4 above. Subsequent versions of F(ab)-1 were engineered to fashion the 3-dimensional structure of the CDR loops into a more favorable conformation for binding IL-8. The relative affinities of the F(ab) variants determined from competition binding experiments using human neutrophils as

described in Section (B)(1) above are presented in Table 4 above. A slight decrease in IL-8 binding (<2 fold) was observed for F(ab)-2-3 while only slight increases in IL-8 binding were noted for F(ab)3-5. Variant F(ab)-6 had the highest increase in affinity for IL-8 (approximately 2 fold), exhibiting an IL-8 binding affinity of 34.6nM compared to the F(ab)-1 IL-8 binding affinity of 63nM. The substitutions of murine Leu for Ile at H69 and murine Ala for Leu at H78 are predicted to influence the packing of CDRs H1 and H2. Further framework substitutions using the F(ab)-6 variant as template were made to bring the binding affinity closer to that of the chimeric F(ab). *In-vitro* binding experiments revealed no change in affinity for F(ab)-7 (38.4nM) but a significant improvement in affinity for F(ab)-8/9 of 14nM and 19 nM, respectively. By analysis of a 3-D computer-generated model of the anti-IL-8 antibody, it was hypothesized that the substitution of murine Lys for Arg at H38 in F(ab)-8 influences CDR-H2 while a change at H6 of murine Gln for Glu in F(ab)-9 affects CDR-H3. Examination of the human antibody sequences with respect to amino acid variability revealed that the frequency of Arg at residue H38 is >99% whereas residue H6 is either Gln ~20% or Glu ~80% (Kabat *et. al.*, Sequences of Proteins of Immunological Interest 5th Ed. (1991)). Therefore, to reduce the likelihood of causing an immune response to the antibody, F(ab)-9 was chosen over F(ab)-8 for further affinity maturation studies. Variant F(ab)-9 was also tested for its ability to inhibit IL-8-mediated chemotaxis (Fig. 30). This antibody was able to block neutrophil migration induced by wild-type human IL-8, human monomeric IL-8 and Rhesus IL-8 with IC₅₀s of approximately 12nM, 15nM, and 22nM, respectively, in IL-8 mediated neutrophil chemotaxis inhibition assays performed as described in Section (B)(2) above. The amino acid sequence for variant F(ab)-8 is provided in Fig. 31c. The F(ab)-8 was found to block human and rhesus IL-8-mediated chemotaxis with IC₅₀s of 12nM and 10nM, respectively, in IL-8 mediated neutrophil chemotaxis inhibition assays performed as described in Section (B)(2) above.

H. CONSTRUCTION OF AN ANTI-IL-8-GENE III FUSION PROTEIN FOR PHAGE DISPLAY AND ALANINE SCANNING MUTAGENESIS

An expression plasmid, pPh6G4.V11, encoding a fusion protein (heavy chain of the humanized 6G4.2.5 version 11 antibody and the M13 phage gene-III coat protein) and the light chain of the humanized 6G4.2.5 version 11 antibody was assembled to produce a monovalent display of the anti-IL-8 antibody on phage particles. The construct was made by digesting the plasmid, pFPHX, with EcoRV and ApaI to remove the existing irrelevant antibody coding sequence and replacing it with a 1305bp EcoRV-ApaI fragment from the plasmid, p6G4.V11, encoding the humanized 6G4.2.5 version 11 anti-IL-8 antibody. The translated sequence of the humanized 6G4.2.5 version 11 heavy chain (SEQ ID NO: 52), peptide linker and gene III coat protein (SEQ ID NO: 53) is shown in Fig. 31A. The pFPHX plasmid is a derivative of pHGHam-3 which contains an in-frame amber codon (TAG) between the human growth hormone and gene-III DNA coding sequences. When transformed into an amber suppressor strain of *E. coli*, the codon (TAG) is read as Glutamate producing a growth hormone (hGH)-gene III fusion protein. Likewise, in a normal strain of *E. coli*, the codon (TAG) is read as a stop preventing translational read-through into the gene-III sequence and thus allowing the production of soluble hGH. The pHGHam-3 plasmid is described in Methods: A Companion to Methods in Enzymology, 3:205 (1991). The final product, pPh6G4.V11, was used as the template for the alanine scanning mutagenesis of the CDRs and for the construction of randomized CDR libraries of the humanized 6G4.V11

antibody.

I. ALANINE SCANNING MUTAGENESIS OF HUMANIZED ANTIBODY 6G4.2.5 VERSION 11

The solvent exposed amino acid residues in the CDRs of the humanized anti-IL-8 6G4.2.5 version 11 antibody (h6G4V11) were identified by analysis of a 3-D computer-generated model of the anti-IL-8 antibody.

- 5 In order to determine which solvent exposed amino acids in the CDRs affect binding to interleukin-8, each of the solvent exposed amino acids was individually changed to alanine, creating a panel of mutant antibodies wherein each mutant contained an alanine substitution at a single solvent exposed residue. The alanine scanning mutagenesis was performed as described by Leong *et. al.*, J. Biol. Chem., 269: 19343 (1994)).

- The IC₅₀'s (relative affinities) of h6G4V11 wt and mutated antibodies were established using a
- 10 Competition Phage ELISA Assay described by Cunningham *et. al.*, (EMBO J. 13:2508 (1994)) and Lee *et. al.*, (Science 270:1657 (1995)). The assay measures the ability of each antibody to bind IL-8 coated onto a 96-well plate in the presence of various concentrations of free IL-8 (0.2 to 1uM) in solution. The first step of the assay requires that the concentrations of the phage carrying the wild type and mutated antibodies be normalized, allowing a comparison of the relative affinities of each antibody. The normalization was accomplished by
- 15 titrating the phage on the IL-8 coated plates and establishing their EC₅₀. Sulfhydryl coated 96-well binding plates (Corning-Costar, Wilmington, MA) were incubated with a 0.1mg/ml solution of K64C IL-8 (Lysine 64 is substituted with Cysteine to allow the formation of a disulfide bond between the free thiol group of K64C IL-8 and the sulfhydryl coated plate, which results in the positioning of the IL-8 receptor binding domains towards the solution interface) in phosphate buffered saline (PBS) pH 6.5 containing 1mM EDTA for 1 hour at
- 20 25°C followed by three washes with PBS and a final incubation with a solution of PBS containing 1.75mg/ml of L-cysteine-HCl and 0.1M NaHCO₃ to block any free reactive sulfhydryl groups on the plate. The plates were washed once more and stored covered at 4°C with 200ul of PBS/well. Phage displaying either the reference antibody, h6G4V11, or the mutant h6G4V11 antibodies were grown and harvested by PEG precipitation. The phage were resuspended in 500ul 10mM Tris-HCl pH 7.5, 1mM EDTA and 100mM NaCl
- 25 and held at 4°C for no longer than 3 hours. An aliquot of each phage was diluted 4-fold in PBS containing 0.05% Tween-20 (BioRad, Richmond, Ca.) and 0.5% BSA RJA grade (Sigma, St. Louis, Mo.) (PBB) and added to IL-8 coated plates blocked for at least 2 hours at 25°C with 50mg/ml skim milk powder in 25mM Carbonate Buffer pH 9.6. The phage were next serially diluted in 3 fold steps down the plate from well A through H. The plates were incubated for 1 hour at 25°C followed by nine quick washes with PBS containing
- 30 0.05% Tween-20 (PBST). The plates were then incubated with a 1:3200 dilution of rabbit anti-phage antibody and a 1:1600 dilution of secondary goat-anti-rabbit Fc HRP-conjugated antibody for 15 minutes at 25°C followed by nine quick washes with PBST. The plates were developed with 80ul/well of 1mg/ml OPD (Sigma, St. Louis, Mo) in Citrate Phosphate buffer pH 5.0 containing 0.015% H₂O₂ for 4 minutes at 25°C and the reaction stopped with the addition of 40ul of 4.5M H₂SO₄. The plates were analyzed at wavelength 8492 in a
- 35 SLT model 340ATTC plate reader (SLT Lab Instruments). The individual EC₅₀'s were determined by analyzing the data using the program Kaleidagraph (Synergy Software, Reading, Pa.) and a 4-parameter fit

equation. The phage held at 4°C were then immediately diluted in PBB to achieve a final concentration corresponding to their respective EC₅₀ or target OD₄₉₂ for the competition segment of the experiment, and dispensed into a 96 well plate containing 4-fold serial dilutions of soluble IL-8 ranging from 1uM in well A and ending with 0.2uM in well H. Using a 12-channel pipet, 100ul of the phage/IL-8 mixture was transferred to an IL-8 coated 96-well-plate and executed as described above. Each sample was done in triplicate - 3 columns/sample.

Table 5 - Relative Affinities (IC₅₀) for Alanine-scan Anti-IL-8 6G4V11 CDR Mutants

CDR	Amino Acid Residue	Avg IC ₅₀ (nM)	Std Dev
V11	Reference	11.5	6.4
CDR-L1	S26	6.3	2.9
	Q27	10.2	2.4
	S28	14.2	5.2
	V30	29.1	12.3
	H31	580.3	243.0
	I33	64.2	14.6
	N35	3.3	0.7
	T36	138.0	nd
	Y37	NDB	nd
CDR-L2	K55	24.2	14.9
	V56	15.5	3.8
	S57	12.4	4.0
	N58	17.6	3.7
	R59	nd	nd
CDR-L3	S96	10.8	4.4
	T97	70.6	55.2
	H98	8.0	1.2
	V99	19.6	1.9

CDR	Amino Acid Residue	Avg IC50 (nM)	Std Dev
CDR-H1	S28	8.6	3.1
	S30	nd	nd
	S31	7.8	2.5
	H32	13.3	5.8
	Y53	48.2	15.8
CDR-H2	Y50	35.6	13.0
	D52	13.3	7.5
	S53	6.0	3.4
	N54	96.0	5.8
	E56	15.8	4.5
	T57	8.4	1.6
	T58	11.3	1.8
	Y59	9.1	3.7
	Q61	12.6	6.4
	K64	18.5	12.1
CDR-H3	D96	NDB	nd
	Y97	NDB	nd
	R98	36.6	15.3
	Y99	199.5	nd
	N100	278.3	169.4
	D102	159.2	44
	W103	NDB	nd
	F104	NDB	nd
	F105	209.4	72.3

CDR	Amino Acid Residue	Avg IC ₅₀ (nM)	Std Dev
	D106	25.3	21.7

Each sample performed in triplicate/experiment.

NDB = No Detectable Binding /nd = value not determined*

Residue numbering is according to Kabat et al.

The results of the alanine-scan are summarized in Table 5 above. The alanine substitutions in of many of the mutant antibodies had little or no adverse effects (<3 fold) on the binding affinity for IL-8. Mutants that were found to exhibit no detectable binding of IL-8 (NDB) presumably contained disruptions in the conformational structure of the antibody conferred by crucial structural or buried amino acids in the CDR. Based on the results of the scan, CDR-H3 (heavy chain, 3rd CDR) was identified as the dominant binding epitope for binding IL-8. Alanine substitutions in this CDR resulted in a 3 to >26 fold decrease in binding affinity. The amino acids, Y597, Y599 and D602 are of particular interest because it was determined from the computer generated model of the anti-IL-8 antibody that these residues are solvent exposed and that these residues might participate in hydrogen bonding or charge interactions with IL-8 or other amino acids of the antibody that influence either binding to IL-8 or the conformation of the CDR-H3 loop structure. (See the model depicted in Fig. 32). Unexpected increases in binding affinity (1.8 > 2.7 fold) were noted for S528 and S531 of CDR-H1 and S553 of CDR-H2.

Surprisingly, a significant increase in binding affinity was observed in the alanine mutant N35A located in CDR-L1 (light chain, 1st CDR). A 3-6 fold increase in affinity was observed compared to the wild-type h6G4V11 antibody. This augmentation of IL-8 binding could be the result of the close proximity of N35A to CDR-H3. The alanine substitution may have imparted a slight change in the conformation of CDR-L1 which alters the packing interaction of neighboring amino acid residues on CDR-H3, thereby tweaking the loop of CDR-H3 into a conformation that facilitates more appropriate contacts with IL-8. Similarly, N35A may also influence the orientation of amino acids in CDR-L1 or its interaction directly with IL-8. Unexpected increases in affinity (~2 fold) were also observed for S26 of CDR-L1 and H98 of CDR-L3.

J. CHARACTERIZATION OF HUMANIZED ANTI-IL-8 ANTIBODY 6G4V11N35A

Soluble 6G4V11N35A Fab antibody was made by transforming an amber non-suppressor strain of *E. coli*, 34B8, with pPh6G4.V11 and growing the culture in low phosphate medium for 24 hours. The periplasmic fraction was collected and passed over a Hi-Trap Protein-G column (Pharmacia, Piscataway, NJ.) followed by a desalting and concentration step. The protein was analyzed by SDS-PAGE, mass spectrometry and amino acid analysis. The protein had the correct size and amino acid composition (Fig. 35). The 6G4V11N35A Fab was tested for its ability to inhibit ¹²⁵I-IL-8 binding to human neutrophils and to inhibit IL-8 mediated neutrophil chemotaxis as described in Section (B)(1) and (B)(2) above. As shown in Fig. 33, hybridoma-derived intact murine antibody (6G4 murine mAb), recombinant 6G4 murine-human chimera Fab, recombinant humanized Fab versions 1 and 11, and 6G4V11N35A Fab were found to inhibit ¹²⁵I-IL-8 binding to human neutrophils with an average IC₅₀ of 5nM, 8nM, 40nM, 10nM and 3nM, respectively. The 6G4V11N35A Fab had at least a 2-fold higher affinity than the 6G4.2.5 chimera Fab and a 3-fold higher

affinity than 6G4V11. As shown in Fig. 34, the 6G4V11N35A Fab was found to inhibit IL-8 mediated neutrophil chemotaxis induced by both wild type and monomeric human IL-8, and by two different animal species of IL-8, namely, rabbit and rhesus. The irrelevant isotype control Fab (4D5) did not inhibit neutrophil migration. The average IC_{50} values were 3nM (wt IL-8), 1 nM (monomeric IL-8), 5nM (Rabbit IL-8), and 10nM (Rhesus IL-8).

K. CONSTRUCTION OF A 6G4V11N35A F(ab')₂ LEUCINE ZIPPER

Production of a F(ab')₂ version of the humanized anti-IL-8 6G4V11N35A Fab was accomplished by constructing a fusion protein with the yeast GCN4 leucine zipper. The expression plasmid p6G4V11N35A.F(ab')₂ was made by digesting the plasmid p6G425chim2.fab2 with the restriction enzymes *bsal* and *apaI* to remove the DNA sequence encoding the 6G4.2.5 murine-human chimeric Fab and replacing it with a 2620bp *bsal*-*apaI* fragment from pPh6G4.V11N35A. The plasmid p6G425chim2.fab2 is a derivative of pS1130 which encodes a fusion protein (the GCN4 leucine zipper fused to the heavy chain of anti-CD18) and the light chain of anti-CD18 antibody. The expression plasmid p6G4V11N35A.F(ab')₂ was deposited on February 20, 1996 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, U.S.A. (ATCC) and assigned ATCC Accession No. 97890. A pepsin cleavage site in the hinge region of the antibody facilitates the removal of the leucine zipper leaving the two immunoglobulin monomers joined by the cysteines that generate the interchain disulfide bonds. The DNA and protein sequence of the h6G4V11N35A.F(ab')₂ are depicted in Figs. 35-37.

~~An expression host cell was obtained by transforming E. coli strain 49D6 with~~ p6G4V11N35A.F(ab')₂ essentially as described in Section (II)(3)(C) above. The transformed host E. coli 49D6 (p6G4V11N35A.F(ab')₂) was deposited on February 20, 1997 at the ATCC and assigned ATCC Accession No. 98332. Transformed host cells were grown in culture, and the 6G4V11N35A F(ab')₂ product was harvested from the host cell periplasmic space essentially as described in Section (II)(3)(F) above.

L. CHARACTERIZATION OF THE HUMANIZED 6G4V11N35A F(ab')₂ LEUCINE ZIPPER

The 6G4V11N35A Fab and F(ab')₂ were tested for their ability to inhibit ¹²⁵I-IL-8 binding to neutrophils according to the procedures described in Section (B)(1) above. The displacement curves from a representative binding experiment performed in duplicate is depicted in Fig. 38. Scatchard analysis of this data shows that 6G4V11N35A F(ab')₂ inhibited ¹²⁵I-IL-8 binding to human neutrophils with an average IC_{50} of 0.7 nM (+/- 0.2). This is at least a 7 fold increase in affinity compared to the hybridoma-derived intact murine antibody (average IC_{50} of 5 nM) and at least a 2.8 fold increase in affinity over the Fab version (average IC_{50} of 2 nM).

The 6G4V11N35A F(ab')₂ was also tested for its ability to inhibit IL-8 mediated neutrophil chemotaxis according to the procedures described in Section (B)(2) above. The results of a representative

chem taxis experiment performed in quadruplicate are depicted in Fig. 39. As shown in Fig. 39, the 6G4V11N35A F(ab')₂ inhibited human IL-8 mediated neutrophil chemotaxis. The 6G4V11N35A F(ab')₂ exhibited an average IC₅₀ value of 1.5nM versus 2.7nM for the 6G4V11N35A Fab, which represents an approximately 2 fold improvement in the antibody's ability to neutralize the effects of IL-8. The irrelevant isotype control Fab (4D5) did not inhibit neutrophil migration. Furthermore, the 6G4V11N35A F(ab')₂ antibody retained its ability to inhibit IL-8 mediated neutrophil chemotaxis by monomeric IL-8 and by two different animal species of IL-8, namely rabbit and rhesus, in neutrophil chemotaxis experiments conducted as described above. An individual experiment is shown in Fig. 40. The average IC₅₀ values were 1nM (monomeric IL-8), 4nM (Rabbit IL-8), and 2.0nM (Rhesus IL-8).

M. RANDOM MUTAGENESIS OF LIGHT CHAIN AMINO ACID (N35A) IN CDR-L1 OF HUMANIZED ANTIBODY 6G4V11

A 3-fold improvement in the IC₅₀ for inhibiting ¹²⁵I-IL-8 binding to human neutrophils was observed when alanine was substituted for asparagine at position 35 in CDR-L1 (light chain) of the humanized 6G4V11 mAb as described in Section (I) above. This result might be attributed to an improvement in the contact between the antigen-antibody binding interfaces as a consequence of the replacement of a less bulky nonpolar side chain (R-group) that may have altered the conformation of CDR-L1 or neighboring CDR-H3 (heavy chain) to become more accessible for antigen docking. The acceptance of alanine at position 35 of CDR-L1 suggested that this position contributed to improved affinity and that an assessment of the re-modeling of CDR loops / antigen-binding region(s) by other amino acids at this location was warranted. Selection of an affinity matured version of the humanized 6G4.V11 mAb (Kunkel, T. A., Proc. Natl. Acad. Sci. USA, 82:488 (1995)) was accomplished by randomly mutagenizing position 35 of CDR-L1 and constructing an antibody-phage library. The codon for Asparagine (N) at position 35 of CDR-L1, was targeted for randomization to any of the 20 known amino acids.

Initially, a stop template, pPh6G4.V11-stop, was made to eliminate contaminating wild-type N35 sequence from the library. This was accomplished by performing site-directed mutagenesis (Muta-Gene Kit, Biorad, Richmond, CA) of pPh6G4V11 (described in Section (H) above) to replace the codon (AAC) for N35 with a stop codon (TAA) using the primer SL.97.2 (SEQ ID NO:63) (Figure 42). The incorporation of the stop codon was confirmed by DNA sequencing. Subsequently, uracil containing single-stranded DNA derived from *E. coli* CJ236 transformed with the stop template was used to generate an antibody-phage library following the method described by Lowman (Methods in Molecular Biology, 87 Chapter 25: 1-15 (1997)). The variants generated from this library were predicted to produce a collection of antibodies containing one of the 20 known amino acids at position N35 in CDR-L1. The amino acid substitutions were accomplished by site-directed mutagenesis using the degenerate oligonucleotide primer (SL.97.3) with the sequence NNS (N = A/G/T/C; S = G/C;) (SEQ ID NO: 64) (Figure 42). This codon usage should allow for the expression of any of the 20 amino acids - including the amber stop codon (TAG). The collection of antibody-phage variants was transfected into *E. coli* strain XL-1 blue (Stratagene, San Diego, CA) by electroporation and grown at 37°C overnight to amplify the library. Selection of tight binding humanized 6G4V11 Fab's were accomplished by

panning the library on IL-8 coated 96-well plates as described in Section (I) above. Prior to panning, the number of phage/library was normalized to 1.1×10^{13} phage/ml (which produces a maximum OD₂₇₀ reading = 1 OD unit) and IL-8 coated plates were incubated with blocking solution (25mN Carbonate buffer containing 50mg/ml skim milk) for 2 hours before the addition of phage (each sort used eight IL-8 coated wells/library).

5 After the blocking and washing steps, every sort began with the addition of 100ul of antibody-phage (titered at 1.1×10^{13} phage/ml) to each of eight IL-8 coated wells followed by an 1 hour incubation at 25°C. The non-specifically bound antibody-phage were removed by 10 quick washes with PBS-0.05% Tween 20 (PBS-Tween). For sort #1, a low stringency wash (100ul PBS-Tween/well for 10 minutes at 25°C) was employed to capture the small proportion of tight binding antibody-phage bound to the immobilized IL-8. The antibody-phage variants specifically bound to IL-8 were eluted with 100ul/well of 200mM Glycine pH 2.0 for 5 minutes

10 at 25°C. The eluted antibody-phage variants from the 8 wells were then pooled and neutralized with 1M Tris-HCl pH 8.0 (1/3 the elution volume). The phage were titered and propagated as described in Section (I) above. The stringency of the washes were successively increased with each round of panning depending upon the percent recovery of phage at the end of a sort. The wash conditions were as follows: sort #2 (4 x 15 minute

15 intervals; total time = 60 minutes) and sort #3 (either #3a: 8 x 15 minute intervals or #3b: 12 x 10 minute intervals; total time = 120 minutes). The total number of phage recovered was progressively reduced after each sort suggesting that non- or weak- binders were being selected against. The recovery of the negative control (the antibody-phage stop variant) was constant throughout the panning (approximately 0.0001 to 0.00001 percent).

20 Eighteen random variants from sort #3 were analyzed by DNA sequencing to look for an amino acid consensus at position 35 of CDR-L1. The data presented in Figure 43A showed that Glycine occupied position 35 in 33% of the variants sequenced. However, after correcting for the number of NNS codon combinations/amino acid, the frequency of Glycine was reduced to 16.6%. Glutamic Acid was represented with the highest frequency (22%) followed by Aspartic Acid and Glycine (16.6%). The frequencies of recovery of

25 the wild-type Asparagine and substituted Alanine were only 5.6%. Interestingly, the high frequency of Glycine may suggest that a much wider range of conformations might be allowed for the loop of CDR-L1 which may be attributed to the reduction in steric hindrance of bond angle (ϕ - ψ) pairing as a result of the single hydrogen atom as the side chain. Conversely, Glutamic Acid at position 35 might restrict the flexibility of the loop by imposing less freedom of rotation imposed by the more rigid and bulky charged polar side chain.

30 Soluble Fab's of the affinity matured variants (N35G, N35D, N35E and N35A) were made as described in Section (J) above for evaluating their ability to block IL-8 binding. As shown in Figure 43B, variants N35A, N35D, N35E and N35G were found to inhibit ¹²⁵I-IL-8 binding to human neutrophils with an approximate IC₅₀ of 0.2nM, 0.9nM, 0.1nM and 3.0nM, respectively. All of the affinity matured variants showed an improvement in binding IL-8 ranging from 3 - 100 fold compared to the humanized 6G4V11 mAb.

35 The affinity-matured variant, 6G4V11N35E, was 2-fold more potent in blocking IL-8 binding to human neutrophils than the alanine-scan variant, 6G4V11N35A. Equilibrium and kinetic measurements of variants 6G4V11N35A and 6G4V11N35E were determined using KinEXA™ automated immunoassay system

(Sapidyne Instruments Inc., Idaho City, ID) as described by Blake et al., *J. Bi. I. Chem.* 271: 27677 (1996). The procedure for preparing the antigen-coated particles was modified as follows: 1 ml of activated agarose beads (Reacti-Gel 6X; Pierce, Rockford, IL) were coated with antigen in 50mM Carbonate buffer pH 9.6 containing 20ug/ml of human IL-8 and incubated with gentle agitation on a rocker overnight at 25°C. The IL-8 coated beads were then washed twice with 1M Tris-HCl pH 7.5 to inactivate any unreactive groups on the beads and blocked with Superblock (Pierce, Rockford, IL) for 1 hour at 25°C to reduce non-specific binding. The beads were resuspended in assay buffer (0.1% bovine serum albumin in PBS) to a final volume of 30 ml. A 550ul aliquot of the IL-8 coated bead suspension was used each time to pack a fresh 4mm high column in the KinEXA observation cell. The amount of unbound antibody from the antibody-antigen mixtures captured by the IL-8-coated beads in both the equilibrium and kinetic experiments was quantified using a fluorescently labeled secondary antibody. Murine 6G4.2.5 was detected with a R-PE AffiniPure F(ab')₂ goat anti-mouse IgG, Fc fragment specific 2° antibody (Jackson Immuno Research Laboratories, West Grove, PA) and humanized affinity matured N35A (Fab and F(ab')₂) and N35E Fab were detected with a R-PE AffiniPure F(ab')₂ donkey anti-human IgG (H+L) 2° antibody (Jackson Immunoresearch Laboratories, West Grove, PA); both at a 1:1000 dilution.

Equilibrium measurements were determined by incubating a constant amount of anti-IL-8 antibody (0.005ug/ml) with various concentrations of human IL-8 (0.009, 0.019, 0.039, 0.078, 0.156, 0.312, 0.525, 1.25, 2.5nM). The antibody-antigen mixture was incubated for 2 hours at 25°C to allow the molecules to reach equilibrium. Subsequently, each sample was passed over a naive IL-8 coated bead pack in the KinEXA observation cell at a flow rate of 0.5ml/minute for a total of 9 minutes/sample. The equilibrium constant (Kd) was calculated using the software provided by Sapidyne Instruments Inc.

Rates of association (ka) and dissociation (kd) were determined by incubating together a constant amount of antibody and antigen, and measuring the amount of uncomplexed anti-IL-8 bound to the IL-8 coated beads over time. The concentration of antibody used in the kinetic experiments was identical to that used in the equilibrium experiment described above. Generally, the amount of human IL-8 used was the concentration derived from the binding curves of the equilibrium experiment that resulted in 70% inhibition of anti-IL-8 binding to the IL-8 coated beads. Measurements were made every 15 minutes to collect approximately nine data points. The ka was calculated using the software provided by Sapidyne Instruments, Inc. The off rate was determined using the equation: $kd = Kd/ka$.

Figure 44 shows the equilibrium constants (Kd) for the affinitymatured variants 6G4V11N35E and 6G4V11N35A Fab's were approximately 54pM and 114pM, respectively. The improvement in affinity of 6G4V11N35E Fab for IL-8 can be attributed to a 2-fold faster rate of association (K_{on}) of 4.7×10^6 for 6G4V11N35E Fab versus 2.0×10^6 for 6G4V11N35A F(ab')₂. (The Kd of the 6G4V11N35A F(ab')₂ and 6G4V11N35A Fab are similar.) The dissociation rates (K_{off}) were not significantly different. Molecular modeling suggests that substitution of Asparagine with Glutamic Acid might either affect the antibody's interaction with IL-8 directly or indirectly by neutralizing the charge of neighboring residues R98 (CDR-H3)

or K50 (CDR-L2) in the CDR's to facilitate contact with IL-8. Another effect might be the formation of a more stable loop conformation for CDR-L1 that could have facilitated more appropriate contacts of other CDR-L1 loop residues with IL-8. The DNA (SEQ ID NO: 65) and amino acid (SEQ ID NO:62) sequences of p6G4V11N35E.Fab showing the Asparagine to Glutamic Acid substitution in the light chain are presented in Figure 45.

N. CHARACTERIZATION OF HUMANIZED ANTI-IL-8 VARIANT 6G4V11N35E Fab

The affinity matured Fab variant, 6G4V11N35E, was tested for its ability to inhibit IL-8 mediated neutrophil chemotaxis as described in Section (B)(2) above. The reusable 96-well chemotaxis chamber described in Section (B)(2) was replaced with endotoxin-free disposable chemotaxis chambers containing 5-micron PVP-free polycarbonate filters (ChemoTx101-5, Neuro Probe, Inc. Cabin John, MD). As illustrated in Figure 46, variant N35E effectively blocks IL-8 mediated neutrophil chemotaxis induced by a 2nM stimulus of either rabbit or human IL-8. In fact, the level of inhibition at antibody concentrations between 3.7nM - 33nM was not significantly different from the buffer control indicating variant N35E could completely inhibit this response. The IC_{50} 's for both rabbit and human IL-8 were approximately 2.8nM and 1.2nM, respectively.

The irrelevant isotype control Fab (4D5) did not inhibit neutrophil migration indicating the results observed for the affinity matured variant, N35E, is IL-8 specific.

O. CONSTRUCTION OF HUMANIZED 6G4V11N35E F(ab')₂ LEUCINE ZIPPER

A F(ab')₂ expression plasmid for 6G4V11N35E was constructed using methods similar to those described in Section (K) above. The expression plasmid, p6G4V11N35E.F(ab')₂, was made by digesting the plasmid p6G4V11N35A.F(ab')₂ (described in Section (K) above) with the restriction enzymes Apal and NdeI to isolate a 2805 bp fragment encoding the heavy chain constant domain -GCN4 leucine zipper and ligating it to a 3758 bp Apal-NdeI fragment of the pPH6G4V11N35E phage display clone (encoding 6G4V11N35E Fab) obtained as described in Section (M) above. The integrity of the entire coding sequence was confirmed by DNA sequencing.

25 P. CONSTRUCTION OF THE FULL LENGTH HUMANIZED 6G4V11N35A IgG EXPRESSION PLASMID

The full length IgG₁ version of the humanized anti-IL8 variant 6G4V11N35A was made using a dicistronic DHFR-Intron expression vector (Lucas et al., Nucleic Acids Res.,24: 1774-1779 (1996)) which contained the full length recombinant murine-human chimera of the 6G4.2.5 anti-IL8 mAb. The expression plasmid encoding the humanized variant 6G4V11N35A was assembled as follows. First an intermediate plasmid (pSL-3) was made to shuttle the sequence encoding the variable heavy chain of humanized anti-IL-8 variant 6G4V11N35A to pRK56G4chim.2Vh - which contains the variable heavy region of the chimeric 6G4.5 anti-IL8 antibody. The vector pRK56G4chim.Vh was digested with PvuII and Apal to remove the heavy chain variable region of the chimeric antibody and religated with an 80bp PvuII - XhoI synthetic oligonucleotide (encoding Leu4 to Phe29 of 6G4V11N35A) (Fig. 47) and a 291bp XhoI - Apal fragment from p6G4V11N35A.7 carrying the remainder of the variable heavy chain sequence of 6G4V11N35A to create pSL-

3. This intermediate plasmid was used in conjunction with 2 other plasmids, p6G4V11N35A.F(ab')₂ and p6G425chim2.choSD, to create the mammalian expression plasmid, p6G4V11N35AchoSD.9 (identified as p6G425V11N35A.choSD in a deposit made on December 16, 1997 with the ATCC and assigned ATCC Accession No. 209552). This expression construct was assembled in a 4-part ligation using the following DNA fragments: a 5,203bp ClaI - BlnI fragment encoding the regulatory elements of the mammalian expression plasmid (p6G425 chim2.choSD), a 451bp ClaI - ApaI fragment containing the heavy chain variable region of the humanized 6G4V11N35A antibody (pSL-3), a 1,921bp ApaI - EcoRV fragment carrying the heavy chain constant region of 6G4V11N35A (p6G425chim2.choSD) and a 554bp EcoRV - BlnI fragment encoding the light chain variable and constant regions of 6G4V11N35A (p6G4V11N35A.F(ab')₂). The DNA sequence (SEQ ID NO: 68) of clone p6G4V11N35A.choSD.9 was confirmed by DNA sequencing and is presented in Figure 48.

Q. CONSTRUCTION OF THE FULL LENGTH HUMANIZED 6G4V11N35E IgG EXPRESSION PLASMID

A mammalian expression vector for the humanized 6G4V11N35E was made by swapping the light chain variable region of 6G4V11N35A with 6G4V11N35E as follows: a 7,566bp EcoRV - BlnI fragment (void of the 554bp fragment encoding the light chain variable region of 6G4V11N35A) from p6G4V11N35A.choSD.9 was ligated to a 554bp EcoRV - BlnI fragment (encoding the light chain variable region of 6G4V11N35E) from pPH6G4V11N35E.7. The mutation at position N35 of the light chain of p6G4V11N35E.choSD.10 was confirmed by DNA sequencing.

R. STABLE CHO CELL LINES FOR VARIANTS N35A AND N35E

For stable expression of the final humanized IgG1 variants (6G4V11N35A and 6G4V11N35E), Chinese hamster ovary (CHO) DP-12 cells were transfected with the above-described dicistronic vectors (p6G4V11N35A.choSD.9 and p6G4V11N35E.choSD.10, respectively) designed to coexpress both heavy and light chains (Lucas et al., *Nucleic Acid Res.* 24:1774-79 (1996)). Plasmids were introduced into CHO DP12 cells via lipofection and selected for growth in GHT-free medium (Chisholm, V. High efficiency gene transfer in mammalian cells. In: Glover, DM, Hames, BD. *DNA Cloning 4. Mammalian systems*. Oxford Univ. Press, Oxford pp 1-41 (1996)). Approximately 20 unamplified clones were randomly chosen and reseeded into 96 well plates. Relative specific productivity of each colony was monitored using an ELISA to quantitate the full length human IgG accumulated in each well after 3 days and a fluorescent dye, Calcein AM, as a surrogate marker of viable cell number per well. Based on these data, several unamplified clones were chosen for further amplification in the presence of increasing concentrations of methotrexate. Individual clones surviving at 10, 50, and 100 nM methotrexate were chosen and transferred to 96 well plates for productivity screening. One clone for each antibody (clone#1933 aIL8.92 NB 28605/12 for 6G4V11N35A; clone#1934 aIL8.42 NB 28605/14 for 6G4V11N35E), which reproducibly exhibited high specific productivity, was expanded in T-flasks and used to inoculate a spinner culture. After several passages, the suspension-adapted cells were used to inoculate production cultures in GHT-containing, serum-free media supplemented with various hormones and protein hydrolysates. Harvested cell culture fluid containing recombinant humanized anti-IL8 was purified using protein A-Sepharose CL-4B. The purity after this step was approximately 99%. Subsequent purification

to homogeneity was carried out using an ion exchange chromatography step. Production titer of the humanized 6G4V11N35E IgG1 antibody after the first round of amplification and 6G4V11N35A IgG1 after the second round of amplification were 250mg/L and 150mg/L, respectively.

S. CHARACTERIZATION OF THE HUMANIZED 6G4V11N35A/E IgG VARIANTS

5 The humanized full length IgG variants of 6G4.2.5 were tested for their ability to inhibit ^{125}I -IL-8 binding and to neutralize activation of human neutrophils; the procedures are described in Sections (B)(1) and (B)(2) above. As shown in Figure 49, the full length IgG1 forms of variants 6G4V11N35A and 6G4V11N35E equally inhibited ^{125}I -IL-8 binding to human neutrophils with approximate IC_{50} 's of 0.3nM and 0.5nM, respectively. This represents a 15 - 25 fold improvement in blocking binding of IL-8 compared to the full
10 length murine mAb ($\text{IC}_{50} = 7.5\text{nM}$). Similarly, the two anti-IL-8 variants showed equivalent neutralizing capabilities with respect to inhibiting IL-8 mediated human neutrophil chemotaxis (Figures 50A-50B). The IC_{50} 's of 6G4V11N35A IgG1 and 6G4V11N35E IgG1 for human IL-8 were 4.0nM and 6.0nM, respectively, and for rabbit IL-8 were 4.0nM and 2.0nM, respectively. The irrelevant isotype control Fab (4D5) did not inhibit neutrophil migration.

15 The affinity for IL-8 of these variants relative to the murine 6G4.2.5 mAb was determined using KinExA as described in Section (M). Figure 51 shows the equilibrium constant (K_d) for the full length affinity matured variants 6G4V11N35E IgG1 and 6G4V11N35A IgG1 were approximately 49pM and 88pM, respectively. The K_d for 6G4V11N35A IgG1 was determined directly from the kinetic experiment. As reported with their respective Fabs, this improvement in affinity might be attributed to an approximate 2-fold
20 increase in the on-rate of 6G4V11N35E IgG1 ($k_a = 3.0 \times 10^6$) compared to that of 6G4V11N35A IgG1 ($k_a = 8.7 \times 10^5$). In addition, these results were confirmed by a competition radio-immune assay using iodinated human IL-8. 50pM of 6G4V11N35A IgG1 or 6G4V11N35E IgG1 was incubated for 2 hours at 25°C with 30-50pM of ^{125}I -IL-8 and varying concentrations (0 to 100nM) of unlabeled IL-8. The antibody-antigen mixture was then incubated for 1 hour at 4°C with 10ul of a 70% slurry of Protein-A beads (pre-blocked with 0.1%
25 BSA). The beads were briefly spun in a microcentrifuge and the supernatant discarded to remove the unbound ^{125}I -IL-8. The amount of ^{125}I -IL-8 specifically bound to the anti-IL-8 antibodies was determined by counting the protein-A pellets in a gamma counter. The approximate K_d values were similar to those determined by KinEXA. The average K_d for 6G4V11N35A IgG1 and 6G4V11N35E IgG1 were 54pM (18 -90pM) and 19pM (5-34pM), respectively (Figure 52).

30 T. CONSTRUCTION OF HUMANIZED 6G4V11N35A/E Fab's FOR MODIFICATION BY POLYETHYLENE GLYCOL

A Fab' expression vector for 6G4V11N35A was constructed by digesting p6G4V11N35A.F(ab')₂ with the restriction enzymes Apal and NdeI to remove the 2805 bp fragment encoding the human IgG₁ constant domain fused with the yeast GCN4 leucine zipper and replacing it with the 2683bp Apal-NdeI fragment from
35 the plasmid pCDNA.18 described in Eigenbrot et al., Proteins: Struct. Funct. Genet., 18: 49-62 (1994). The

pCDNA.18 Apal-NdeI fragment carries the coding sequence for the human constant IgG1 heavy domain, including the free cysteine in the hinge region that was used to attach the PEG molecule. The 3758bp Apal-NdeI fragment (encodes the light chain and heavy variable domain of 6G4V11N35A) isolated from p6G4V11N35A.F(ab')₂ was ligated to the 2683bp Apal-NdeI fragment of pCDNA.18 to create p6G4V11N35A.PEG-1. The integrity of the entire coding sequence was confirmed by DNA sequencing. The nucleotide and translated amino acid sequences of heavy chain constant domain with the cysteine in the hinge are presented in Figure 53.

A Fab' expression plasmid for 6G4V11N35E was made similarly by digesting pPH6G4V11N35E (from Section (O) above) with the restriction enzymes Apal and NdeI to isolate the 3758bp Apal-NdeI DNA fragment carrying the intact light chain and heavy variable domain of 6G4V11N35E and ligating it to the 2683 bp Apal-NdeI DNA fragment from p6G4V11N35A.PEG-1 to create p6G4V11N35E.PEG-3. The integrity of the entire coding sequence was confirmed by DNA sequencing.

Anti-IL-8 6G4V11N35A Fab' variant was modified with 20 kD linear methoxy-PEG-maleimide, 30 kD linear methoxy-PEG-maleimide, 40 kD linear methoxy-PEG-maleimide, or 40 kD branched methoxy-PEG-maleimide as described below. All PEG's used were obtained commercially from Shearwater Polymers, Inc.

a. MATERIALS AND METHODS

Fab'-SH Purification

A Fab'-SH antibody fragment of the affinity matured antibody 6G4V11N35A was expressed in *E. coli* grown to high cell density in the fermentor as described by Carter *et al.*, *Bio/Technology* 10, 163-167 (1992). Preparation of Fab'-SH fragments was accomplished by protecting the Fab'-SH fragments with 4',4'-dithiodipyridine (PDS), partially purifying the protected Fab'-PDS fragments, deprotect the Fab'-PDS with dithiothreitol (DTT) and finally isolate the free Fab'-SH by using gel permeation chromatography.

Protection of Fab'-SH with PDS

Fermentation paste samples were dissolved in 3 volumes of 20mM MES, 5mM EDTA, pH 6.0 containing 10.7mg of 4',4'-dithiodipyridine per gram fermentation paste, resulting in a suspension with a pH close to 6.0. The suspension was passed through a homogenizer followed by addition of 5% PEI (w/v), pH 6 to the homogenate to a final concentration of 0.25%. The mixture was then centrifuged to remove solids and the clear supernatant was conditioned to a conductivity of less than 3mS by the addition of cold water.

Partial purification of the Fab'-SH molecule using ion exchange chromatography

The conditioned supernatant was loaded onto an ABX (Baker) column equilibrated in 20 mM MES, pH 6.0. The column was washed with the equilibration buffer followed by elution of the Fab'-SH with a 15 column volume linear gradient from 20 mM MES, pH 6.0 to 20 mM MES, 350 mM sodium chloride. The column was monitored by absorbance at 280nm, and the eluate was collected in fractions.

Deprotection of the Fab'-SH antibody fragments with DTT

The pH of the ABX pool was adjusted to 4.0 by the addition of dilute HCl. The pH adjusted solution was then deprotected by adding DTT to a final concentration of 0.2mM. The solution was incubated for about 30 minutes and then applied to a gel filtration Sephadex G25 column, equilibrated with 15mM sodium

phosphate, 25mM MES, pH 4.0. After elution, the pH of the pool was raised to pH 5.5 and immediately flash frozen at -70°C for storage or derivatized with PEG-MAL as described below.

Alternative Fab'-SH Purification

Alternatively Fab'-SH fragments can be purified using the following procedure. 100 g fermentation paste is thawed in the presence of 200 ml 50 mM acetic acid, pH 2.8, 2 mM EDTA, 1 mM PMSF. After mixing vigorously for 30 min at room temperature, the extract is incubated with 100 mg hen egg white lysozyme. DEAE fast flow resin (approximately 100 mL) is equilibrated with 10 mM MES, pH 5.5, 1 mM EDTA on a sintered glass funnel. The osmotic shock extract containing the Fab'-SH fragment is then filtered through the resin.

A protein G Sepharose column is equilibrated with 10 mM MES, pH 5.5, 1 mM EDTA and then loaded with the DEAE flow-through sample. The column is washed followed by three 4 column volume washes with 10 mM MES, pH 5.5, 1 mM EDTA. The Fab'-SH antibody fragment containing a free thiol is eluted from the column with 100 mM acetic acid, pH 2.8, 1 mM EDTA. After elution, the pH of the pool is raised to pH 5.5 and immediately flash frozen at -70°C for storage or derivatized with PEG-MAL as described below.

Preparation of Fab'-S-PEG

The free thiol content of the Fab'-SH preparation obtained as described above was determined by reaction with 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB) analysis according to the method of Creighton in Protein Structure: A Practical Approach, Creighton, T.E., ed, IRL Press (Oxford, UK: 1990), pp. 155-167. The concentration of free thiol was calculated from the increase on absorbance at 412 nm, using $\epsilon_{412} = 14,150 \text{ cm}^{-1} \text{ M}^{-1}$ for the thionitrobenzoate anion and a $M_r = 48,690$ and $c_{280} = 1.5$ for the Fab'-SH antibody. To the Fab'-SH protein G Sepharose pool, or the deprotected Fab'-SH gel permeation pool, 5 molar equivalents of PEG-MAL were added and the pH was immediately adjusted to pH 6.5 with 10% NaOH.

The Fab'-S-PEG was purified using a 2.5 x 20 cm cation exchange column (Poros 50-HS). The column was equilibrated with a buffer containing 20 mM MES, pH 5.5. The coupling reaction containing the PEGylated antibody fragment was diluted with deionized water to a conductivity of approximately 2.0 mS. The conditioned coupling reaction was then loaded onto the equilibrated Poros 50 HS column. Unreacted PEG-MAL was washed from the column with 2 column volumes of 20 mM MES, pH 5.5. The Fab'-S-PEG was eluted from the column using a linear gradient from 0 to 400 mM NaCl, in 20 mM MES pH 5.5, over 15 column volumes.

Alternatively a Bakerbond ABX column can be used to purify the Fab'-S-PEG molecule. The column is equilibrated with 20 mM MES, pH 6.0 (Buffer A). The coupling reaction is diluted with deionized water until the conductivity equaled that of the Buffer A (approximately 2.0 mS) and loaded onto the column. Unreacted PEG-MAL is washed from the column with 2 column volumes of 20 mM MES, pH 6.0. The Fab'-S-PEG is eluted from the column using a linear gradient from 0 to 100 mM $(\text{NH}_4)_2\text{SO}_4$, in 20 mM MES pH 6.0, over 15 column volumes.

Size Exclusion Chromatography

The hydrodynamic effective size of each molecule was determined using a Pharmacia Superose-6 HR 10/30 column (10x300mm). The mobile phase was 200 mM NaCl, 50 mM sodium phosphate pH 6.0. Flow rate was at 0.5 ml/min and the column was kept at ambient temperature. Absorbance at 280 nm was monitored where PEG contributed little signal. Biorad MW standards containing cyanocobalamin, myoglobin, ovalbumin, IgG, Thyroglobulin monomer and dimer were used to generate a standard curve from which the effective size of the pegylated species was estimated.

b. RESULTS

Size Exclusion Chromatography

The effective size of each modified species was characterized using size exclusion chromatography. The results are shown in Fig. 60 below. The theoretical molecular weight of the anti-IL8 Fab fragments modified with PEG 5kD, 10kD, 20kD, 30kD, 40kD (linear), 40kD (branched) or 100,000kD is shown along with the apparent molecular weight of the PEGylated fragments obtained by HPLC size exclusion chromatography. When compared to the theoretical molecular weight of the Fab'-S-PEG fragments, the apparent molecular weight (calculated by size exclusion HPLC) increases dramatically by increasing the size of the PEG attached to the fragments. Attachment of a small molecular weight PEG, for example PEG 10,000D only increases the theoretical molecular weight of the PEGylated antibody fragment (59,700 D) by 3 fold to an apparent molecular weight of 180,000D. In contrast attachment of a larger molecular weight PEG for example 100,000D PEG to the antibody fragment increases the theoretical molecular weight of the PEGylated antibody fragment (158,700 D) by 12 fold to an apparent molecular weight of 2,000,000D.

SDS-PAGE

In Fig. 61, the upper panel shows the size of the anti-IL-8 Fab fragments modified with PEG of molecular weight 5kD (linear), 10kD (linear), 20kD (linear), 30kD (linear), 40kD (linear), 40kD (branched) or 100kD (linear) under reduced conditions. The unmodified Fab is shown in lane 2 from right to left. Both the heavy and light chains of the Fab had a molecular weight of approximately 30kD as determined by PAGE. Each PEGylated fragment sample produced two bands: (1) a first band (attributed to the light chain) exhibiting a molecular weight of 30kD; and (2) a second band (attributed to the heavy chain to which the PEG is attached specifically at the hinge SH) exhibiting increasing molecular weights of 40, 45, 70, 110, 125, 150 and 300kD. This result suggested that PEGylation was specifically restricted to the heavy chain of the Fab's whereas the light chain remained unmodified.

The lower panel is non-reduced PAGE showing the size of the anti-IL-8 Fab fragments modified with PEG of molecular weight 5kD (linear), 20kD (linear), 30kD (linear), 40kD (linear), 40kD (branched), or 100kD (linear). The PEGylated fragments exhibited molecular weights of approximately 70kD, 115kD, 120kD, 140kD, 200kD and 300kD.

The SDS PAGE gels confirm that all Fab'-S-PEG molecules were purified to homogeneity and that the molecules differed only with respect to the size of the PEG molecule attached to them.

U. AMINE SPECIFIC PEGYLATION OF ANTI-IL-8 F(ab')₂ FRAGMENTS

Pegylated F(ab')₂ species were generated by using large MW or branched PEGs in order to achieve a large effective size with minimal protein modification which might affect activity. Modification involved N-

hydroxysuccinamide chemistry which reacts with primary amines (lysines and the N-terminus). To decrease the probability of modifying the N-terminus, which is in close proximity to the CDR region, a reaction pH of 8, rather than the commonly used pH of 7, was employed. At pH 8.0, the amount of the reactive species (charged NH_3^+) would be considerably more for the ϵ -NH₂ group of lysines ($\text{pK}_a=10.3$) than for the α -NH₂ group (pK_a

5 of approximately 7) of the amino-terminus. For the linear PEGs, a methoxy-succinimidyl derivative of an NHS-PEG was used because of the significantly longer half-life in solution (17 minutes at 25°C at pH 8.0) compared to the NHS esters of PEGs (which have 5-7 minute half life under the above conditions). By using a PEG that is less prone to hydrolysis, a greater extent of modification is achieved with less PEG. Branched PEGs were used to induce a large increase in effective size of the antibody fragments.

10 a. MATERIALS

All PEG reagents were purchased from Shearwater Polymers and stored at -70°C in a desiccator: branched N-hydroxysuccinamide-PEG (PEG2-NHS-40KDa) has a 20 kDa PEG on each of the two branches, methoxy-succinimidyl-propionic acid-PEG (M-SPA-20000) is a linear PEG molecule with 20 kDa PEG. Protein was recombinantly produced in *E. coli* and purified as a (Fab)₂ as described in Sections (K) and (O)

15 above.

b. METHODS

IEX method: A J. T. Baker Wide-Pore Carboxy-sulfone (CSX), 5 micron, 7.75 x 100 mm HPLC column was used for fractionation of the different pegylated products, taking advantage of the difference in charge as the lysines are modified. The column was heated at 40°C. A gradient as shown in Table 7 below was

20 used where Buffer A was 25 mM sodium Borate/25 mM sodium phosphate pH 6.0, and Buffer B was 1 M ammonium sulfate, and Buffer C was 50 mM sodium acetate pH 5.0.

Table 7

	Time (min)	%B	%C	flow mL/min
25	0	10	10	1.5
	20	18	7.5	1.5
	25	25	7.5	1.5
	27	70	3.0	2.5
	29	70	3.0	2.5
30	30	10	10	2.5
	33	10	10	2.5

SEC-HPLC: The hydrodynamic or effective size of each molecule was determined using a Pharmacia Superose-6 HR 10/30 column (10x300mm). The mobile phase was 200 mM NaCl, 50 mM sodium phosphate pH 6.0. Flow rate was at 0.5 mL/min and the column was kept at ambient temperature. Absorbance

35 at 280 nm was monitored where PEG contributed little signal. Biorad MW standards containing cyanocobalamin, myoglobin, ovalbumin, IgG, Thyroglobulin monomer and dimer were used to generate a standard curve from which the effective size of the pegylated species was estimated.

SEC-HPLC-Light Scattering: For determination of the exact molecular weight, this column was

40 connected to an on-line light scattering detector (Wyatt Minidawn) equipped with three detection angles of 50°, 90°, and 135° C. A refractive index detector (Wyatt) was also placed on-line to determine concentration. All

buffers were filtered with Millipore 0.1 μ filters; in addition a 0.02 μ Whatman Anodisc 47 was placed in-line prior to the column.

The intensity of scattered light is directly proportional to the molecular weight (M) of the scattering species, independent of shape, according to:

$$M = R_0 / K \cdot c$$

where R_0 is the Rayleigh ratio, K is an optical constant relating to the refractive index of the solvent, the wavelength of the incident light, and dn/dc , the differential refractive index between the solvent and the solute with respect to the change in solute concentration, c. The system was calibrated with toluene (R_0 of 1.406×10^{-5} at 632.8 nm); a dn/dc of 0.18, and an extinction coefficient of 1.2 was used. The system had a mass accuracy of ~5%.

SDS-PAGE: 4-12% Tris-Glycine Novex minigels were used along with the Novex supplied Tris-Glycine running buffers. 10-20 μ g of protein was applied in each well and the gels were run in a cold box at 150 mV/gel for 45 minutes. Gels were then stained with colloidal Coomassie Blue (Novex) and then washed with water for a few hours and then preserved and dried in drying buffer (Novex)

Preparation of a linear(1)20KDa-(N)-(Fab')₂: A 4 mg/ml solution of anti-IL8 formulated initially in a pH 5.5 buffer was dialyzed overnight against a pH 8.0 sodium phosphate buffer. 5 mL protein was mixed at a molar ratio of 3:1. The reaction was carried out in a 15 mL polypropylene Falcon tube and the PEG was added while vortexing the sample at low speed for 5 seconds. It was then placed on a nutator for 30 minutes. The extent of modification was evaluated by SDS-PAGE. The whole 5 ml reaction mixture was injected on the IEX for removal of any unreacted PEG and purification of singly or doubly pegylated species. The above reaction generated a mixture of 50% singly-labeled anti-IL8. The other 50% unreacted anti-IL8 was recycled through the pegylation/purification steps. The pooled pegylated product was dialyzed against a pH 5.5 buffer for in vitro assays and animal PK studies. Endotoxin levels were measured before administration to animals or for the cell based assays. Levels were below 0.5 eu/ml. The fractions were also run on SDS-PAGE to confirm homogeneity. Concentration of the final product was assessed by absorbance at 280 nm using an extinction coefficient of 1.34, as well as by amino acid analysis.

Preparation of a branched(1)40KDa-(N)-(Fab')₂: A 4 mg/mL solution of anti-IL8 (Fab')₂ formulated in a pH 5.5 buffer was dialyzed overnight against a pH 8.0 phosphate buffer. Solid PEG powder was added to 5 mL protein in two aliquots to give a final PEG:protein molar ratio of 6:1. Each solid PEG aliquot was added to the protein in a 15 mL polypropylene Falcon tube while vortexing at low speed for 5 sec, and then placing the sample on a nutator for 15 minutes. The extent of modification was evaluated by SDS-PAGE using a 4-12% Tris-Glycine (Novex) gel and stained with colloidal Coomassie blue (Novex). The 5 mL PEG-protein mixture was injected on the ion exchange column for removal of any unreacted PEG. The above reaction generated a mixture of unreacted (37%), singly-labelled (45%), doubly and triply-labeled (18%) species. These were the optimal conditions for obtaining the greatest recovery of the protein with only 1 PEG per antibody rather than the higher molecular weight adducts. The unmodified anti-IL8 was recycled. The pegylated products were separated and fractionated in falcon tubes and then dialyzed against a pH 5.5 buffer

for assays and animal PK studies. Endotoxin levels were below 0.5 eu/ml. The fractions were also run on SDS-PAGE to confirm homogeneity. The concentration of the final product was assessed by absorbance at 280 nm using an extinction coefficient of 1.34, as well as by amino acid analysis.

5 Preparation of branched(2)-40KDa-(N)(Fab')₂: This molecule was most efficiently made by adding three times in 15 minute intervals a 3:1 molar ratio of PEG to the already modified branched(1)-40KDa-(N)-(Fab')₂. The molecule was purified on IEX as 50% branched(2)-40KDa-(N)-(Fab')₂. The unmodified molecule was recycled until ~20 mg protein was isolated for animal PK studies. The product was characterized by SEC-light scattering and SDS-PAGE.

c. RESULTS

10 PEGs increased the hydrodynamic or effective size of the product significantly as determined by gel filtration (SEC-HPLC). Figure 62 shows the SEC profile of the pegylated F(ab')₂ species with UV detection at 280 nm. The hydrodynamic size of each molecule was estimated by reference to the standard MW calibrators.

As summarized in Figure 62, the increase in the effective size of (Fab')₂ was about 7-fold by adding one linear 20 kDa PEG molecule and about 11-fold by adding one branched ("Br(1)") 40 kDa PEG molecule, and
15 somewhat more with addition of two branched ("Br(2)") PEG molecules

Light scattering detection gave the exact molecular weight of the products and confirmed the extent of modification (Figure 63). The homogeneity of the purified material was shown by SDS-PAGE (Figure 64). Underivatized F(ab')₂ migrated as a 120 kDa species, the linear(1)20KDa-(N)-F(ab')₂ migrated as a band at 220kDa, the Br(1)-40KD(N)-F(ab')₂ migrated as one major band at 400 kDa, and the Br(2)-40KD-(N)-F(ab')₂
20 migrated as a major band at around 500 kDa. The proteins appeared somewhat larger than their absolute MW due to the steric effect of PEG.

V. IN VITRO ACTIVITY CHARACTERIZATION OF PEG MODIFIED Fab' FRAGMENTS OF 6G4V11N35A (MALEIMIDE CHEMICAL COUPLING METHOD)

Anti-IL-8 6G4V11N35A Fab' variants modified with 5-40kD linear PEG molecules and a 40kD
25 branched PEG molecule were tested for their ability to inhibit both IL-8 binding and activation of human neutrophils; the procedures were described in Sections (B)(1), (B)(2) and (B)(3) above. The binding curves and IC₅₀'s for PEG-maleimide modified 6G4V11N35A Fab' molecules are presented in Figures 54A-54C. The IC₅₀ of the 5kD pegylated Fab' (350pM) and the average IC₅₀ of the Fab control (366pM) were not significantly different, suggesting that the addition of a 5kD MW PEG did not affect the binding of IL-8 to the modified Fab' (Figure 54A). However, a decrease in the binding of IL-8 to the 10kD and 20kD pegylated
30 Fab' molecules was observed as depicted by the progressively higher IC₅₀'s (537pM and 732pM, respectively) compared to the average IC₅₀ of the native Fab. These values represent only a minimal loss of binding activity (between 1.5- and 2.0-fold). A less pronounced difference in IL-8 binding was observed for the 30kD and 40kD linear PEG antibodies (Figure 54B). The IC₅₀'s were 624pM and 1.1nM, respectively,
35 compared to the 802pM value of the Fab control. The 40kD branched PEG Fab' showed the largest decrease

in IL-8 binding (2.5 fold) relative to the native Fab (Figure 54C). Nevertheless, the reduction in binding of IL-8 by these pegylated Fab's is minimal.

The ability of the pegylated antibodies to block IL-8 mediated activation of human neutrophils was demonstrated using the PMN chemotaxis (according to the method described in Section B(2) above) and β -glucuronidase release (according to the method described in Lowman et al., *J. Biol. Chem.*, 271: 14344 (1996)) assays. The IC_{50} 's for blocking IL-8 mediated chemotaxis are shown in Figures 55A-55C. The 5-20kD linear pegylated Fab' antibodies were able to block IL-8 mediated chemotaxis within 2-3 fold of the unpegylated Fab control (Figure 55A). This difference is not significant because the inherent variation can be up to 2 fold for this type of assay. However, a significant difference was detected for the 30kD and 40kD linear pegylated Fab' antibodies as illustrated by the higher IC_{50} 's of the 30kD linear PEG-Fab' (2.5nM) and 40kD linear PEG-Fab' (3.7nM) compared to the Fab control (0.8nM) (Figure 55B). The ability of the 40kD branched PEG Fab' molecule to block IL-8 mediated chemotaxis was similar to that of the 40kD linear PEG Fab' (Figure 55C). At most, the ability of the pegylated Fab' antibodies to block IL-8 mediated chemotaxis was only reduced 2-3 fold. Furthermore, release of β -glucuronidase from the granules of neutrophils was used as another criteria for assessing IL-8 mediated activation of human PMNs. Figure 56A (depicting results obtained with 5 kD, 10 kD and 20 kD linear PEGs), Figure 56B (depicting results obtained with 30 kD and 40 kD linear PEGs), and Figure 56C (depicting results obtained with 40 kD branched PEG) show that all the pegylated Fab' antibodies were able to inhibit IL-8 mediated release of β -glucuronidase as well as or better than the unpegylated Fab control. The data collectively shows that the pegylated Fab' variants are biological active and are capable of inhibiting high amounts of exogenous IL-8 in in-vitro assays using human neutrophils.

W. IN VITRO ACTIVITY CHARACTERIZATION OF PEG MODIFIED $F(ab')_2$ FRAGMENTS OF 6G4V11N35A (SUCCINIMIDYL CHEMICAL COUPLING METHOD)

The anti-IL-8 variant 6G4V11N35A $F(ab')_2$ modified with (a) a single 20kD linear PEG molecule per $F(ab')_2$, (b) a single 40kD branched PEG molecule per $F(ab')_2$, (c) with three, four, or five 20 kD linear PEG molecules per $F(ab')_2$ (a mixture of: (1) species having three 20 kD linear PEG molecules per $F(ab')_2$; (2) species having four 20 kD linear PEG molecules per $F(ab')_2$; and (3) species having five 20 kD linear PEG molecules per $F(ab')_2$; denoted as "20 kD linear PEG (3,4,5) $F(ab')_2$ "), or (d) with two 40kD branched PEG molecules per $F(ab')_2$ (denoted as "40 kD branch PEG (2) $F(ab')_2$ "), were tested for their ability to inhibit ^{125}I -IL-8 binding and to neutralize activation of human neutrophils. The procedures used are described in Sections (B)(1), (B)(2) and (B)(3) above. The binding curves for pegylated $F(ab')_2$ variants are shown in Figures 57A-57B. No significant differences were observed amongst the $F(ab')_2$ control, the single 20kD linear PEG-modified $F(ab')_2$, and the single 40kD branched PEG-modified $F(ab')_2$ (Figure 57A). However, the $F(ab')_2$ variants containing multiple PEG molecules showed a slight reduction (less than 2-fold) in their ability to bind

IL-8. The IC_{50} 's of the 20kD linear PEG (3,4,5) $F(ab')_2$ and 40kD branch PEG (2) $F(ab')_2$ variants were 437pM and 510pM, respectively, compared to 349pM of the $F(ab')_2$ control (Figure 57B).

The ability of these pegylated $F(ab')_2$ variants to block IL-8 mediated neutrophil chemotaxis is presented in Figures 58A-58B. Consistent with the PMN binding data, the single linear and branched PEG $F(ab')_2$ variants were able to block IL-8 mediated chemotaxis similar to the unpegylated $F(ab')_2$ control (Figure 58A). The ability of the 40kD branch PEG (2) $F(ab')_2$ variant to inhibit PMN chemotaxis was identical to the control $F(ab')_2$ while the 20kD linear PEG (3,4,5) $F(ab')_2$ mixture was able to inhibit within 3-fold of the control antibody (Figure 58B).

Shown in Figures 59A and 59B are the results of the β -glucuronidase release assay which is a measure of degranulation by IL-8 stimulated human neutrophils. The single 20kD linear PEG-modified $F(ab')_2$ and the single 40kD branched PEG-modified $F(ab')_2$ variants were able to inhibit release of β -glucuronidase as well as the $F(ab')_2$ control (Figure 59A). The 40kD branch PEG (2) $F(ab')_2$ inhibited this response within 2-fold of the $F(ab')_2$ control (Figure 59B). The 20kD linear PEG (3,4,5) molecule was not tested. Overall, the $F(ab')_2$ pegylated anti-IL-8 antibodies were biologically active and effectively prevented IL-8 binding to human neutrophils and the signaling events leading to cellular activation.

X. PHARMACOKINETIC AND SAFETY STUDY OF EIGHT CONSTRUCTS OF PEGYLATED ANTI-IL-8 (HUMANIZED) $F(ab')_2$ AND $F(ab')$ FRAGMENTS IN NORMAL RABBITS FOLLOWING INTRAVENOUS ADMINISTRATION

The objective of this study was to evaluate the effect of pegylation on the pharmacokinetics and safety of six pegylated humanized anti-IL-8 constructs (pegylated 6G4V11_{1-35A}. Fab' and pegylated 6G4V11N35A. $F(ab')_2$ obtained as described in Sections (T) and (U) above) relative to the non-pegylated fragments in normal rabbits. Eight groups of two/three male rabbits received equivalent protein amounts of pegylated 6G4V11N35A. Fab' or pegylated 6G4V11N35A. $F(ab')_2$ constructs (2 mg/kg) via a single intravenous (IV) bolus dose of one anti-IL8 construct. Serum samples were collected according to the schedule shown in Table 8 below and analyzed for anti-IL8 protein concentrations and antibody formation against anti-IL8 constructs by ELISA.

Table 8

Group No.	Dose level/ Route	Material	Blood Collection
1		Fab' control	0,5,30 min; 1,2,3,4,6,8,10, 14,20,24,360 hr
2		linear(1)20K(s) Fab'	0,5,30 min; 1,2,4,6,8,10,12, 24,28,32,48,72,96,168,216,
3		linear(1)40K(s) Fab'	

4	2 mg/kg (protein conc.) IV bolus	branched(1)40K(N)F(ab') ₂	264,336,360 hr
5		F(ab') ₂ control	0,5,30 min; 1,2,4,6,8,10,12, 24,28,32,48,52,56,336 hr
6		branched(2)40K(s)Fab'	0,5,30 min; 1,2,4,6,8,10,12, 24,28,32,48,72,96,168,216,264,3 36 hr; Day 17,21, 25
7		branched(2)40K(N)F(ab') ₂	0,5,30 min; 1,2,4,6,8,10,12, 24,28,32,48,72,144,192, 240 hr; Day 13, 16, 20, 23
8		linear(1)30K(s)Fab'	0,5,30 min; 1,2,4,6,8,10,12, 24,28,32,48,72,96,168,216,264,3 36 hr; Day 17,21, 25

a. METHODS

Three male New Zealand White (NZW) rabbits per group (with exception to Group 7, n=2) received an equivalent amount of 6G4V11N35A protein (Fab' or F(ab')₂) construct at 2 mg/kg via an IV bolus dose in a marginal ear vein. Amino acid composition analysis and absorbance at 280 nm using extinction coefficients of 1.26 for 6G4V11N35A Fab' constructs and 1.34 for 6G4V11N35A F(ab')₂ constructs were performed to determine the protein concentration. Whole blood samples were collected via an ear artery cannulation (ear opposing dosing ear) at the above time points. Samples were harvested for serum and assayed for free 6G4V11N35A Fab' or F(ab')₂ constructs using an IL-8 Binding ELISA. Assays were conducted throughout the study as samples became available. All animals were sacrificed following the last blood draw, and necropsies were performed on all animals in Groups 1, 4–8. Due to the development of antibodies against the 6G4V11N35A constructs, non-compartmental pharmacokinetic analysis was conducted on concentration versus time data only up to 168 hours.

b. RESULTS

In four animals (Animals B, P, Q, V), interference to rabbit serum in the ELISA assay was detected (i.e. measurable concentrations of anti-IL8 antibodies at pre-dose). However, because these values were at insignificant levels and did not effect the pharmacokinetic analysis, the data were not corrected for this interference.

One animal (Animal G; Group 3) was exsanguinated before the termination of the study and was excluded from the pharmacokinetic analysis. At 4 hours, the animal showed signs of a stroke that was not believed to be drug related, as this can occur in rabbits following blood draws via ear artery cannulation.

The mean concentration–time profiles of the eight anti-IL8 constructs in normal rabbits are depicted in Fig. 65, and the pharmacokinetic parameters for the eight constructs are summarized in Table 9 below.

Significant antibodies to the anti-IL-8 constructs were present at Day 13/14 in all dose groups except Group 1 (Fab' control).

Table 9. Pharmacokinetic parameters.

Molecule	Fab'					F(ab') ₂		
Group No.	1	2	8	3	6	5	4	7
PEG structure	—	linear	linear	linear	branched	—	branched	branched
Number of PEGs	—	1	1	1	1	—	1	2
PEG MW	—	20K	30K	40K	40K	—	40K	40K
Dose (mg/kg)	2	2	2	2	2	2	2	2
V _c (mL/kg) ^a	58±3	36±3	35±1	34	44±1	45±5	36±1	32
V _{ss} (mL/kg) ^b	68±8	80±8	110±15	79	88±21	59±4	50±3	52
C _{max} (µg/mL) ^c	35±1	58±3	57±1	60	45±1	45±6	56±2	62
T _{max} (min) ^d	5	5	5	5	5	5	5	5
t _{1/2} term (hr) ^e	3.0±0.9	44±2	43±7	50	105±11	8.5±2.1	45±3	48
AUC ₀₋ (hr·µg/mL) ^f	18±3	80±74	910±140	1600	3400±1300	140±3	2200±77	2500
CL (mL/hr/kg) ^g	110±17	2.5±0.2	2.2±0.4	1.3	0.63±0.20	14±0	0.92±0.03	0.83
MRT (hr) ^h	0.61±0.15	32±2	45±9	63	140±18	4.2±0.3	55±3	64
No. of Animals	3	3	3	2	3	3	3	2

^a Initial volume of distribution.

^b Volume of distribution at steady state.

^c Observed maximum concentration.

^d Observed time to C_{max}.

^e t_{1/2} term = half-life associated with the terminal phase of the concentration vs. time profile.

^f Area under the concentration versus time curve (extrapolated to infinity).

^g CL = serum clearance.

^h MRT = Mean residence time.

The initial volume of distribution approximated the plasma volume for both the Fab' and F(ab')₂.

Pegylation decreased serum CL of anti-IL8 fragments and extended both the terminal half-life and MRT as shown in Table 10 below.

Table 10. Fold decrease/increase in clearance, terminal half-life & MRT of pegylated anti-IL8 fragments.

anti-IL8 fragment	Fab'					F(ab') ₂		
Group No.	1	2	8	3	6	5	4	7
PEG structure	—	linear	linear	linear	bran.	—	bran.	bran.
No. of PEGs	—	1	1	1	1	—	1	2
PEG MW	—	20K	30K	40K	40K	—	40K	40K
CL : mean (mL/hr/kg)	110	2.5	2.2	1.3	0.63	14	0.92	0.83
fold decrease	1	46	51	90	180	1	15	17
t _{1/2} term : mean (hr)	3.0	44	43	50	110	8.5	45	48
fold increase	1	14	14	17	35	1	5.3	5.7
MRT : mean (hr)	0.61	32	45	63	140	4.2	55	64
fold increase	1	53	73	100	240	1	13	15

For the pegylated anti-IL8 Fab' fragments, CL decreased by 46 to 180-fold. Terminal half-life and MRT increased 14 to 35-fold and 53 to 240-fold, respectively. For pegylated anti-IL8 F(ab')₂ molecules, CL decreased 15 to 17-fold with pegylation, and terminal half-life and MRT increased by greater than 5-fold and 13-fold, respectively. The changes in these parameters increased for both pegylated Fab' and F(ab')₂ molecules with increasing PEG molecular weight and approached the values of the full-length anti-IL8 (terminal half-life of 74 hours, MRT of 99 hours and CL of 0.47 mL/hr/kg). In comparing the branched(1)40K Fab' (Group 6) and branched(1)40K F(ab')₂ (Group 4), unexpected pharmacokinetics were observed. The pegylated Fab' molecule appeared to remain in the serum longer than the pegylated F(ab')₂ (see Figure 66). The mean CL of branched(1)40K Fab' was 0.63 mL/hr/kg, but a higher CL was observed for branched(1)40kD F(ab')₂ (CL 0.92 mL/hr/kg). The terminal half-life, likewise, was longer for the Fab' than the F(ab')₂ pegylated molecule (110 vs 45 hours).

The pharmacokinetic data demonstrated that pegylation decreased CL and increased terminal t_{1/2} and MRT of anti-IL8 fragments (Fab' and F(ab')₂) to approach that of the full-length anti-IL8. Clearance was decreased with pegylation 46 to 180-fold for the Fab' and approximately 16-fold for the F(ab')₂. The terminal half-life of the Fab' anti-IL8 fragment was increased by 14 to 35-fold and approximately 5-fold for the F(ab')₂ anti-IL8. MRT, likewise, were extended by 53 to 240-fold for the Fab' and approximately 14-fold for the F(ab')₂. The branched(1) 40kD Fab' had a longer terminal half-life and lower clearance compared to the branched(1) 40kD F(ab')₂.

Y. IN VIVO EFFICACY TESTING OF ANTI-IL-8 ANTIBODY REAGENTS IN RABBIT MODEL OF ISCHEMIA/REPERFUSION AND ACID ASPIRATION-INDUCED ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

Full length murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5, 40 kD branched PEG-6G4V11N35A Fab', and control antibody (anti-HIV gp120 monoclonal antibody 9E3.1F10) were tested in a rabbit ARDS model. The animals were weighed and anaesthetized by intramuscular injection of ketamine (50 mg/kg body weight), xylazine (5 mg/kg body weight), and acepromazine (0.75 mg/kg body weight). A second dose (20% of the first dosage) was given IM 15 minutes before removal of vascular clip, and third dose (60% of the first dosage) was given at tracheotomy. Intra-arterial catheter (22G, 1 in. Angiocath) and intra-venous catheter (24G, 1 in. angiocath) were be placed in the ear central artery and posterior marginal ear vein for blood samplings (arterial blood gases and CBC) and anti-IL-8 and fluid administration, respectively. The anaesthetized animals were transferred in a supine position to an operating tray; the abdominal area was shaved and prepared for surgery. Via a midline laparotomy, the superior mesenteric artery (SMA) was isolated and a microvascular arterial clip applied at the aortic origin. Before the temporary closure of the abdomen using 9 mm wound clip (Autoclip, Baxter), 15 ml of normal saline was given intraperitoneally as fluid supplement. After 110 minutes of intestinal ischemia, the abdominal incision was reopened and the arterial clip was released to allow reperfusion. Before closure, 5 ml of normal saline was given intraperitoneally for fluid replacement.

The laparotomy incision was closed in two layers and the animals allowed to awaken.

After surgery, the animals were placed on a heating pad (38°C) and continuously monitored for up to 6 hours post reperfusion and lactated Ringer's 8-12 ml/kg/hr IV was given as fluid supplement.

At 22-24 hr post-reperfusion, a tracheotomy was performed under anesthesia. Normal physiologic saline was diluted 1:3 with water and adjusted to pH 7.5 (adjusted by using 1N HCL); 3 ml/kg body weight was then instilled intra-tracheally. Rectal temperature was maintained at 37 +/- 1 degree C using a homeothermic heat therapy pad (K-Mod II, Baxter). Fluid supplements (LRS) at a rate of 5 ml/kg/hour IV were given. Blood gases were monitored every hour. The rabbits were returned to the cage after 6 hr of continuous monitoring.

Just prior to aspiration, animals were treated with saline, the control monoclonal antibody (anti-HIV gp-120 IgG 9E3.1F10), the full length murine anti-rabbit IL8 (6g4.2.5 murine IgG2a anti-rabbit IL8) or the pegylated 6G4V11N35A Fab' (6G4V1N35A Fab' modified with 40kD branched PEG-maleimide as described in Section T above, denoted as "40 kD branched PEG-6G4V11N35A Fab' "). Data from saline or control antibody treated animals was combined and presented as "Control". Arterial blood gases and A-a PO2 gradient measurements were taken daily, and IV fluid supplementation was performed daily. A-a PO2 gradient was measured at 96 hr of reperfusion. The A-a PO2 gradient was calculated as:

$$A-a PO_2 = [FIO_2(PB - PH_2O) - (PaCO_2/RQ)] - PaO_2.$$

PaO2/FiO2 ratios were measured at 24hr and 48hr in room air and 100% oxygen.

After the final A-a PO2 gradient measurement, the animals were anesthetized with Nembutal 100mg/kg i.v. and the animals were euthanized by transecting the abdominal aorta in order to reduce red blood cell contamination of bronchoalveolar lavage fluid (BAL). The lungs were removed en bloc. The entire lung was weighed and then lavaged with an intratracheal tube (Hi-Lo tracheal tube, 3mm) using 30 ml of HBSS and lidocain. Total and differential leukocyte counts in the BAL were determined. Lesions/changes were verified by histological examination of each lobe of the right lung of each animal.

The gross lung weight, total leukocyte and polymorphonuclear cell counts in BAL, and PaO2/FiO2 data obtained are depicted in Figs. 67, 68 and 69, respectively. Treatment with 40 kD branched PEG-6G4V11N35A Fab' exhibited no effect on the biological parameters measured in the model as compared to the "Control" group. However, the data do not contradict the pharmacokinetic analysis or the in vitro activity analysis for the 40 kD branched PEG-6G4V11N35A Fab' presented in Sections (V) and (X) above. In addition, these data do not contradict the ability of the 40 kD branched PEG-6G4V11N35A Fab' to reach and act on disease effector targets in circulation or other tissues.

Z. ADDITIONAL IN VIVO EFFICACY TESTING OF ANTI-IL-8 ANTIBODY REAGENTS IN RABBIT MODEL OF ISCHEMIA/REPERFUSION AND ACID ASPIRATION-INDUCED ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

Full length murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 and 20 kD linear PEG-6G4V11N35E Fab' were tested in a rabbit model of ischemia/reperfusion- and acid aspiration-induced acute respiratory distress syndrome (ARDS).

Antibodies

A Fab'-SH antibody fragment of the affinity matured anti-IL-8 antibody 6G4V11N35E was expressed using the Fab' expression plasmid for 6G4V11N35E (described in Section (T) above) in *E. coli* grown to high

density in the fermentor as described by Carter et al., Bio/Technology, 10: 163-167 (1992). Anti-IL-8 6G4V11N35E Fab' variant was purified from fermentation paste and modified with 20 kD linear methoxy-PEG-maleimide as described in Example T above. Pegylated material was formulated in phosphate buffered saline (PBS) at physiological pH. Full length 6G4.2.5 antibody was obtained from hybridoma cell line 6G4.2.5 as described in Section (B) above and formulated in phosphate buffered saline (PBS) at physiological pH.

Sterile Surgical Procedures and Post-Operative Care

Male New Zealand White rabbits weighing 2.2 to 2.5 kg (obtained from Western Oregon Rabbit Company) were anaesthetized by intramuscular injection of ketamine (50 mg/kg body weight), xylazine (5 mg/kg body weight), and acepromazine (0.75 mg/kg body weight). Intra-arterial catheter (22G, 1 in. Angiocath) and intra-venous catheter (24G, 1 in. angiocath) were be placed in the ear central artery and posterior marginal ear vein for blood samplings (arterial blood gases and CBC) and anti-IL-8 (or fluid) administration, respectively. The anaesthetized animals were transferred in a supine position to an operating tray; the abdominal area was shaved and prepared for surgery. Via a midline laparotomy, the superior mesenteric artery (SMA) was isolated and a microvascular arterial clip applied at the aortic origin. Before the temporary closure of the abdomen using 9 mm wound clip (Autoclip, Baxter), 15 ml of normal saline (38°C) was given intraperitoneally as fluid supplement. After 110 minutes of intestinal ischemia, the abdominal incision was reopened and the arterial clip was released to allow reperfusion. Before closure, 5 ml of normal saline (38°C) was given intraperitoneally for fluid replacement. The laparotomy incision was closed in two layers and the animals allowed to awaken.

After surgery, the animals were placed on a heating pad (38°C) and continuously monitored for up to 6 hours post reperfusion and lactated Ringer's 8-12 ml/kg/hr IV was given as fluid supplement.

At 22-24 hr post-reperfusion, a tracheotomy was performed under anesthesia using ketamine, xylazine and acepromazine as described above. Normal physiologic saline was diluted 1:3 with water and adjusted to pH 1.5 (adjusted by using 1N HCL), and 3 ml/kg body weight was then instilled intra-tracheally through an uncuffed tracheal tube (2.0mm I.D., Mallinckrodt Medical, Inc.). After instillation, the trachea was closed with 3-0 silk suture and the rabbits were allowed to recover. Rectal temperature was maintained at 37°C +/- 1°C using a homeothermic heat therapy pad (K-Mod II, Baxter). Fluid supplements (LRS) at a rate of 5 ml/kg/hour IV were given. The rabbits were observed and blood gases in room air and in 100% oxygen were measured daily.

Dose Administration

Treated animals received an intravenous injection of 7 mg/kg 20 kD linear PEG-6G4V11N35E Fab' (n=5 animals) at 10 minutes before and 6 hours after acid instillation.

Oxygenation Measurement

Alveolar-arterial oxygen pressure gradient (A-a PO₂ gradient) was calculated as follows:

$$A-a PO_2 = [FiO_2(PB - PH_2O) - (PaCO_2/RQ)] - PaO_2$$

where FiO₂ is fraction of inspired oxygen, PB is barometric pressure, PH₂O is partial pressure of water vapor, PaCO₂ is arterial carbon dioxide pressure, RQ is respiratory quotient, and PaO₂ is arterial oxygen pressure.

A-a PO₂ gradient and PaO₂/FiO₂ ratios for each rabbit were measured at baseline (pre-op), before acid instillation, every hour up to 6 hours after acid instillation, and every 24 hours thereafter.

Bronchoalveolar Lavage (BAL)

After blood gases measurement at 72 hours post reperfusion, the rabbits were anesthetized with Nembutal 50 mg/kg i.v. and were euthanized by exsanguination. The abdominal aorta was transected to reduce red blood cell contamination of bronchial alveolar lavage fluid (BALF). The lung and heart were removed en bloc. The right lung was lavaged with an intratracheal tube (Hi-Lo-tracheal-tube, 3.0-mm) using 20-ml of HBSS and lidocain. Total and differential leukocyte counts of BALF were determined.

Gross Lung Weight

The whole lung from each rabbit was weighed immediately after harvest and was expressed as g/kg of body weight.

10 Peripheral Blood Count

Blood samples (0.05 ml for CBC, 0.2 ml for blood gases) were collected from the ear central artery catheter at baseline (pre-op), 2 hours, 4 hours, 6 hours, and 22 hours post reperfusion (prior to acid or saline instillation) and at 1 hour, 2 hours, 3 hours, 4 hours, 6 hours and every 24 hours after acid instillation. Hematology parameters were determined by Automated Hematology Analyzer according to the standard hematological procedures.

Pharmacokinetics

Blood samples (0.5 ml) were collected from the ear central artery catheter at baseline (pre-op), 4 hours, and 22 hours post reperfusion and at 1 hour, 4 hours, and every 24 hours after acid instillation.

Results and Discussion

20 In the rabbit model of ARDS, lung injury is manifested by hypoxemia (low PaO₂ - the pressure of O₂ in the arterial blood, as measured by a blood gas machine), lung edema (evidenced by an elevated lung weight to body weight ratio) and pro-inflammatory infiltrates into the alveolar space (evidenced by high white blood cell (WBC) and neutrophil (PMN) numbers). Although 40 kD branched PEG-6G4V11N35A Fab' did not protect rabbits from lung injury at any
25 of the doses tried (5 mg/kg and 20 mg/kg) (see Section (Y) above), the 20 kD linear PEG-6G4V11N35E Fab' had efficacy equal to, and, for some end-points, superior to that of the full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 and prevented lung injury in the rabbits as shown in Figs. 70A-70E. (The data points for 40 kD branched PEG-6G4V11N35A Fab' treated animals, full length 6G4.2.5 treated animals, and saline treated animals appearing in Figs. 70A-70E are taken from the data displayed in Figs. 67-69 and
30 generated in Example Y above.) In addition, these data indicate that large effective size anti-IL-8 Fab'-PEG conjugates can exhibit useful levels of efficacy in acute lung injury and ARDS.

AA. IN VIVO EFFICACY TESTING OF ANTI-IL-8 ANTIBODY REAGENTS IN RABBIT EAR MODEL OF TISSUE ISCHEMIA AND REPERFUSION

35 Full length murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5, 20 kD linear PEG-6G4V11N35E Fab', 30 kD linear PEG-6G4V11N35E Fab', and 40 kD branched PEG-6G4V11N35E Fab' were tested in a rabbit ear model of tissue ischemia and reperfusion injury.

Antibodies

A Fab'-SH antibody fragment of the affinity matured anti-IL-8 antibody 6G4V11N35E was expressed using the Fab' expression plasmid for 6G4V11N35E (described in Example T above) in *E. coli* grown to high

density in the fermentor as described by Carter et al., Bio/Technol, 10: 163-167 (1992). Anti-IL-8 6G4V11N35E Fab' variant was purified from fermentation paste and modified with 20 kD linear methoxy-PEG-maleimide, 30 kD linear methoxy-PEG-maleimide, or 40 kD branched methoxy-PEG-maleimide as described in Example T above. Pegylated material was formulated in phosphate buffered saline (PBS) at physiological pH.

Animals

1.0 to 1.5 kg New Zealand White rabbits were obtained from Western Oregon Rabbit Company.

Surgical procedure and animal evaluation

The procedure was essentially described by Vedder et al., Proc. Natl. Acad. Sci. (USA), 87: 2643-2646 (1990). Briefly, general anesthesia was achieved by intramuscular injections of Ketamine (50 mg/kg) plus Xylazine (5 mg/kg) and Acepromazine (2 mg/kg). The right external ear was prepared for surgery and under sterile procedure the ear was transected at its base, leaving intact only the central artery and vein. All nerves were transected to ensure that the ear was completely anesthetic. A straight microaneurysm clip (1.5x10mm) was placed across the artery to produce complete ischemia. The ear was reattached with the clip exiting through the wound. The rabbits were then housed at 26°C and 6 hours later the clip was removed to effect reperfusion. Untreated rabbits (n=11 animals) received an intravenous injection of vehicle (10 mM sodium acetate, 8% trehalose and 0.01% polysorbate-20 at pH 5.5) immediately prior to reperfusion. Treated animals received 5 mg/kg full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 (n=4 animals), 20 kD linear PEG-6G4V11N35E Fab' (n=3 animals), 30 kD linear PEG-6G4V11N35E Fab' (n=3 animals), or 40 kD branched PEG-6G4V11N35E Fab' (n=3 animals) immediately prior to reperfusion.

The ear volume and necrosis were measured daily by procedures described in Vedder et al., supra. Briefly, the ear was submerged in a beaker of water containing 1.2% Povidone iodine (Baxter) up to the intertragic incisure and the ear volume determined by the volume of fluid displaced. The ears were monitored in this manner for 7 days. The data are represented (in Fig. 71) as percent change in ear volume calculated as follows:

$$\% \text{ change in ear volume} = \frac{(\text{Ear vol. at day } x - \text{Ear vol. at day } 0) \times 100\%}{\text{Ear vol. at day } 0}$$

Animals were sacrificed at day 1 and day 7 for histological evaluation of the ear and the same section of ear was taken from all animals.

To determine that the therapeutic agents did not adversely affect any hematological parameter, aliquots of blood were withdrawn for complete blood counts and differentials immediately before reperfusion and at 24 hour intervals. In a separate experiment, blood samples were taken at 1, 5, 15, and 30 minutes and at 1 hour and 4 hours.

Results and Discussion

In the rabbit model of ear ischemia reperfusion injury, antibody was administered intravenously at a single dose (5 mg/kg) at the time of reperfusion. In this model, ischemia reperfusion injury is characterized by tissue damage, edema and sometimes necrosis; all attributable in part to neutrophil-mediated damage. Monitoring of ear volume over time is a surrogate end-point for evaluating edema in the ear tissue. The resulting data (depicted in Fig. 71) showed that treatment with 20 kD linear PEG-, 30 kD linear PEG- and 40

kD branched PEG-conjugated Fab's effectively reduced ear swelling and edema at all time points of observation (days 1, 3 and 5). In fact, the efficacy of all three PEGylated Fab's was statistically indistinguishable from that of the full length IgG murine anti-rabbit IL-8 monoclonal antibody 6G4.2.5 at all time points observed. These data support the efficacy of large effective size anti-IL-8 Fab'-PEG conjugates in ischemic reperfusion injury and specifically support the ability of 40 kD branched PEG-conjugated Fab' molecules to reach and act on disease effector targets in circulation and other tissues.

The following biological materials have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, USA (ATCC):

	<u>Material</u>	<u>ATCC Accession No.</u>	<u>Deposit Date</u>
10	hybridoma cell line 5.12.14	HB 11553	February 15, 1993
	hybridoma cell line 6G4.2.5	HB 11722	September 28, 1994
	pantiIL-8.2, E. coli strain 294 mm	97056	February 10, 1995
	p6G425chim2, E. coli strain 294 mm	97055	February 10, 1995
	p6G4V11N35A.F(ab') ₂	97890	February 20, 1997
15	E. coli strain 49D6(p6G4V11N35A.F(ab') ₂)	98332	February 20, 1997
	p6G425V11N35A.choSD	209552	December 16, 1997
	clone#1933 aIL8.92 NB 28605/12	CRL-12444	December 11, 1997
	clone#1934 aIL8.42 NB 28605/14	CRL-12445	December 11, 1997

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable deposit for 30 years from the date of deposit. These cell lines will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the cell lines to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the cell lines to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if the deposited cell lines should be lost or destroyed when cultivated under suitable conditions, they will be promptly replaced on notification with a specimen of the same cell line. Availability of the deposited cell lines is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

WE CLAIM:

1. A conjugate consisting essentially of one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein the apparent size of the conjugate is at least about 500 kD.
5
2. The conjugate of claim 1, wherein the apparent size of the conjugate is at least about 800 kD.
3. The conjugate of claim 1, wherein the apparent size of the conjugate is at least about 1,400
10 kD.
4. The conjugate of claim 1, wherein the apparent size of the conjugate is at least about 1,800 kD.
5. The conjugate of claim 1, wherein the apparent size of the conjugate is at least about 8 fold greater than the apparent size of at least one antibody fragment.
15
6. The conjugate of claim 5, wherein the apparent size of the conjugate is at least about 15 fold greater than the apparent size of at least one antibody fragment.
20
7. The conjugate of claim 6, wherein the apparent size of the conjugate is at least about 25 fold greater than the apparent size of at least one antibody fragment.
8. The conjugate of claim 1, wherein the conjugate contains no more than one antibody
25 fragment, and wherein the antibody fragment is selected from the group consisting of Fab, Fab', Fab'-SH, Fv, scFv and F(ab')₂.
9. The conjugate of claim 8 wherein the antibody fragment is F(ab')₂.
10. The conjugate of claim 1 wherein at least one antibody fragment is covalently attached to no more than about 10 nonproteinaceous polymer molecules.
30
11. The conjugate of claim 10 wherein the antibody fragment is covalently attached to no more than about 5 nonproteinaceous polymer molecules.
35
12. The conjugate of claim 11 wherein the antibody fragment is covalently attached to no more than about 2 nonproteinaceous polymer molecules.

13. The conjugate of claim 12 wherein the antibody fragment is attached to no more than 1 nonproteinaceous polymer molecule.

14. The conjugate of claim 12, wherein the antibody fragment comprises a heavy chain and a light chain derived from a parental antibody, wherein in the parental antibody the heavy and light chains are covalently linked by a disulfide bond between a cysteine residue in the light chain and a cysteine residue in the heavy chain, wherein in the antibody fragment the cysteine residue in the light or heavy chain is substituted with another amino acid and the cysteine residue in the opposite chain is covalently linked to a nonproteinaceous polymer molecule.

10

15. The conjugate of claim 8 wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH.

16. The conjugate of claim 15 wherein the antibody fragment is covalently attached to no more than 1 nonproteinaceous polymer molecule.

15

17. The conjugate of claim 16 wherein the nonproteinaceous polymer molecule in the conjugate is covalently attached to the hinge region of the antibody fragment.

20

18. The conjugate of claim 1 wherein at least one nonproteinaceous polymer is a polyethylene glycol (PEG).

19. The conjugate of claim 18 wherein the PEG has an average molecular weight of at least about 20 kD.

25

20. The conjugate of claim 19 wherein the PEG has an average molecular weight of at least about 40 kD.

30

21. The conjugate of claim 19 wherein the PEG is a single chain molecule.

22. The conjugate of claim 20 wherein the PEG is a branched chain molecule.

35

23. The conjugate of claim 19, wherein the conjugate contains no more than one antibody fragment, and wherein the antibody fragment is a F(ab')₂ and is covalently attached to no more than about 2 PEG molecules.

24. The conjugate of claim 19, wherein the conjugate contains no more than one antibody fragment, and wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH and is covalently attached to no more than one PEG molecule.

5 25. The conjugate of claim 24 wherein the PEG molecule is covalently attached to the hinge region of the antibody fragment.

26. The conjugate of claim 1 wherein at least one antibody fragment comprises an antigen binding site that binds to human interleukin-8 (IL-8).

10 27. The conjugate of claim 26, wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a polyethylene glycol having an average molecular weight of at least about 30 kD.

15 28. The conjugate of claim 26 wherein the antibody fragment comprising the anti-human IL-8 antigen binding site is humanized.

20 29. The conjugate of claim 28 wherein the anti-human IL-8 antigen binding site comprises the complementarity determining regions of a light chain polypeptide amino acid sequence selected from the group consisting of the 6G4V11N35A light chain polypeptide amino acid sequence of Fig. 36 (SEQ ID NO:56) and the 6G4V11N35E light chain polypeptide amino acid sequence of Fig. 45 (SEQ ID NO:62).

25 30. The conjugate of claim 1 wherein the conjugate contains no more than one antibody fragment.

31. A composition comprising the conjugate of claim 1 and a carrier.

30 32. The composition of claim 31 that is sterile.

33. The conjugate of claim 1, wherein the covalent structure of the conjugate is free of any matter other than the antibody fragment and nonproteinaceous polymer molecules that form the conjugate.

35 34. The conjugate of claim 1, wherein the covalent structure of the conjugate incorporates one or more nonproteinaceous labels, and wherein the covalent structure of the conjugate is free of any matter other than the antibody fragment, nonproteinaceous polymer and nonproteinaceous label molecules that form the conjugate.

35. The conjugate of claim 34 wherein at least one nonproteinaceous label is a radiolabel.

36. A method of treating an inflammatory disorder in a mammal comprising administering to the mammal an effective amount of a conjugate consisting essentially of one or more antibody fragments covalently attached to one or more nonproteinaceous polymer molecules, wherein at least one antibody fragment comprises an antigen binding site that binds to human interleukin-8 (IL-8), and wherein the apparent size of the conjugate is at least about 500 kD.

37. The method of claim 36 wherein the inflammatory disorder is acute lung injury.

10

38. The method of claim 37 wherein the acute lung injury includes adult respiratory distress syndrome (ARDS).

39. The method of claim 36 wherein the inflammatory disorder is an ischemic reperfusion disorder.

15

40. The method of claim 36, wherein the inflammatory disorder is hypovolemic shock.

41. The method of claim 39, wherein the ischemic reperfusion disorder is a surgical tissue reperfusion injury.

20

42. The method of claim 39, wherein the ischemic reperfusion disorder is myocardial ischemia.

43. The method of claim 39, wherein the ischemic reperfusion disorder is acute myocardial infarction.

25

44. The method of claim 36, wherein the inflammatory disorder is inflammatory bowel disease.

45. The method of claim 44, wherein the inflammatory bowel disease is ulcerative colitis.

30

46. The method of claim 36, wherein the inflammatory disorder is bacterial pneumonia.

47. The method of claim 36, wherein the apparent size of the conjugate is at least about 800 kD.

48. The method of claim 47, wherein the apparent size of the conjugate is at least about 1,400 kD.

35

49. The method of claim 37, wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a polyethylene glycol (PEG) having an average molecular weight of at least about 20 kD.

50. The method of claim 49, wherein the PEG is a single chain molecule.

51. The method of claim 39, wherein the conjugate contains no more than one antibody fragment, wherein the antibody fragment is selected from the group consisting of Fab, Fab' and Fab'-SH, wherein the antibody fragment is covalently attached to no more than one nonproteinaceous polymer molecule, and wherein the nonproteinaceous polymer molecule is a polyethylene glycol (PEG) having a molecular an average weight of at least about 20 kD.

52. The method of claim 51, wherein the PEG has an average molecular weight of at least about 30 kD.

53. The method of claim 52, wherein the PEG has an average molecular weight of at least about 40 kD.

54. The method of claim 51, wherein the PEG is a single chain molecule.

55. The method of claim 53, wherein the PEG is a branched chain molecule.

56. The method of claim 36, wherein at least one nonproteinaceous polymer molecule is a polyethylene glycol (PEG) having an average molecular weight of at least about 20 kD.

57. The method of claim 56, wherein the PEG has an average molecular weight of at least about 30 kD.

58. The method of claim 57, wherein the PEG has an average molecular weight of at least about 40 kD.

59. The method of claim 56, wherein the PEG is a single chain molecule.

60. The method of claim 58, wherein the PEG is a branched chain molecule.


61. The method of claim 36, wherein the antibody fragment comprising the anti-human IL-8 antigen binding site is monoclonal and humanized.



62. The method of claim 61, wherein the anti-human IL-8 antigen binding site comprises the complementarity determining regions of a light chain polypeptide amino acid sequence selected from the group consisting of the 6G4V11N35A light chain polypeptide amino acid sequence of Fig. 36 (SEQ ID NO:56) and the 6G4V11N35E light chain polypeptide amino acid sequence of Fig. 45 (SEQ ID NO:62).



63. The method of claim 36, wherein the covalent structure of the conjugate is free of any matter other than the antibody fragment and nonproteinaceous polymer molecules that form the conjugate.

10 64. The method of claim 36 wherein the mammal is human.

FIG. 1










FIG. 6




FIG. 7

FIG. 8




FIG. 9





FIG. 10






FIG. 11B




FIG. 11A




FIG. 11D




FIG. 11C




FIG. 11F




FIG. 11E




FIG. 11H




FIG. 11G




FIG. 11J





FIG. 11I

FIG. 12



Light Chain Primers:

MKLC-1, 22mer

FIG. 13

5' CAGTCCAACTGTTCAGGACGCC 3'

(SEQ ID NO: 1)

MKLC-2, 22mer

5' GTGCTGCTCATGCTGTAGGTGC 3'

(SEQ ID NO: 2)

MKLC-3, 23mer

5' GAAGTTGATGTCTTGTGAGTGGC

3' (SEQ ID NO: 3)

Heavy Chain Primers:

IGG2AC-1, 24mer

5' GCATCCTAGAGTCACCGAGGAGCC

3' (SEQ ID NO: 4)

IGG2AC-2, 22mer

5' CACTGGCTCAGGGAAATAACCC 3'

(SEQ ID NO: 5)

IGG2AC-3, 22mer

5' GGAGAGCTGGGAAGGTGTGCAC 3'

(SEQ ID NO: 6)

FIG. 14

Light chain forward primer

SL001A-2 35 mer

5' ACAAACGCGTACGCT GACATCGTCATGACCCAGTC 3'
 T T T
 A

(SEQ ID NO: 7)
(SEQ ID NO: 8)
(SEQ ID NO: 9)

Light chain reverse primer

SL001B 37 mer

5' GCTCTTCGAATG GTGGGAAGATGCATACAGTTGGTGC 3' (SEQ ID NO: 10)

Heavy chain forward primer

FIG. 15

SL002B 39 mer

5' CGATGGGCCCCG ATAGACCGATGGGGCTGTTGTTTGGC 3' (SEQ ID NO: 11)
 T C (SEQ ID NO: 12)
 G (SEQ ID NO: 13)
 A (SEQ ID NO: 14)

Heavy chain reverse primer

SL002B 39-MER

5' CGATGGGCCCCG ATAGACCGATGGGGCTGTTGTTTGGC 3' (SEQ ID NO: 11)
 T (SEQ ID NO: 15)
 A (SEQ ID NO: 14)
 G (SEQ ID NO: 13)

FIG. 16

1 TTCTATTGCT ACAAACGCGT ACGCTGAGGT GCAGCTGGTG GAGTCTGGGG GAGGCTTAGT
 AAGATAACGA TGTTTGCGCA TGC GACTCCA CGTCGACCAC CTCAGACCCC CTCCGAATCA
 1 E V Q L V E S G G G L V

61 GCCGCCTGGA GGGTCCCTGA AACTCTCCTG TGCAGCCTCT GGATTCATAT TCAGTAGTTA
 CGGCGGACCT CCCAGGGACT TTGAGAGGAC ACGTCGGAGA CCTAAGTATA AGTCATCAAT
 13 P P G G S L K L S C A A S G F I F S S Y * *

CDR #1

121 TGGCATGTCT TGGGTTCGCC AGACTCCAGG CAAGAGCCTG GAGTTGGTCTG CAACCATTAA
 ACCGTACAGA ACCCAAGCGG TCTGAGGTCC GTTCTCGGAC CTCAACCAGC GTTGTAATT
 33 G M S W V R Q T P G K S L E L V A T I N * * *

181 TAATAATGGT GATAGCACCT ATTATCCAGA CAGTGTGAAG GGCCGATTCA CCATCTCCCG
 ATTATTACCA CTATCGTGGA TAATAGGTCT GTCACACTTC CCGGCTAAGT GGTAGAGGGC
 53 N N G D S T Y Y P D S V K G R F T I S R * * * * *

CDR #2

241 AGACAATGCC AAGAACACCC TGTACCTGCA AATGAGCAGT CTGAAGTCTG AGGACACAGC
 TCTGTTACGG TTCTTGTTGGG ACATGGACGT TTACTCGTCA GACTTCAGAC TCCTGTGTCG
 73 D N A K N T L Y L Q M S S L K S E D T A

301 CATGTTTTAC TGTGCAAGAG CCCTCATTAG TTCGGCTACT TGGTTTGGTT ACTGGGGCCA
 GTACAAAATG ACACGTTCTC GGGAGTAATC AAGCCGATGA ACCAAACCAA TGACCCCGGT
 93 M F Y C A R A L I S S A T W F G Y W G Q * * * * *

CDR #3

361 AGGGACTCTG GTCAGTGTCT CTGCAGCCAA AACAACAGCC CCATCTGTCT
 TCCCTGAGAC CAGTGACAGA GACGTCGGTT TTGTTGTCGG GGTAGACAGA
 113 G T L V T V S A A K T T A P S V Y

411 ^{Apal} ATCCGGG (SEQ ID NO: 18)
 TAGGCCC
 130 P (SEQ ID NO: 19)

FIG. 17

FIG. 18

VL.front 31-MER

5' ACAAACGCGTACGCTGATATCGTCATGACAG 3' (SEQ ID NO: 20)

VL.rear 31-MER

5' GCAGCATCAGCTCTTCGAAGCTCCAGCTTGG 3' (SEQ ID NO: 21)

VH.front.SPE 21-MER

5' CCACTAGTACGCAAGTTCACG 3' (SEQ ID NO: 22)

VH.rear 33-MER

5' GATGGGCCCTTGGTGGAGGCTGCAGAGACAGTG 3' (SEQ ID NO: 23)

1 ATGAAGAAGA ATATCGCATT TCTTCTTGCA TCTATGTTCTG TTTTCTCTAT TGCTACAAAC
TACTTCTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTTG
-23 M K K N I A F L L A S M F V F S I A T N

61 GCGTACGCTG ATATCGTCAT GACACAGTCT CAAAAATTCA TGTCCACATC AGTAGGAGAC
CGCATGCGAC TATAGCAGTA CTGTGTCAGA GTTTTAAAGT ACAGGTGTAG TCATCCTCTG
-3 A Y A D I V M T Q S Q K F M S T S V G D

121 AGGGTCAGCG TCACCTGCAA GGCCAGTCAG AATGTGGGTA CTAATGTAGC CTGGTATCAA
TCCCAGTCGC AGTGGACGTT CCGGTCAGTC TTACACCCAT GATTACATCG GACCATAGTT
18 R V S V T C K A S O N V G T N V A W Y Q

CDR #1

181 CAGAAACCAG GGCAATCTCC TAAAGCACTG ATTTACTCGT CATCCTACCG GTACAGTGGA
GTCTTTGGTC CCGTTAGAGG ATTTCGTGAC TAAATGAGCA GTAGGATGGC CATGTCACCT
38 Q K P G Q S P K A L I Y S S S Y R Y S G

CDR #2

241 GTCCCTGATC GCTTCACAGG CAGTGGATCT GGGACAGATT TCACTCTCAC CATCAGCCAT
CAGGGACTAG CGAAGTGTCC GTCACCTAGA CCCTGTCTAA AGTGAGAGTG GTAGTCGGTA
58 V P D R F T G S G S G T D F T L T I S H

301 GTGCAGTCTG AAGACTTGGC AGACTATTTT TGTCAGCAAT ATAACATCTA TCCTCTCAGC
CACGTCAGAC TTCTGAACCG TCTGATAAAG ACAGTCGTTA TATTGTAGAT AGGAGAGTGC
78 V Q S E D L A D Y F C Q Q Y N I Y P L T

CDR #3

BstBI

361 TTCGGTCCTG GGACCAAGCT GGAGCTTCGA AGAGCTGTGG CTGCACCATC TGTCTTCATC
AAGCCAGGAC CCTGGTTTGA CCTCGAAGCT TCTCGACACC GACGTGGTAG ACAGAAGTAG
98 F G P G T K L E L R R A V A A P S V F I

421 TTCCCGCCAT CTGATGAGCA GTTGAAATCT GGAAGTGTCT CTGTTGTGTG CCTGCTGAAT
AAGGGCGGTA GACTACTCGT CAACTTTAGA CCTTGACGAA GACAACACAC GGACGACTTA
118 F P P S D E Q L K S G T A S V V C L L N

481 AACTTCTATC CCAGAGAGGC CAAAGTACAG TGGAAGGTGG ATAACGCCCT CCAATCGGGT
TTGAAGATAG GGTCTCTCCG GTTTCATGTC ACCTTCCACC TATTGCGGGA GGTTAGCCCA
138 N F Y P R E A K V Q W K V D N A L Q S G

541 AACTCCCAGG AGAGTGTCAC AGAGCAGGAC AGCAAGGACA GCACCTACAG CCTCAGCAGC
TTGAGGGTCC TCTCACAGTG TCTCGTCCTG TCGTTCCTGT CGTGGATGTC GGAGTCGTCG
158 N S Q E S V T E Q D S K D S T Y S L S S

601 ACCCTGACGC TGAGCAAAGC AGACTACGAG AAACACAAAG TCTACGCCTG CGAAGTCACC
TGGGACTGCG ACTCGTTTTC TCTGATGCTC TTTGTGTTTC AGATGCGGAC GCTTCAGTGG
178 T L T L S K A D Y E K H K V Y A C E V T

661 CATCAGGGCC TGAGCTCGCC CGTCACAAAG AGCTTCAACA GGGGAGAGTG
GTAGTCCCGG ACTCGAGCGG GCAGTGTTTC TCGAAGTTGT CCCCTCTCAC
198 H Q G L S S P V T K S F N R G E C (SEQ ID NO: 25)

711 TTAA (SEQ ID NO: 24)
AATT

216 O

FIG. 19

1 ATGAAAAGA ATATCGCATT TCTTCTTGCA TCTATGTTCTG TTTTCTCTAT TGCTACAAAC
TACTTTTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTG
-23 M K K N I A F L L A S M F V F S I A T N

61 GCGTACGCTG AGGTGCAGCT GGTGGAGTCT GGGGGAGGCT TAGTGCCGCC TGGAGGGTCC
CGCATGCGAC TCCACGTCGA CCACCTCAGA CCCCTCCGA ATCACGCGG ACCTCCAGG
-3 A Y A E V Q L V E S G G G L V P P G G S

121 CTGAAACTCT CCTGTGCAGC CTCTGGATTC ATATTCAGTA GTTATGGCAT GTCTTGGGTT
GACTTTGAGA GGACACGTCG GAGACCTAAG TATAAGTCAT CAATACCGTA CAGAACCCAA
18 L K L S C A A S G F I F S S Y G M S W V
* * * * *

CDR #1

181 CGCCAGACTC CAGGCAAGAG CCTGGAGTTG GTCGCAACCA TTAATAATAA TGGTGATAGC
GCGGTCTGAG GTCCGTTCTC GGACCTCAAC CAGCGTTGGT AATTATTATT ACCACTATCG
38 R Q T P G K S L E L V A T I N N N G D S
* * * * *

241 ACCTATTATC CAGACAGTGT GAAGGGCCGA TTCACCATCT CCCGAGACAA TGCCAAGAAC
TGGATAATAG GTCTGTCACA CTCCCCGGCT AAGTGGTAGA GGGCTCTGTT ACGGTTCTTG
58 T Y Y P D S V K G R F T I S R D N A K N
* * * * *

CDR #2

301 ACCCTGTACC TGCAAATGAG CAGTCTGAAG TCTGAGGACA CAGCCATGTT TTACTGTGCA
TGGGACATGG ACGTTTACTC ~~GTCAGACTTC~~ ~~AGACTCCTGT~~ GTCGGTACAA AATGACACGT
78 T L Y L Q M S S L K S E D T A M F Y C A

361 AGAGCCCTCA TTAGTTCGGC TACTTGGTTT GGTACTGGG GCCAAGGGAC TCTGGTCACT
TCTCGGGAGT AATCAAGCCG ATGAACCAA CCAATGACCC CGGTTCCCTG AGACCAGTGA
98 R A L I S S A T W F G Y W G Q G T L V T
* * * * *

CDR #3

ApaI

421 GTCTCTGCAG CCTCCACCAA GGGCCCATCG GTCTTCCCCC TGGCACCCCTC CTCCAAGAGC
CAGAGACGTC GGAGGTGGTT CCCGGGTAGC CAGAAGGGGG ACCGTGGGAG GAGGTTCTCG
118 V S A A S T K G P S V F P L A P S S K S

481 ACCTCTGGGG GCACAGCGGC CCTGGGCTGC CTGGTCAAGG ACTACTTCCC CGAACCGGTG
TGGAGACCCC CGTGTGCGCG GGACCCGACG GACCAGTTCC TGATGAAGGG GCTTGGCCAC
138 T S G G T A A L G C L V K D Y F P E P V

541 ACGGTGTCGT GGAAGTCAGG CGCCCTGACC AGCGGCGTGC ACACCTTCCC GGCTGTCCTA
TGCCACAGCA CCTTGAGTCC GCGGGACTGG TCGCCGCACG TGTGGAAGGG CCGACAGGAT
158 T V S W N S G A L T S G V H T F P A V L

601 CAGTCCTCAG GACTCTACTC CCTCAGCAGC GTGGTGACCG TGCCCTCCAG CAGCTTGGGC
GTCAGGAGTC CTGAGATGAG GGAGTCGTCG CACCACTGGC ACGGGAGGTC GTCGAACCCG
178 Q S S G L Y S L S S V V T V P S S S L G

FIG. 20A

661 ACCCAGACCT ACATCTGCAA CGTGAATCAC AAGCCCAGCA ACACCAAGGT GGACAAGAAA
TGGGTCTGGA TGTAGACGTT GCACTTAGTG TTCGGGTCGT TGTGGTTCCA CCTGTTCTTT
198 T Q T Y I C N V N H K P S N T K V D K K
721 GTTGAGCCCA AATCTTGTGA CAAAACTCAC ACATGA (SEQ ID NO: 26)
CAACTCGGGT TTAGAACACT GTTTGAGTG TGTACT
218 V E P K S C D K T H T O (SEQ ID NO: 27)

FIG. 20B

Light Chain Primers:

MKLC-1, 22mer

5' CAGTCCAACCTGTTTCAGGACGCC 3'

(SEQ ID NO: 1)

MKLC-2, 22mer

5' GTGCTGCTCATGCTGTAGGTGC 3'

(SEQ ID NO: 2)

MKLC-3, 23mer

5' GAAGTTGATGTCTTGTGAGTGGC

3' (SEQ ID NO: 3)

Heavy Chain Primers:

IGG2AC-1, 24mer

5' GCATCCTAGAGTCACCGAGGAGCC

3' (SEQ ID NO: 4)

IGG2AC-2, 22mer

5' CACTGGCTCAGGGAAATAACCC 3'

(SEQ ID NO: 5)

IGG2AC-3, 22mer

5' GGAGAGCTGGGAAGGTGTGCAC 3'

(SEQ ID NO: 6)

FIG. 21

Light chain forward primer

6G4.light.Nsi 36-MER

5' CCAATGCATACGCT GAC ATC GTG ATG ACC CAG ACC CC 3' (SEQ ID NO: 28)
T T T A A (SEQ ID NO: 29)
A A (SEQ ID NO: 30)

Light chain reverse primer

6G4.light.Mun 35-MER

5' AGA TGT CAA TTG CTC ACT GGA TGG TGG GAA GAT GG 3' (SEQ ID NO: 31)

FIG. 22

Heavy chain forward primer

6G4.heavy.Mlu 32-MER

5' CAAACGCGTACGCT GAG ATC CAG CTG CAG CAG 3' (SEQ ID NO: 32)
T C (SEQ ID NO: 33)

Heavy chain reverse primer

SL002B 39-MER

5' CGATGGGCCCCG ATAGACCGATGGGGCTGTTGTTTGGC 3' (SEQ ID NO: 11)
T (SEQ ID NO: 15)
A (SEQ ID NO: 14)
G (SEQ ID NO: 13)

FIG. 23

70 G ATATCGTGAT GACACAGACA CCACTCTCCC TGCCTGTCAG TCTTGGAGAT
 C TATAGCACTA CTGTGTCTGT GGTGAGAGGG ACGGACAGTC AGAACCTCTA
 1 D I V M T Q T P L S L P V S L G D

121 CAGGCCTCCA TCTCTTGCAG ATCTAGTCAG AGCCTTGTAC ACGGTATTGG AAACACCTAT
 GTCCGGAGGT AGAGAACGTC TAGATCAGTC TCGGAACATG TGCCATAACC TTTGTGGATA
 18 Q A S I S C R S S O S L V H G I G N T Y
 * * * * *

CDR #1

181 TTACATTGGT ACCTGCAGAA GCCAGGCCAG TCTCCAAAGC TCCTGATCTA CAAAGTTTCC
 AATGTAACCA TGGACGTCTT CGGTCCGGTC AGAGGTTTCG AGGACTAGAT GTTTCAAAGG
 38 L H W Y L Q K P G Q S P K L L I Y K V S
 * * * * *

CDR #2

241 AACCGATTTT CTGGGGTCCC AGACAGGTTC AGTGGCAGTG GATCAGGGAC AGATTTTACA
 TTGGCTAAAA GACCCAGGG TCTGTCCAAG TCACCGTCAC CTAGTCCCTG TCTAAAGTGT
 58 N R F S G V P D R F S G S G S G T D F T
 * * * * *

301 CTCAGGATCA GCAGAGTGGA GGCTGAGGAT CTGGGACTTT ATTTCTGCTC TCAAAGTACA
 GAGTCCTAGT CGTCTCACCT CCGACTCCTA GACCCTGAAA TAAAGACGAG AGTTTCATGT
 78 L R I S R V E A E D L G L Y F C S Q S T
 * * * * *

CDR #3

361 CATGTTCCGC TCACGTTCGG TGCTGGGACC AAGCTGGAGC TGAAACGGGC TGATGCTGCA
 GTACAAGGCG AGTGCAAGCC ACGACCCTGG TTCGACCTCG ACTTTGCCCCG ACTACGACGT
 98 H V P L T F G A G T K L E L K R A D A A
 * * * * *

MunI

421 CCAACTGTAT CCATCTTCCC ACCATCCAGT GAGCAATTGA (SEQ ID NO: 34)
 GGTTGACATA GGTAGAAGGG TGGTAGGTCA CTCGTAACT
 118 P T V S I F P P S S E Q L K (SEQ ID NO: 35)

FIG. 24

70 G AGATTCAGCT GCAGCAGTCT GGACCTGAGC TGATGAAGCC TGGGGCTTCA
 C TCTAAGTCGA CGTCGTCAGA CCTGGACTCG ACTACTTCGG ACCCCGAAGT
 1 E I Q L Q Q S G P E L M K P G A S

121 GTGAAGATAT CCTGCAAGGC TTCTGGTTAT TCATTAGTA GCCACTACAT GCACTGGGTG
 CACTTCTATA GGACGTTCCG AAGACCAATA AGTAAGTCAT CCGTGATGTA CGTGACCCAC
 18 V K I S C K A S G Y S F S S H Y M H W V
 * * * * *

CDR #1

181 AAGCAGAGCC ATGGAAAGAG CCTTGAGTGG ATTGGCTACA TTGATCCTTC CAATGGTGAA
 TTCGTCTCGG TACCTTTCTC GGAATCACC TAACCGATGT AACTAGGAAG GTTACCACTT
 38 K Q S H G K S L E W I G Y I D P S N G E
 * * * * *

CDR #2

241 ACTACTTACA ACCAGAAATT CAAGGGCAAG GCCACATTGA CTGTAGACAC ATCTTCCAGC
 TGATGAATGT TGGTCTTTAA GTTCCCGTTC CCGTGTAAT GACATCTGTG TAGAAGGTG
 58 T T Y N Q K F K G K A T L T V D T S S S
 * * * * *

301 ACAGCCAACG TGCATCTCAG CAGCCTGACA TCTGATGACT CTGCAGTCTA TTTCTGTGCA
 TGTCGGTTGC ACGTAGAGTC GTCGGACTGT AGACTACTGA GACGTCAGAT AAAGACACGT
 78 T A N V H L S S L T S D D S A V Y F C A

361 AGAGGGGACT ATAGATACAA CGGCGACTGG TTTTTCGATG TCTGGGGCGC AGGGACCACG
 TCTCCCCTGA TATCTATGTT GCCGCTGACC AAAAAGCTAC AGACCCCGCG TCCCTGGTGC
 98 R G D Y R Y N G D W F F D V W G A G T T
 * * * * *

CDR #3

BstEII

421 GTCACCGTCT CCTCCGCCAA AACCGACAGC CCCATCGGTC TATCCGGGGC
 CAGTGGCAGA GGAGGCGGAT TTGGCTGTGCG GGGTAGCCAG ATAGGCCCCG
 118 V T V S S A K T D S P I G L S G P

ApaI

471 CATC (SEQ ID NO: 36)
 GTAG
 135 I (SEQ ID NO: 37)

FIG. 25

5' CTTGGTGGAGGCGGAGGAGACG 3' (SEQ ID NO: 38)

Mutagenesis Primer for 6G425VL

DS/VF 38MER

5' GAAACGGGCTGTTGCTGCACCAACTGTATTCATCTTCC 3' (SEQ ID NO: 39)

SYN.BstEII 31 MER

5' GTCACCGTCT CCTCCGCCTC CACCAAGGGC C 3' (SEQ ID NO: 40)

SYN.Apa 22 MER

5' CTTGGTGGAGGCGGAGGAGACG 3' (SEQ ID NO: 38)

FIG. 26

1 ATGAAGAAGA ATATCGCATT TCTTCTTGCA TCTATGTTTCG TTTTCTCTAT TGCTACAAAT
TACTTCTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTTA
-23 M K K N I A F L L A S M F V F S I A T N

61 GCATACGCTG ATATCGTGAT GACACAGACA CCACTCTCCC TGCCTGTCAG TCTTGGAGAT
CGTATGCGAC TATAGCACTA CTGTGTCTGT GGTGAGAGGG ACGGACAGTC AGAACCTCTA
-3 A Y A D I V M T Q T P L S L P V S L G D

121 CAGGCCTCCA TCTCTTGCGAG ATCTAGTCAG AGCCTTGTTAC ACGGTATTGG AAACACCTAT
GTCCGGAGGT AGAGAACGTC TAGATCAGTC TCGGAACATG TGCCATAACC TTTGTGGATA
18 Q A S I S C R S S O S L V H G I G N T Y
* * * * *

CDR #1

181 TTACATTGGT ACCTGCAGAA GCCAGGCCAG TCTCCAAAGC TCCTGATCTA CAAAGTTTCC
AATGTAACCA TGGACGTCTT CGGTCCGGTC AGAGGTTTCG AGGACTAGAT GTTTCAAAGG
38 L H W Y L Q K P G Q S P K L L I Y K V S
* * * * *

CDR #2

241 AACCGATTTT CTGGGGTCCC AGACAGGTTT AGTGGCAGTG GATCAGGGAC AGATTTTACA
TTGGCTAAAA GACCCAGGG TCTGTCCAAG TCACCGTCAC CTAGTCCCTG TCTAAAGTGT
58 N R F S G V P D R F S G S G S G T D F T
* * * * *

301 CTCAGGATCA GCAGAGTGGA GGCTGAGGAT CTGGGACTTT ATTTCTGCTC TCAAAGTACA
GAGTCCCTAGT CGTCTCACCT CCGACTCCTA GACCCTGAAA TAAAGACGAG AGTTTCATGT
78 L R I S R V E A E D L G L Y F C S Q S T
* * * * *

CDR #3

361 CATGTTCCGC TCACGTTCCG TGCTGGGACC AAGCTGGAGC TGAAACGGGC TGTTGCTGCA
GTACAAGGCG AGTGCAAGCC ACGACCCTGG TTCGACCTCG ACTTTGCCCG ACAACGACGT
98 H V P L T F G A G T K L E L K R A V A A
* * * * *

421 CCAACTGTAT TCATCTTCCC ACCATCCAGT GAGCAATTGA AATCTGGAAC TGCCTCTGTT
GGTTGACATA AGTAGAAGGG TGGTAGGTCA CTCGTAACT TTAGACCTTG ACGGAGACAA
118 P T V F I F P P S S E Q L K S G T A S V

481 GTGTGCCTGC TGAATAACTT CTATCCCAGA GAGGCCAAAG TACAGTGGAA GGTGGATAAC
CACACGGACG ACTTATTGAA GATAGGGTCT CTCCGGTTTC ATGTCACCTT CCACCTATTG
138 V C L L N N F Y P R E A K V Q W K V D N

541 GCCCTCCAAT CGGGTAACTC CCAGGAGAGT GTCACAGAGC AGGACAGCAA GGACAGCACC
CGGGAGGTGA GCCCATTGAG GGTCCTCTCA CAGTGTCTCG TCCTGTCGTT CCTGTCTGTT
158 A L Q S G N S Q E S V T E Q D S K D S T

601 TACAGCCTCA GCAGCACCCT GACGCTGAGC AAAGCAGACT ACGAGAAACA CAAAGTCTAC
ATGTCGGAGT CGTCGTGGGA CTGCGACTCG TTTCGTCTGA TGCTCTTTGT GTTTCAGATG
178 Y S L S S T L T L S K A D Y E K H K V Y

FIG. 27A

661 GCCTGCGAAG TCACCCATCA GGGCTGAGC TCGCCCGTCA CAAAGAGCTT CAACAGGGGA
CGGACGCTTC AGTGGTAGT CCGGACTCG AGCGGGCAGT GTTCTCGAA GTTGTCCTT
198 A C E V T H Q G L S S P V T K S F N R G

721 GAGTGTTAA (SEQ ID NO: 41)
CTCACAATT
218 E C O (SEQ ID NO: 42)

FIG. 27B

1 ATGAAAAAGA ATATCGCATT TCTTCTTGCA TCTATGTTCTG TTTTCTCTAT TGCTACAAAC
TACTTTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTTG
-23 M K K N I A F L L A S M F V F S I A T N

61 GCGTACGCTG AGATTCAAGCT GCAGCAGTCT GGACCTGAGC TGATGAAGCC TGGGGCTTCA
CGCATGCGAC TCTAAGTCGA CGTCGTCAGA CCTGGACTCG ACTACTTCGG ACCCCGAAGT
-3 A Y A E I Q L Q Q S G P E L M K P G A S

121 GTGAAGATAT CCTGCAAGGC TTCTGGTTAT TCATTCAAGTA GCCACTACAT GCACTGGGTG
CACTTCTATA GGACGTTCCG AAGACCAATA AGTAAGTCAT CGGTGATGTA CGTGACCCAC
18 V K I S C K A S G Y S F S S H Y M H W V
* * * *

CDR #1

181 AAGCAGAGCC ATGGAAAGAG CCTTGAGTGG ATTGGCTACA TTGATCCTTC CAATGGTGAA
TTCGTCTCGG TACCTTTCTC GGAACCTACC TAACCGATGT AACTAGGAAG GTTACCACTT
38 K Q S H G K S L E W I G Y I D P S N G E
* * * * *

CDR #2

241 ACTACTTACA ACCAGAAATT CAAGGGCAAG GCCACATTGA CTGTAGACAC ATCTTCCAGC
TGATGAATGT TGGTCTTTAA GTTCCCGTTC CGGTGTAAT GACATCTGTG TAGAAGGTGC
58 T T Y N Q K F K G K A T L T V D T S S S
* * * * *

301 ACAGCCAACG TGCATCTCAG CAGCCTGACA TCTGATGACT CTGCAGTCTA TTTCTGTGCA
TGTCGGTTGC ACGTAGAGTC GTCGGACTGT AGACTACTGA GAGGTGAGAT AAAGACACGT
78 T A N V H L S S L T S D D S A V Y F C A

361 AGAGGGGACT ATAGATACAA CGGCGACTGG TTTTTCGATG TCTGGGGCGC AGGGACCACG
TCTCCCCTGA TATCTATGTT GCCGCTGACC AAAAAGCTAC AGACCCCGCG TCCCTGGTGC
98 R G D Y R Y N G D W F F D V W G A G T T
* * * * *

CDR #3

421 GTCACCGTCT CCTCCGCCTC CACCAAGGGC CCATCGGTCT TCCCCCTGGC ACCCTCCTCC
CAGTGGCAGA GGAGGCGGAG GTGGTTCCCG GGTAGCCAGA AGGGGGACCG TGGGAGGAGG
118 V T V S S A S T K G P S V F P L A P S S

481 AAGAGCACCT CTGGGGGCAC AGCGGCCCTG GGCTGCCTGG TCAAGGACTA CTTCCCCGAA
TTCTCGTGGA GACCCCGTG TCGCCGGGAC CCGACGGACC AGTTCCTGAT GAAGGGGCTT
138 K S T S G G T A A L G C L V K D Y F P E

541 CCGGTGACCG TGTCGTGGAA CTCAGGCGCC CTGACCAGCG GCGTGCACAC CTTCCCGGCT
GGCCACTGCC ACAGCACCTT GAGTCCGCGG GACTGGTTCG CGCACGTGTG GAAGGGCCGA
158 P V T V S W N S G A L T S G V H T F P A

601 GTCCTACAGT CCTCAGGACT CTACTCCCTC AGCAGCGTGG TGACCGTGCC CTCCAGCAGC
CAGGATGTCA GGAGTCCTGA GATGAGGGAG TCGTCGCACC ACTGGCACGG GAGGTCTGCG
178 V L Q S S G L Y S L S S V V T V P S S S

FIG. 28A

661 TTGGGCACCC AGACCTACAT CTGCAACGTG AATCACAAGC CCAGCAACAC CAAGGTGGAC
AACCCGTGGG TCTGGATGTA GACGTTGCAC TTAGTGTTCTG GGTCTGTTGTG GTTCCACCTG
198 L G T Q T Y I C N V N H K P S N T K V D

721 AAGAAAGTTG AGCCCAATC TTGTGACAAA ACTCACACAT GA (SEQ ID NO: 43)
TTCCTTCAAC TCGGGTTTAG AACACTGTCTT TGAGTGTGTA CT
218 K K V E P K S C D K T H T O (SEQ ID NO: 44)

FIG. 28B

Variable Light Chain Domain

[illegible]

```

50          60          70          80          90          100
6G425      YKVSNRFGVTD RFS DSGSGTDFTLRISRVEAEDLGLYFCSQSTH VPLTFGAGTKLELYR (SEQ ID NO: 45)
          *      *      *      *      *      *      *      *
F(ab)-1    YKVSNRFGVPSRFS GSGSGTDFTLT ISSLPQEDFATYYCSQSTH VPLTFGQGTKVEIKR (SEQ ID NO: 46)
          *      *      *      *      *      *      *      *
humk1      YSGSTLES GVP SRFS GSGSGTDFTLT ISSLPQEDFATYYCQ HNEYPLTFGQGTKVEIKR (SEQ ID NO: 47)
          ***      *****
          +++++++      +++++++
          L2          L3

```

Variable Heavy Chain Domain

```

10      20      30      40
6G425  EIQLQQSGPELHKPGASVKISCKASGYSFSSHYHMKVVKQSHCKSLEWI
      . . . . .
F(ab)-1 EVQLVESGGGLVQPGGSLRLSCAASGYSFSSHYHMKVVRQAPGKGLEWV
      . . . . .
humIII  EVQLVESGGGLVQPGGSLRLSCAASGFSFTGHYMKVVRQAPGKGLEWV
      =====
      ++++++
      H1

```

```

      50   a              70             80 abc          90           100         110
6G425    GYIDPSNGGETTYNQKFKGKATLTVDTSSTANVHLSSLTSDSAVYFCAARGDYRYNGDWFFDVWGAGT (SEQ ID NO: 48)
            ||| | | ||||| ||| | |
F(ab)-I  GYIDPSNGGETTYNQKFGRTISRDN SKNTLYLQHNLSRAEDTAVYYCAARGDYRYNGDWFFDVWGGQT (SEQ ID NO: 49)
            ||| ||| ||||| ||| | |
humIII   GHINPDSDETRYADSVKGRTISRDN SKNTLYLQHNLSRAEDTAVYYCAARGIFY-GTTYFDYWGGQT (SEQ ID NO: 50)
                |||||
               ++++++
            ++++++
            H2                                     H3

```

FIG. 29




FIG. 30A




FIG. 30B




FIG. 30C

Amino Acid Sequence of the humanized anti-IL-8 6G4.2.5V11 Light Chain

MKKNIAFLLASMFVFSIATNAYADIQMTQSPSSLSASVGDRTTITCRSSQSLVHGIGNTY
LHWYQQKPGKAPKLLIYKVSNRFSGVPSRFSGSGGTDFLTITSSLPEDFATYYCSQST
HVPLTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDSYSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRG
EC (seq ID NO: 51)

Amino Acid Sequence of the humanized anti-IL-8 6G4.2.5V11 Heavy Chain

MKKNIAFLLASMFVFSIATNAYAEVQLVQSGGGLVQPGGSLRLSCAASGYFSFSSHYMH
WVRQAPGKGLEWVGVIDPSNGETTYNQKFKGRFTLSRDNSKNTAYLQMNSLRAEDTAVYY
CARGDYRYNGDWFDVWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF
PEPVTISWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTK
VDKKVEPKSCDKTHT (seq ID NO: 52)

Amino Acid Sequence of the peptide linker and M13 Phage Coat (gene-III)

SGGGSGGDFDYKMANANKGAMTENADENALQSDAKGKLDVATDYGAIDGFIGDVS
GLANGATGDFAGSSNSQMAQVGDGDN SPLMNNFRQYLP SLPQSVCECRPFVFSAGKPY
EFSIDCDKINLFRGVFAFLLYVATFMVVFSTFANILRNKES (seq ID NO: 53)

FIG. 31A

37 / 142

1 ATGAAAAAGA ATATCGCATT TCTTCTTGCA TCTATGTTTCG TTTTCTCTAT TGCTACAAAC
 TACTTTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTTG
 -23 M K K N I A F L L A S M F V F S I A T N
 61 GCATACGCTG ATATCCAGAT GACCCAGTCC CCGAGCTCCC TGTCCGCCTC TGTGGGCGAT
 CGTATGCGAC TATAGGTCTA CTGGGTCAGG GGCTCGAGGG ACAGGCGGAG ACACCCGCTA
 -3 A Y A D I Q M T Q S P S S L S A S V G D
 121 AGGGTCACCA TCACCTGCAG GTCAAGTCAA AGCTTAGTAC ATGGTATAGG TAACACGTAT
 TCCAGTGGT AGTGGACGTC CAGTTCAGTT TCGAATCATG TACCATATCC ACGATGCATA
 18 R V T I T C R S S Q S L V H G I G N T Y
 181 TTACACTGGT ATCAACAGAA ACCAGGAAAA GCTCCGAAAC TACTGATTTA CAAAGTATCC
 AATGTGACCA TAGTTGTCTT TGGTCTTTT CGAGGCTTTG ATGACTAAAT GTTTCATAGG
 38 L H W Y Q Q K P G K A P K L L I Y K V S
 241 AATCGATTCT CTGGAGTCCC TTCTCGCTTC TCTGGATCCG GTTCTGGGAC GGATTTCACT
 TTAGCTAAGA GACCTCAGGG AAGAGCGAAG AGACCTAGGC CAAGACCCTG CCTAAAGTGA
 58 N R F S G V P S R F S G S G S G T D F T
 301 CTGACCATCA GCAGTCTGCA GCCAGAAGAC TTCGCAACTT ATTACTGTTC ACAGAGTACT
 GACTGGTAGT CGTCAGACGT CCGTCTTCTG AAGCGTTGAA TAATGACAAG TGTCTCATGA
 78 L T I S S L Q P E D F A T Y Y C S Q S T
 361 CATGTCCCGC TCACGTTTGG ACAGGGTACC AAGGTGGAGA TCAAACGAAC TGTGGCTGCA
 GTACAGGGCG AGTGCAAACC TGTCCCATGG TTCCACCTCT AGTTTGCTTG ACACCGACGT
 98 H V P L T F G Q G T K V E I K R T V A A
 421 CCATCTGTCT TCATCTTCCC GCCATCTGAT GAGCAGTTGA AATCTGGAAC TGCTTCTGTT
 GGTAGACAGA AGTAGAAGGG CGGTAGACTA CTCGTCAACT TTAGACCTTG ACGAAGACAA
 118 P S V F I F P P S D E Q L K S G T A S V
 481 GTGTGCCTGC TGAATAACTT CTATCCCAGA GAGGCCAAAG TACAGTGGAA GGTGGATAAC
 CACACGGACG ACTTATTGAA GATAGGGTCT CTCCGGTTTC ATGTCACCTT CCACCTATTG
 138 V C L L N N F Y P R E A K V Q W K V D N
 541 GCCCTCCAAT CGGGTAACTC CCAGGAGAGT GTCACAGAGC AGGACAGCAA GGACAGCACC
 CGGGAGGTTA GCCCATTGAG GGTCTCTCTA CAGTGTCTCG TCCTGTCGTT CCTGTCTGTTG
 158 A L Q S G N S Q E S V T E Q D S K D S T
 601 TACAGCCTCA GCAGCACCCT GACGCTGAGC AAAGCAGACT ACGAGAAACA CAAAGTCTAC
 ATGTCGGAGT CGTCGTGGGA CTGCGACTCG TTTCGTCTGA TGCTCTTTGT GTTTCAGATG
 178 Y S L S S T L T L S K A D Y E K H K V Y
 661 GCCTGCGAAG TCACCCATCA GGGCCTGAGC TCGCCCGTCA CAAAGAGCTT CAACAGGGGA
 CGGACGCTTC AGTGGGTAGT CCCGGAAGTC AGCGGGCAGT GTTCTCTGAA GTTGTCCCTT
 198 A C E V T H Q G L S S P V T K S F N R G
 721 GAGTGTTAAG CTGATCCTCT ACGCCGGACG CATCGTGGCC CTAGTACGCA ACTAGTCGTA (Seq ID: 54)
 CTCACAATTC GACTAGGAGA TGCGGCCTGC GTAGCACCGG GATCATGCGT TGATCAGCAT
 218 E C O (Seq ID: 51)

FIG. 31B

Amino Acid Sequence of the humanized anti-IL-8 6G4.2.5V19 Light Chain

MKKNIAFLASMFVFSIATNAYADIQMTQSPSSLSASVGDRTITCRSSQSLVHGIGNTY
LHWYQQKPGKAPKLLIYKVSNRFSGVPSRFSGSGGTDFTLTISLQPEDFATYYCSQST
HVPLTFGQGTKEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFYFPREAKVQWKVDN
ALQSGNSQESVTEQDSKDSITYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRG
EC (seq ID NO: 51)


Amino Acid Sequence of the humanized anti-IL-8 6G4.2.5V19 Heavy Chain



MKKNIAFLASMFVFSIATNAYAEVQLVESGGGLVQPGGSLRLSCAASGYSFSSHYMH
WVKQAPGKGLEWVGYIDPSNGETTYNQKFKGRFTLSRDNSKNTAYLQMNSLRAEDTAVYY
CARGDYRYNGDWFDDVWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF
PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHHKPSNTK
VDKKVEPKSCDKTHT (seq ID NO: 55)



FIG. 31C



Fig. 32







Amino Acid Sequence of the humanized anti-IL-8 6G4.2.5V11N35A Light Chain

MKKNIAFLASMFVFSIATNAYADIQMTQSPSSLSASVGDRTTITCRSSQSLVHGIGATY
LHWYQQKPGKAPKLLIYKVSNRFSRFSGSGTDFLTITSSLPEDFATYYCSQST
HVPLTFCQGQTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDSYSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRG
EC (SEQ ID NO: 56)

Amino Acid Sequence of the humanized anti-IL-8 6G4.2.5V11N35A Heavy Chain

MKKNIAFLASMFVFSIATNAYAEVQLVQSGGGLVQPGGSLRLSCAASGYSFSSHYMH
WVRQAPGKGLEWVGYIDPSNGETTYNQKFKGRFTLSRDNSKNTAYLQMNSLRAEDTAVYY
CARGDYRYNGDWFDDVWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF
PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHHKPSNTK
VDKKVEPKSCDKTHT (SEQ ID NO: 57)

Amino Acid Sequence of the putative Pepsin Cleavage Site and GCN4 Leucine Zipper

CPPCPAPELLGGRMKQLEDKVEELLSSKNYHLENEVARLKKLVGER (SEQ ID NO: 57)

FIG. 35

44 / 142

1 ATGAAAAGA ATATCGCATT TCTTCTTGCA TCTATGTTTCG TTTTCTAT TGCTACAAAC
 TACTTTTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTTG
 -23 M K K N I A F L L A S M F V F S I A T N
 61 GCATACGCTG ATATCCAGAT GACCCAGTCC CCGAGCTCCC TGTCCGCCTC TGTGGGCGAT
 CGTATGCGAC TATAGGTCTA CTGGGTCAGG GGCTCGAGGG ACAGGCGGAG ACACCCGCTA
 -3 A Y A D I Q M T Q S P S S L S A S V G D
 121 AGGGTCACCA TCACCTGCAG GTCAAGTCAA AGCTTAGTAC ATGGTATAGG TGCTACGTAT
 TCCCAGTGGT AGTGGACGTC CAGTTCAGTT TCGAATCATG TACCATATCC ACGATGCATA
 18 R V T I T C R S S O S L V H G I G A T Y
 181 TTACACTGGT ATCAACAGAA ACCAGGAAAA GCTCCGAAAC TACTGATTAA CAAAGTATCC
 AATGTGACCA TAGTTGTCTT TGGTCTTTT CGAGGCTTTG ATGACTAAAT GTTTCATAGG
 38 L H W Y Q Q K P G K A P K L L I Y K V S
 241 AATCGATTCT CTGGAGTCCC TTCTCGCTTC TCTGGATCCG GTTCTGGGAC GGATTTCACT
 TTAGCTAAGA GACCTCAGGG AAGAGCGAAG AGACCTAGGC CAAGACCCTG CCTAAAGTGA
 58 N R F S G V P S R F S G S G S G T D F T
 301 CTGACCATCA GCAGTCTGCA GCCAGAAGAC TTCGCAACTT ATTACTGTTC ACAGAGTACT
 GACTGGTAGT CGTCAGACGT CGGTCTTCTG AAGCGTTGAA TAATGACAAG TGTCTCATGA
 78 L T I S S L Q P E D F A T Y Y C S O S T
 361 CATGTCCCGC TCACGTTTGG ACAGGGTACC AAGGTGGAGA TCAAACGAAC TGTGGCTGCA
 GTACAGGGCG AGTGCAAACC TGTCCTATGG TTCCACCTCT AGTTTGCTTG ACACCGACGT
 98 H V P L T F G Q G T K V E I K R T V A A
 421 CCATCTGTCT TCATCTTCCC GCCATCTGAT GAGCAGTTGA AATCTGGAAC TGCTTCTGTT
 GGTAGACAGA AGTAGAAGGG CGGTAGACTA CTCGTCAACT TTAGACCTTG ACGAAGACAA
 118 P S V F I F P P S D E Q L K S G T A S V
 481 GTGTGCCTGC TGAATAACTT CTATCCAGA GAGGCCAAAG TACAGTGGAA GGTGGATAAC
 CACACGGACG ACTTATTGAA GATAGGGTCT CTCCGGTTTC ATGTCACCTT CCACCTATTG
 138 V C L L N N F Y P R E A K V Q W K V D N
 541 GCCCTCCAAT CGGGTAACTC CCAGGAGAGT GTCACAGAGC AGGACAGCAA GGACAGCACC
 CGGGAGGTGA GCCCATTGAG GGTCTCTCA CAGTGTCTCG TCCTGTGCTT CCTGTCTGTTG
 158 A L Q S G N S Q E S V T E Q D S K D S T
 601 TACAGCCTCA GCAGCACCTT GACGCTGAGC AAAGCAGACT ACGAGAAACA CAAAGTCTAC
 ATGTCGGAGT CGTCGTGGGA CTGCGACTCG TTTCGTCTGA TGCTCTTGT GTTTCAGATG
 178 Y S L S S T L T L S K A D Y E K H K V Y
 661 GCCTGCGAAG TCACCCATCA GGGCTGAGC TCGCCCGTCA CAAAGAGCTT CAACAGGGGA
 CGGACGCTTC AGTGGGTAGT CCGGACTCG AGCGGGCAGT GTTCTCTGAA GTTGTCCCTT
 198 A C E V T H Q G L S S P V T K S F N R G
 721 GAGTGTTAAG CTGATCCTCT ACGCCGGACG CATCGTGGCC CTAGTACGCA ACTAGTCGTA (SEQ ID NO: 52)
 CTCACAATTC GACTAGGAGA TCGCGCCTGC GTAGCACCAG GATCATGCGT TGATCAGCAT
 218 E C O (SEQ ID NO: 56)

FIG. 36

781 AAAAGGGTAT CTAGAGGTTG AGGTGATTTT ATGAAAAAGA ATATCGCATT TCTTCTTGCA
TTTTCCCAT A GATCTCCAAC TCCACTAAAA TACTTTTCT TATAGCGTAA AGAAGAACGT
-1 M K K N I A F L L A

841 TCTATGTTTCG TTTTCTCTAT TGCTACAAAC GCGTACGCTG AGGTTCAGCT AGTGCAGTCT
AGATACAAGC AAAAAAGATA ACGATGTTTG CGCATGCGAC TCCAAGTCGA TCACGTCAGA
-11 S M F V F S I A T N A Y A E V Q L V Q S

901 GGCGGTGGCC TGGTGCAGCC AGGGGGCTCA CTCCGTTTGT CCTGTGCAGC TTCTGGCTAC
CCGCCACCGG ACCACGTCGG TCCCCGAGT GAGGCAAACA GGACACGTCG AAGACCGATG
8 G G G L V Q P G G S L R L S C A A S G Y

961 TCCTTCTCGA GTCACTATAT GCACTGGGTC CGTCAGGCCG CGGGTAAGGG CCTGGAATGG
AGGAAGAGCT CAGTGATATA CGTGACCCAG GCAGTCCGGG GCCCATTTCC GGACCTTACC
28 S F S S H Y M H W V R Q A P G K G L E W

1021 GTTGGATATA TTGATCCTTC CAATGGTGAA ACTACGTATA ATCAAAAGTT CAAGGGCCGT
CAACCTATAT AACTAGGAAG GTTACCACTT TGATGCATAT TAGTTTTCAA GTTCCCGGCA
48 V G Y I D P S N G E T T Y N O K F K G R

1081 TTCACCTTAT CTCGCGACAA CTCCAAAAAC ACAGCATACC TGCAGATGAA CAGCCTGCGT
AAGTGAAATA GAGCGCTGTT GAGGTTTTTG TGTCGTATGG ACGTCTACTT GTCGGACGCA
68 F T L S R D N S K N T A Y L Q M N S L R

1141 GCTGAGGACA CTGCCGTCTA TTA CTGTGCA AGAGGGGATT ATCGCTACAA TGGTGACTGG
CGACTCCTGT GACGGCAGAT AATGACACGT TCTCCCCTAA TAGCGATGTT ACCACTGACC
88 A E D T A V Y Y C A R G D Y R Y N G D W

1201 TTCTTCGACG TCTGGGGTCA AGGAACCCTG GTCACCGTCT CCTCGGCCTC CACCAAGGGC
AAGAAGCTGC AGACCCCACT TCCTTGGGAC CAGTGGCAGA GGAGCCGGAG GTGGTTCCCG
108 F F D V W G Q G T L V T V S S A S T K G

1261 CCATCGGTCT TCCCCCTGGC ACCCTCCTCC AAGAGCACCT CTGGGGGCAC AGCGGCCCTG
GGTAGCCAGA AGGGGGACCG TGGGAGGAGG TTCTCGTGGA GACCCCGTG TCGCCGGGAC
128 P S V F P L A P S S K S T S G G T A A L

1321 GGCTGCCTGG TCAAGGACTA CTTCCCCGAA CCGGTGACGG TGTCGTGGAA CTCAGGCGCC
CCGACGGACC AGTTCCTGAT GAAGGGGCTT GGCCACTGCC ACAGCACCTT GAGTCCGCGG
148 G C L V K D Y F P E P V T V S W N S G A

1381 CTGACCAGCG GCGTGCACAC CTTCCCGGCT GTCCTACAGT CCTCAGGACT CTACTCCCTC
GACTGGTTCG CGCACGTGTG GAAGGGCCGA CAGGATGTCA GGAGTCCTGA GATGAGGGAG
168 L T S G V H T F P A V L Q S S G L Y S L

1441 AGCAGCGTGG TGACCGTGCC CTCCAGCAGC TTGGGCACCC AGACCTACAT CTGCAACGTG
TCGTGCGACC ACTGGCACGG GAGGTCGTCG AACCCGTGGG TCTGGATGTA GACGTTGCAC
188 S S V V T V P S S S L G T Q T Y I C N V

1501 AATCACAAGC CCAGCAACAC CAAGGTCGAC AAGAAAGTTG AGCCCAAATC TTGTGACAAA
TTAGTGTTTCG GGTGCTTGTG GTTCCAGCTG TTCTTTCAAC TCGGGTTTAG AACACTGTTT
208 N H K P S N T K V D K K V E P K S C D K


1561 ACTCACACAT GCCCGCCGTG CCCAGCACCA GAACTGCTGG GCGGCCGCAT GAAACAGCTA
TGAGTGTA CGGGCGGCAC GGGTCGTGGT CTTGACGACC CGCCGGCGTA CTTTGTGCGT
228 T H T C P P C P A P E L L G G R M K Q L

FIG. 37A

1621 GAGGACAAGG TCGAAGAGCT ACTCTCCAAG AACTACCACC TAGAGAATGA AGTGGCAAGA
CTCCTGTTCC AGCTTCTCGA TGAGAGGTTC TTGATGGTGG ATCTCTTACT TCACCGTTCT
248 E D K V E E L L S K N Y H L E N E V A R

1681 CTCAAAAAGC TTGTCGGGGA GCGCTAA (SEQ ID NO: 59)
GAGTTTTTCG AACAGCCCCT CGCGATT
268 L K K L V G E R O (SEQ ID NO: 60)

FIG. 37B



Cold Ligand (nM)

FIG. 38

FIG. 39





FIG. 40



ecorI pflMI pIeI
 apoI bslI mboII taqI
 1 GAATTCAACT TCTCCATACT TTGGATAGG AATACAGAC ATGNAANAATC TCNTTGTCTGA GTTGTATATT AAGCTTGCCC AAAAGAGA AGAGTCGAAT
 CTTAAGTGA AGAGGTATGA AACCTATTC TTTATGTCTG TACTTTTAG AGAACGACT CAACAATAA TTCCAACGGG TTTTCTTCT TCTCAGCTTA
 bspMI hlnPI aluI
 hlnPI hlnPI hlnPI
 hhaI/cfoI hhaI/cfoI hhaI/cfoI
 mstI avII/fspI hlnPI
 101 GAAGTGTG CGCAGGTAGA AGCTTGGAG ATTATCTCA CTGCATGCT TCGCAATGCT TCGCAATG CGGCAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG
 CTTGACACAC GCGTCCATCT TCGAACCTC TAATAGCAGT GACGTACGA AGGTTATAC CGGTTTAC TGGTTGCG CAACTAACTA GTCCATCTC
 hlnPI hlnPI hlnPI
 hhaI/cfoI hhaI/cfoI hhaI/cfoI
 mstI avII/fspI hlnPI
 201 GAAGTGTG CGCAGGTAGA AGCTTGGAG ATTATCTCA CTGCATGCT TCGCAATGCT TCGCAATG CGGCAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG
 CTTGACACAC GCGTCCATCT TCGAACCTC TAATAGCAGT GACGTACGA AGGTTATAC CGGTTTAC TGGTTGCG CAACTAACTA GTCCATCTC
 hlnPI hlnPI hlnPI
 hhaI/cfoI hhaI/cfoI hhaI/cfoI
 mstI avII/fspI hlnPI
 301 GAAGTGTG CGCAGGTAGA AGCTTGGAG ATTATCTCA CTGCATGCT TCGCAATGCT TCGCAATG CGGCAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG
 CTTGACACAC GCGTCCATCT TCGAACCTC TAATAGCAGT GACGTACGA AGGTTATAC CGGTTTAC TGGTTGCG CAACTAACTA GTCCATCTC

FIG. 41A

scrFI
 nciI
 mepI
 hpaII
 dsav
 xmaI/pspAI
 smaI
 scrFI
 nciI
 dsav
 cauII
 bsaJI
 avai
 rsaI
 csp6I
 nlaIV
 kpnI cauII dpmII(dam-)
 hgiCI
 bani bsaJI alwI(dam-)
 asp718
 acc65I
 401 TCGGTACCCG GGGATCCTCT CGAGGTGAG GIGATTTAT GAAAGAAT ATGCGATTTT TTCTGCATC TATGTTCTGTT TTTTCTATTG CTACAAACGC
 AGCCATGGCG CCTAGGAGA GCTCCAATC CACTAAATA CTTTCTTA TAGCGTAAAG AAGACGTAG ATACAAGCAA AAAAGATAAC GATGTTGCG
 M K K N I A F L L A S M F V F S I A T N A
 a mutation was found that inactivated the mluI site. The penultimate nucleotide was changed fr G to T ^

sstI
 sacI
 hgiIII
 hgiAI/aspHI
 ecl136II
 bsp1286
 bsiHKAI
 bsmFI bmyI
 bsrI avai aluI
 tth111I/aspI banII
 ecovR
 501 ATACGCTGAT ATCCAGATGA CCCAGTCCC GAGTCCCTG TCCGCCTCTG TGGCGATAG GGTCAACATC ACCTGCAGGT CAAGTCAAG CTTACTACAT
 TATCGACTA TAGGTCTACT GGTGAGGAG CCGGAGAC AGCGGTATC ACCGCTATC CCAGTGGTAG TGGACGTCCA GTTCAGTTTC GAATCATGTA
 I Q H T Q S P S S L S A S V G D R V T I T C R S S Q S L V H

mboII sfaNI
 hphI
 maeIII
 bstEII hphI bsgI
 hphI
 maeIII
 bstEII hphI bsgI
 bspMI
 sse8387I
 pstI
 scfI
 bspMI
 ddeI nlaIII
 aluI rsaI
 hindIII csp6I
 CAAGTCAAG CTTACTACAT

FIG. 41B

FIG. 41C

FIG. 41D

FIG. 41E

pleI
hinfI
taqI
xhoI
paer7I
avaI maeIII
11401 CTTCTCGACT CACTATATGC ACTGGGTCCG TCAGGCCCGG GGTAGGGCC TGGATGGGT TGGATATATT GATCCTTCCA ATGGTGAAC TACGTATAAT
GAAGAGCTCA GTGATATACG TGACCCAGGC AGTCCGGGCG CCATCCCGG ACCTTACCCA ACCTATATNA CTAGGAAGGT TACCACITTG ATGCATATTA
29 F S S H Y M H W V R Q A P G K G L E W V G Y I D P S N G E T T Y N

bsaJI dsav
avaI bstNI
bsaJI bsli
sau96I apyI[dcM+] mboI/ndeII[dam-] bsli
nlaIV sau96I sau3AI
haeIII/paII asuI dpmI[dam+] maeII
asuI ecoO109I/draII dpmII[dam-] snaBI
ecoO109I/draII haeIII/paII alwI[dam-] hphI bsaAI
1501 CTTCTCGACT CACTATATGC ACTGGGTCCG TCAGGCCCGG GGTAGGGCC TGGATGGGT TGGATATATT GATCCTTCCA ATGGTGAAC TACGTATAAT
GTTTTCAGT TCCCGGCCAA GTGAATAGA GCGCTGTGA GGTTTTGTG TCGTATGAC GTCTACTGT CCGACGCACG ACTCCTGTGA CCGCAGATAA
62 Q K F K G R F T L S R D N S K N T A Y L Q M N S L R A E D T A V Y Y

thai
fnuDII/mvNI scfI
bstOI pstI
bsh1236I bsgI
nruI bspMI
haeIII/paII cac8I moli
sau96I cac8I ddeI drdI
asuI AGGCCGTTT CACTTATCT CCGGACNACT CCAAAACAC AGCATACTG CAGATGAACA GCTCGGTGC TGAGGACACT GCCGCTATT
TCCCGGCCAA GTGAATAGA GCGCTGTGA GGTTTTGTG TCGTATGAC GTCTACTGT CCGACGCACG ACTCCTGTGA CCGCAGATAA
62 Q K F K G R F T L S R D N S K N T A Y L Q M N S L R A E D T A V Y Y

sau96I
haeIII/paII
sau96I
nlaIV
hgiJII
bsp1286
bsp120I
maeIII
bstEII
scrFI
mvaI
ecoRII
dsaV
bstNI
bsaJI hphI bsmBI
nlaIV apyI[dcM+] bsmAI
maeIII/paII ecoO109I/draII
1601 ACTGTGCAAG AGGGGATTAT CGCTACATG GTGACTGGTT CTTGACGTC TGGGTCAAG GMACTTGGT CACGTCTCC TCGGCTTCCA CCAAGGGCCC
TGACACGTTT TCCCCTATA CGGATGTAC CACTGACCAA GAAGCTGCAG ACCCCAGTTC CTTGGACCA GTGCAGAGG AGCCGGAGGT GGTTCGCGG
96 C A R G D Y R Y N G D W F F D V W G Q G T L V T V S S A S T K G P
seq right is from p6G425chim2.fab2

FIG. 41F

[illegible]

FIG. 41G

fnu4HI
 bsoFI
 haeIII/pali
 mcrI
 eagI/xmaIII/ecI XI
 eaeI
 cfrI
 .bsiEI
 noti
 fnu4HI
 bsoFI
 nlaIII
 nspi
 bsp1286
 nspBI
 acII bmyI
 cac8I
 nlaIII
 nspi
 bsp1286
 acII bmyI
 CCGCGTGGC CAGCACAGA ACTGCTGGC
 2001 TCACACATGC CCGCGTGGC CAGCACAGA ACTGCTGGC GCGCGCATGA AACAGCTAGA GGACAAAGTC GAAGAGCTAC TCTCCAAGAA CTACCACCTA
 AGTGTGTAGG GCGCGCACGG GTCGTGTCT TGACGACCG CCGCGTACT TTGCGATCT CCTGTTCCAG CTCTCGATG AGAGTTCTT GATGGTGGAT
 229 H T C P P C P A P E L L G G R M K Q L E D K V E E L L S K N Y H L
 ^junction between antibody and leucine zipper
 scrFI
 nciI
 mspI
 hpaiI
 dsav
 cauII
 acII
 fnu4HI
 bsoFI
 rmai
 maeI
 bfaI
 bsmFI
 sau96I
 pleI
 haeIII/pali
 asuI
 hinfI
 acII
 fnu4HI
 bsoFI
 2101 GAGAAATGAG TGGCAAGACT CAAAAGCTT GTCGGGAGC GCTAAGCATG CGACGGCCT AGAGTCCCTA ACGTCGGT GCGCGCGGC GTTTTATT
 CTCTTACTTC ACCGTTCTGA GTTTTTCGA CAGCCCTCG CGATTCGTAC GCTGCGGGA TCTCAGGAT TCGAGCCAA CCGCGCGCCG CAAAATAA
 262 E N E V A R L K K L V G E R O (354 bp) 60
 aluI
 taqI
 hindIII
 clal/bsp106
 tru9I
 msel
 bspDI[dam-]
 msel
 bspDI
 aluI
 2201 GTTAACTCAT GTTGACAGC TTATCATCGA TAAGCTTTAA TCGGGTAGTT TATCACAGTT AAATTGCTAA CCGAGTCAGG CACCGGTAT GAATCTAAC
 CAACTGAGTA CAACTGTCG AATAGTAGCT ATTGGAATT AGCCATCAA ATAGTGTCAA TTTAAGGATT CCGTCAGTCC GTGGACATA CTTTAGATTG

FIG. 41H

haeIII/palI
 sau96I
 scrFI
 nciI
 mspI mnlI
 rsaI
 csp6I hpaII
 mspI dsav bsII
 hpaII caulI
 cfr10I/bsrFI asuI
 acII
 ecorV
 sfanI
 scrFI
 mvaI
 ecorII
 dsav
 bstNI
 bsajI
 hphI apyI[dam+]
 mnlI hglCI
 bsajI
 hlnPI
 hhal/cfoI
 foki
 maeIII
 foki
 scfI
 2301 ANTGGCTCA TCGTCATCCT CGGCACCGTC ACCCTGGATG CTGTAGGCAT AGGCTTGTT ATGCCGGTAC TGCCGGGCCT CTGCGGGAT ATCGTCCATT
 TTACGGGAGT AGCAGTAGGA GCGTGGCAG TGGACCTAC GACNTCCGA TCCGACCAA TACGGCCATG ACGGCCCGGA GAAGCCCTA TAGCAGGTAA
 hlnPI
 hhal/cfoI
 xmai
 maeI
 nheI
 fnu4HI haeII
 bsOFI eco47III
 bbvI bfaI
 maeIII
 cac8I
 bsrI
 sfanI
 cac8I
 cac8I
 cac8I
 2401 CCGACAGCAT CCGCAGTCAC TATGGCGTGC TGCTAGCGCT ATATGCGTTG ATGCAATTTC TATGCCGACC CGTTCTCGGA GCACGTGCG ACCGCTTTGG
 GCGTGTGTA GCGGTGAGT ATACCGCAGC ACGATCGCGA TATACGCAAC TAGCTTAAG ATACGCGTGG GCAAGAGCCT CGTGACAGGC TGCGAACC
 fnu4HI
 bs FI
 haeIII/palI
 acII
 mcrI
 bsIKAI
 bsIEI
 bmyI
 hglAI/aspHI
 bsp1286
 hlnPI
 hhal/cfoI
 mstI bsII
 avII/fspi
 sfanI
 sau3AI
 mboI/ndeII[dam-]
 dpnI[dam+]
 dpnII[dam-]
 alwI[dam-]
 nlaIV
 bstXI/xhoII
 bamHI
 alwI[dam-]
 hpaII sfanI
 mspI
 hgaI
 acII
 fnu4HI
 bsOFI
 cac8I
 cac8I
 cac8I
 2501 CCGCGGCCCA GTCCTGCTCG CTTCGCTACT TGGAGCCACT ATCGACTACG CGATCATGCG GACCAACCC GTCTGTGGA TCCTCTACGC CGGACGCATC
 GCGCGCGGGT CAGGACGAGC GAAGCGATGA ACCTCGGTGA TACCTGATGC GCTAGTACCG CTGGTGTGGG CAGGACACCT AGGAGATGCG GCCTGCGTAG

FIG. 411

[illegible]

FIG. 41J

[illegible]

FIG. 41L

hpaI hpaI
 hpaI/cfoI hpaI/cfoI
 mstI mstI
 avIII/fspI avIII/fspI
 bsmI bsmI
 acII acII
 nlaIV nlaIV
 pflMI pflMI
 bslI bslI
 hphI hphI
 tfII tfII
 hinfI hinfI
 3501 CTAACGGATT CACCACCTCA AGAATTGGAG CCAATCAATT CTTGGGAGA ACTGTGAATG CGCAACCAA CCCTTGGCAG AACATATCCA TCGCGTCCGC
 GATTGCTAA GTGGTGGGT TCTTAACCTC GTTAGTTAA GAAGCCTCT TGACACTTAC CGGTTGGTT GGAACCCGTC TTGTATAGGT AGCGAGGCG
 haeIII/palI haeIII/palI
 mscI/balI mscI/balI
 haeI haeI
 scrFI scrFI
 mvaI dsaI mvaI dsaI
 ecorII ecorII
 dsav dsav
 bstNI bstNI
 bslI bsaJI bslI bsaJI
 apyI(dcm+) apyI(dcm+)
 sau96I sau96I
 avall avall
 asuI eaeI asuI eaeI
 ppuHI ppuHI
 nlaIV cfrI nlaIV cfrI
 ecoO109I/draII ecoO109I/draII
 bpmI/gsuI(dcm-) acII sfaNI bpmI/gsuI(dcm-) acII sfaNI
 3601 CATCTCCAGC AGCCGCACGC GCGCATCTC GGCAGCGTT GGTCTCTGGC CACGGGTGCG CATGATCGTG CTCCTGTCTG TGAGGACCCG GCTAGGCTGG
 GTACAGGTGCG TCGGCGTGG CCGGTAGAG CCCGTCCGCA
 fnu4HI fnu4HI
 thaI hlnPI thaI hlnPI
 fnu4HI bsoFI fnu4HI bsoFI
 bsoFI fnuDII/mvnI bsoFI fnuDII/mvnI
 fnu4HI bstUI fnu4HI bstUI
 cac8I cac8I
 hhaI/cfoI hhaI/cfoI
 bsh1236I bsh1236I
 bbvI acII bsh1236I bbvI acII
 bsmI/gsuI(dcm-) acII sfaNI bsmI/gsuI(dcm-) acII sfaNI
 3701 CGGGGTGTC TTACTGTTA GCAGATGAA TCACCGATAC CGGAGCGAAC GTGAAGGCAC TGCTGCTGCA AAACGTCTGC GACCTGAGCA ACAACATGAA
 GCCCCAACGG AATGACCAAT CGTCTTACTT AGTGGCTATG CGCTCGCTTG CACTTCCGTG ACGACGACGT TTTGCAGACG CTGGACTCGT TGTGTACTT
 hgaI hgaI
 thaI acII thaI acII
 fnuDII/mvnI fnuDII/mvnI
 bstUI bstUI
 bsh1236I bsh1236I
 nlaIII nlaIII
 ddeI ddeI
 maeII maeII
 fnu4HI fnu4HI
 bsoFI bsoFI
 bbvI bbvI
 fnu4HI fnu4HI
 bsoFI bsoFI
 bbvI bbvI
 maeII maeII
 nlaIII nlaIII

FIG. 41M

sau3AI
mboI/ndelII[dam-]
mamI[dam-]
dpuI[dam+]
dpuII[dam-]
bstYI/xhoII
alwI[dam-]
mspI
hpaII
mroI bsaBI[dam-]
bspHII
bspEI[dam-]
bsaWI sfaNI
accIII[dam-] foki cac8I
TTATGTTCCG GATCGCATC GCAGGATGCT GCTGGCTACC
CTACACGCTAG CTGCTTACGA GCACCGATGG
3801 TGGTCTTCGG TTTCGGTGT TCGTAAAGTC TGGAAACCGG GAAGTCAGCG CCCTGCACCA
ACCAAGAGCC AAGGCACAA AGCATTTTCAG ACCTTTGGCG CTTCACTCGC GGGACGTGGT AATACAAGGC
cac8I
hinPI
hhaI/cfoI
tr9I haeII
mseI eco47III
GATGATAGAC ATAATGCTT CGCGNCCGTA ACTGGGACTC ACTAANAAGA GACCAGGGCG GCGTAGGTAT GCGGTCAAC AATGGGAGT
3901 CTGTGGAACA CCTACATCTG TATTAACGAA CGGCTGGCAT TGACCCCTGAG TGATTTTCT CTGGTCCCGC CGCATCCATA CGGCCAGTTG TTTACCTCA
GACACCTTGT GGATGATAGAC ATAATGCTT CGCGNCCGTA ACTGGGACTC ACTAANAAGA GACCAGGGCG GCGTAGGTAT GCGGTCAAC AATGGGAGT
4001 CAACGTTCCA GTAACCGGC ATGTTTCATCA TCAGTAACCC GTATCGTGAG CATCCTCTCT CGTTTCATCG GTATCATTAC CCCCATGAAC AGAATTCCC
GTTGCAAGGT CATGGCCCC TACAAGTAGT AGTCATTGGG CATAGCACTC GTAGGAGAGA GCAAGTAGC CATAGTAATG GGGGTACTTG TCTTTAAGG

cac8I
sau96I
tru9I haeIII/palI
mseI
sfanI maeIII
mnII
bpmI/gsuI(dcm-)

4101 CCTTACACGG AGGCATCAAG TGACCAACA GGA AAAAACC GCCCTTAACA TGSCCGCTT TATCAGAAGC CAGACATTAA CGCTTCTGGA GAAACTCAAC
GGATGTGCC TCCGAGTTC ACTGGTTTGT CCTTTTITGG CGGGAATTGT ACCGGGCGAA ATAGTCTTCG GTCTGTAATT CGGNAGACCT CTTTGAGTTG

4201 GAGCTGGACG CGGATGAACA GGCAGACATC TGTGAATCGC TTCACGACCA CGCTGATGAG CTTTACCGCA CTGCTCTCGC GCGTTTCGGT GATGACGGTG
CTCGACCTGC GCCTACTTGT CCGTCTGTAG ACACTTAGCG AAGTGTCTGT GCGACTACTC GAAATGGCCT CGACGGAGCG CGCAAGCCA CTACTGCCAC

4301 AAAACCTCTG ACACATGCAG CTCCTGGAGA CGGTCAACGC TTCTCTGTAA CGGGATGCCG GGAGCAGACA AGCCCGTTCAG GCGCGTCAG CGGGTGTTCG
TTTTGGAGAC TGTGTACGTC GAGGGCCTCT GCCAGTGTGC AACAGACATT CGCCTACGCC CCTCTCTGT TCGGGCAGTC CGCGGAGTC GCCACAACC

FIG. 410

FIG. 41P

FIG. 41Q

mspI
 hpaiI
 sau3AI
 mboI/ndeII[dam-]
 dpnI[dam+]
 dpnII[dam-]
 aluI
 maeIII
 eco57I bsrI
 hhaI/cfoI
 thaI
 fnuDII/mvuI
 bstUI
 bsh1236I
 5301 ATTACGCGCA GAAAAAAGG ATCTCAGAA GATCCTTGA TCTTTTCTAC GGGGTCTGAC GCTCAGTGA ACAGAACTC AGGTTAAGG ATTTTGGTCA
 TAATGCGCGT CTTTTTTC TAGAGTTCTT CTAGGAACCT AGAAAGATG CCCAGACTG CGAGTCACTT GCCTTTGAG TGCATTTCCC TAAACCAGT
 nlaIII
 rcaI
 bspHI
 tru9I
 mseI
 maeII
 hgaI ddeI
 sau3AI
 mboI/ndeII[dam-]
 mboI/ndeII[dam-]
 sau3AI mboII[dam-] dpnI[dam+]
 mboI/ndeII[dam-] dpnII[dam-]
 sau3AI mboII[dam-] dpnI[dam+]
 dpnI[dam+] dpnI[dam+]
 dpnII[dam-] dpnII[dam-]
 bstYI/xhoII aluI[dam-]
 aluI[dam-] bstYI/xhoII
 aluI/draI
 sau3AI
 mboI/ndeII[dam-]
 rmaI
 hphI dpnI[dam+]
 mboII[dam-]
 sau3AI maeI tru9I
 mboI/ndeII[dam-] mseI
 dpnI[dam+] dpnII[dam-]
 dpnII[dam-] aluI[dam-] tru9I
 bstYI/xhoII bstYI/xhoII mseI
 aluI[dam-] bfaI ahaIII/draI
 5401 TGACATTATC AAAAAGGATC TTCACCTAGA TCTTTTAA TTAATAATGA AGTTTAAAT CAATCTAAAG TATATATGAG TAAACTTGGT CTGACAGTTA
 ACTCTAATAG TTTTTCCTAG AAGTGGATCT AGGAATTTT AATTTTACT TCMAAATTTA GTTAGATTTC ATATATACTC ATTTGAACCA GACTGTCAAT
 nlaIV
 hgiCI
 banI
 mnlI
 ddeI
 sau3AI
 mboI/ndeII[dam-]
 dpnI[dam+]
 dpnII[dam-]
 foki
 pleI
 hinfI
 ahdI/eam1105I
 mnlI
 5501 CCAATGCTTA ATCAGTGAGG CACCTATCTC AGCGATCTGT CTATTTTCGT CATCCATAGT TGCTGACTC CCCGTCGT AGATAACTAC GATACGGGAG
 GGTTAGGAAT TAGTCACCTC GTGGATAGAG TCGCTAGACA GATAAAGCA GTAGGTATCA ACCGACTGAG GGGCAGCACA TCTATTGATG CTATGCCCTC

FIG. 41B

[illegible]

FIG. 41S

```

mcrI
bsiEI
bcgI
fnu4HI
bsoFI
acII
rsal
bsrI
scal
maeIII hphI csp6I
sfanI
6001 ATTCTCTTAC TGTGATGCCA TCCGTAAGAT GCTTTTCTGT GACTGGTGAG TACTCAACCA AGTCATTCTG AGAATAGTGT ATGCGGCGAC CGAGTTGCTC
TAAGAGAATG ACAGTACGGT AGGCATTCTA CGAAAGACA CTGACCACCTC ATGAGTTGGT TCAGTAAGAC TCTTATCACA TACGCCGCTG GCTCAACGAG
        hgaI
        hinII/acyI
        ahaII/bsaHI
        mspI
        hpaII
        scrFI
        nciI
        dsav
        cauII hincII/hindII
        hlnPI
        hhaI/cfoI
        thal
        fnuDII/mvni
        bstUI
        bsh1236I
        acII
        hgaI/aspHI
        bsp1286
        tru9I bsiHKAI
        mseI bmyI
        ahaII/draI
        asp700 mboII
        6101 TTGCCCGCGG TCAACACGGG ATAATACCGC GCCACATAGC AGAATCTTAA AGTGCTCAT CATTTGAAA CGTTCTCGG GCGGAAACT CTCAGGATC
AACGGCGCGC AGTGTGCCC TATTATGGC GGTGTATCG TCTTGAATT TTCACGAGTA GTNACCTTT GCNAGAAGCC CGCTTTTGA GAGTTCCTAG
        bsrI
        sau3AI
        mboI/ndeII(dam-)
        dpaI(dam+)
        dpaII(dam-)
        alwI(dam-)
        bstYI/xhoII
        alwI(dam-)
        hphI
        nsp8II
        acII
        bstYI/xhoII
        maeIII bssSI
        6201 TTACCGTGT TGAGATCCAG TTCGATGTAA CCACCTCGTG CACCCACTG ATCTTCAGCA TCTTTTACTT TCACCAGCGT TTCTGGGTGA GCAAAAACAG
AATGGGACA ACTCTAGGTC AAGCTACATT GGTGAGCAGC GTGGGTGAC TAGAAGTCGT AGAAATGAA AGTGGTCGCA AGACCCCACT CGTTTGTG
        acII
        fnu4HI
        bsoFI
        mboII
        earI/ksp632I sspI
        6301 GAAGGCAAAA TGCCGCAAAA AAGGGAATAA GGGGACACG GAAATGTTGA ATACTCATAC TCTTCTTTT TCAATATTAT TGAAGCATTT ATCAGGGTTA
CTTCCGTTTT ACGGCGTTTT TTCCCTTATT CCCGCTGTGC CTTTCAACT TATGAGTATG AGAAGGAAA AGTTATAATA ACTTCGTAAA TAGTCCCAAT

```

FIG. 41T

hinPI
 thai
 fnuDI1/mvni
 bstOI
 bsh1236I
 acII
 nlaIV hhai/cfoI
 nlaIII
 rcaI
 bspHI acII
 bsmAI bsrBI
 bsmAI
 TGTCTCATG AGCGGATACA TATTGCAATG TATTAGNAA AATAACMAA TAGGGTTCC GCGCACATTT CCCCAGAAAG TGCACCTGA CGTCTAAGAA
 AACAGAGTAC TCGCCTATCT ATAACTTAC ATAAATCTTT TTATTGTTT ATCCCCAAGG CGCGTGTAAG GGGGCTTTTC ACGGTGGACT GCAGATCTT
 6401

sau96I
 haeIII/pali
 asuI
 ecoOI09I/draII
 mnlI
 bssSI
 rcaI tru9I
 bspHI mseI
 nlaIII
 bpuAI
 bbsI
 CTTTCGTCTT CAA
 GAAAGCAGAA GTT
 6501

(SEQ ID NO: 61)

FIG. 41U


```

>length: 6563

aatII(GACGTC): 1645 6489
acc65I(GGTACC): 403 823
accI(GTHKAC): 1093 1963 4449
accIII(TCCGGA): 3867(dam-)
acII(CCGC): 178 542 805 877 1340 1750 1826 2011 2039 2043 2182 2242 2384 2492 2501 2504
2628 2781 2784 2787 2906 2926 3005 3045 3094 3141 3226 3241 3309 3342 3367 3412
3436 3448 3490 3544 3597 3613 3619 3700 3838 3967 3970 3981 4139 4155 4210 4266
4351 4390 4400 4442 4467 4505 4518 4544 4561 4604 4611 4632 4723 4751 4878 4897
5018 5128 5263 5272 5634 5725 5916 5962 6083 6127 6204 6313 6412 6459
see hnlI
acyI 1307 4678
afIII(ACRYGT): 1788
ageI(ACCGT): 1645 1813 2616 2637 2751 3408 6107 6489
ahaII/psaHI(GRCGYC): 5435 5454 6146
ahaII/draI(TTAAA): 346 5566
ahdI/eam1105I(GACNNNNNGTC): 72 121 252 320 398 532 589 648 1126 1144 1167 1325 1386 1906 2054 2075 2126
alulI(AGCT): 2218 2233 2889 3292 4202 4259 4270 4319 4338 4619 4845 4935 4981 5238 5759 5859
5922
alw44I/snoI(GTGCAC): 1831 4494 4992 6238
alwI(dam-)(GGATC): 412 413 712 713 1171 1471 2578 2579 3300 3870 5245 5319 5331 5416 5429 5893
6196 6214
alwNI(dcm-)(CAGNNNCTG): 1117 1385 5089
apaI(GGGCCC): 1695
apaLI/snoI(GTGCAC): 1831 4494 4992 6238
apoI(RAATTY): 1 391 4093
apyI(dcm+)(CCWGG): 640 999 1347 1357 1449 1665 1713 1755 1764 2333 3262 3645 4705 4826 4839
aseI/asnI/vspI(ATTAAAT): 5742
asnI see aseI
asp700(GAANNNTTC): 905 930 4234 6166
asp718(GGTACC): 403 823
asphi see hgiAI
aspi see tth111I
asuI(GGNCC): 1119 1195 1425 1434 1446 1512 1695 1696 1752 2155 2375 2727 3002 3090 3339 3463

```

FIG. 41V

Stop Template Primer

SL.97.2 5' CAT GGT ATA GGT TAA ACT TAT TTA CAC 3' (SEQ ID NO: 63)

NNS Randomization Primer

SL.97.3 5' CAT GGT ATA GGT NNS ACT TAT TTA CAC 3' (SEQ ID NO: 64)


FIG. 42

Randomization of Position N35 of Variable Light Chain CDR-1
Amino Acid Frequency

Phage Display (NNS Codon Library) Sort #3

Amino Acid	Frequency	% Total	IC50 (nM)
Asparagine (wt)	1	5.6	4.9
Glycine	6	16.6	3.1
Aspartic Acid	3	16.6	3.1
Glutamic Acid	4	22.2	0.1
Alanine	2	5.6	0.2
Lysine	1	5.6	ND
Serine	1	1.9	ND

FIG. 43A



1 Fig 43B




Fig. 43C




Fig. 43D





Fig. 43E



Representative Conc versus Time Plot. Shown is the kinetic data for 6G4V11N35A.F(ab')₂.

SAMPLE	ka	kd	Kd
6G4V11N35A-Fab	ND	ND	114pM
6G4V11N35A-F(ab') ₂	2.0x10 ⁶	2.1x10 ⁻⁴	109pM
6G4V11N35E-Fab	4.7x10 ⁶	2.6x10 ⁻⁴	54pM

FIG. 44

77 / 142

1 ATGAAAAAGA ATATCGCATT TCTTCTTGCA TCTATGTTTCG TTTTCTCTAT TGCTACAAAC
 TACTTTTCT TATAGCGTAA AGAAGAACGT AGATACAAGC AAAAAAGATA ACGATGTTTG
 -23 M K K N I A F L L A S M F V F S I A T N
 61 GCATACGCTG ATATCCAGAT GACCCAGTCC CCGAGCTCCC TGTCCGCCTC TGTGGGCGAT
 CGTATGCGAC TATAGGTCTA CTGGGTCAGG GGCTCGAGGG ACAGGCGGAG ACACCCGCTA
 -3 A Y A D I Q M T Q S P S S L S A S V G D
 121 AGGGTCACCA TCACCTGCAG GTCAAGTCAA AGCTTAGTAC ATGGTATAGG TGAGACGTAT
 TCCAGTGGT AGTGGACGTC CAGTTCAGTT TCGAATCATG TACCATATCC ACTCTGCATA
 18 R V T I T C R S S O S L V H G I G E T Y
 181 TTACACTGGT ATCAACAGAA ACCAGGAAAA GCTCCGAAAC TACTGATTTA CAAAGTATCC
 AATGTGACCA TAGTTGTCTT TGGTCCTTTT CGAGGCTTTG ATGACTAAAT GTTTCATAGG
 38 L H W Y Q Q K P G K A P K L L I Y K V S
 241 AATCGATTCT CTGGAGTCCC TTCTCGCTTC TCTGGATCCG GTTCTGGGAC GGATTTCACT
 TTAGCTAAGA GACCTCAGGG AAGAGCGAAG AGACCTAGGC CAAGACCCTG CCTAAAGTGA
 58 N R F S G V P S R F S G S G S G T D F T
 301 CTGACCATCA GCAGTCTGCA GCCAGAAGAC TTCGCAACTT ATTACTGTTT ACAGAGTACT
 GACTGGTAGT CGTCAGACGT CGGTCTTCTG AAGCGTTGAA TAATGACAAG TGTCTCATGA
 78 L T I S S L Q P E D F A T Y Y C S O S T
 361 CATGTCCCGC TCACGTTTGG ACAGGGTACC AAGGTGGAGA TCAAACGAAC TGTGGCTGCA
 GTACAGGGCG AGTGCAAACC TGTCCCATGG TTCCACCTCT AGTTTGCTTG ACACCCGACGT
 98 H V P L T F G Q G T K V E I K R T V A A
 421 CCATCTGTCT TCATCTTCCC GCCATCTGAT GAGCAGTTGA AATCTGGAAC TGCTTCTGTT
 GGTAGACAGA AGTAGAAGGG CCGTAGACTA CTCGTCAACT TTAGACCTTG ACGAAGACAA
 118 P S V F I F P P S D E Q L K S G T A S V
 481 GTGTGCCTGC TGAATAACTT CTATCCCAGA GAGGCCAAAG TACAGTGGAA GGTGGATAAC
 CACACGGACG ACTTATTGAA GATAGGGTCT CTCCGGTTTC ATGTCACCTT CCACCTATTG
 138 V C L L N N F Y P R E A K V Q W K V D N
 541 GCCCTCCAAT CGGGTAACTC CCAGGAGAGT GTCACAGAGC AGGACAGCAA GGACAGCACC
 CGGGAGGTTA GCGCATGAG GGTCTCTCA CAGTGTCTCG TCCTGTCTGT CCTGTCTGTG
 158 A L Q S G N S Q E S V T E Q D S K D S T
 601 TACAGCCTCA GCAGCACCTT GACGCTGAGC AAAGCAGACT ACGAGAAACA CAAAGTCTAC
 ATGTCGGAGT CGTCGTGGGA CTGCGACTCG TTTCGTCTGA TGCTCTTTGT GTTTCAGATG
 178 Y S L S S T L T L S K A D Y E K H K V Y
 661 GCCTGCCAAG TCACCCATCA GGGCCTGAGC TCGCCCGTCA CAAAGAGCTT CAACAGGGGA
 CGGACGCTTC AGTGGGTAGT CCGGACTCG AGCGGGCAGT GTTCTCTGAA GTTGTCCCTT
 198 A C E V T H Q G L S S P V T K S F N R G
 721 GAGTGTTAAG CTGATCCTCT ACGCCGGAGC CATCGTGGCC CTAGTACGCA ACTAGTCGTA (SEQ ID NO: 15)
 CTCACAATTC GACTAGGAGA TGGGCGCTGC GTAGCACCGG GATCATGCGT TGATCAGCAT
 218 E C O (SEQ ID NO: 62)

FIG. 45




FIG. 46

N35AH1upr
5'-TAGTGCAGTCTGGCGGTGGCCTGGTGCAGCCAGGGGCTCACTCCGTTTGTCTGTGCAGCTTCTGGCTACTCCTTC-3' (SEQ ID NO: 66)

N35AH1lwr
5'-TCGAGAAGGAGTAGCCAGAAGCTGCACAGGACAAACGGAGTGAGCCCCCTGGCTGCACAGGCCACCGCCAGACTGCACT-3' (SEQ ID NO: 67)

AG-3'

79 / 142

Bold indicates nucleotide change destroying PvuII site.

FIG. 47

> length: 8120 (circular)
 > This has the PSVI backbone with the PRK7 cloning linker (PSVI7) and the intron DHFR(ID)
 > This has the PSVI backbone with the PRK7 cloning linker (LL) into the HpaI site
 > made from PSVI.WTSD.D by adding a linearization linker(LL) into the HpaI site

```

cac8I
alul
sati
saci
hglJII
hgIAI/aspHI
ecII36II
bsp1286
bslHKAI
bmyI
banII
tagI
1 TTCGAGCTCG CCCGACATTG ATTATTGACT AGAGTCGATC GACAGCTGTG GAATGTGTCT CAGTTAGGCT GTGGAAAGTC CCCAGGCTCC CCAGCAGGCA
AAGCTCGAGC GGGCTGTAAC TAATACTGA TCTCAGCTAG CTGTCGACAC CITACACACA GTCAATCCCA CACCTTTCAG GGGTCCGAGG GGTCGTCCGT
cac8I
sau3AI alul
mboI/ndelI[dam-]
dpmI[dam+]
pvul/bspCI
pleI dpmII[dam-]
hinfi taqI[dam-]
rmaI mcrI pvulI
maeI bseI nspBI
bfai taqI[dam-]
sfanI
ppu10I
nslI/avaIII
nlaIII
sphi
nspI
nspHI
cac8I
101 GAAGTATGCA AAGCATGCAT CTCATTAGT CAGCAACCGC GTGTGGAAAG TCCCAGGCT CCCAGCAGG CAGAGTATG CAAGCATGC ATCTCAATTA
CTTCATACGT TTCGTACGTA GAGTTAATCA GTCGTTGGTC CACACCTTTC AGGGTCCGA GGGGTCTGCC GTCTTCATAC GTTCTGTTAG TAGAGTTAAT
nlaIII
styI
ncol
bsaI dsal
acII bsrI acII
acII foki
bsmFI
201 GTCAGCAACC ATAGTCCCGC CCTTAACCTC GCCATCCCG CCCCTACTC CGCCAGTTC CGCCATTCT CGCCCATCT CGCCCATCT GCTGACTAAT TTTTATTATT
GAGTCGTTGG TATCAGGGCG GGGATTGAG GGGTAGGGC GGGATTGAG GCGGTCMAG CCGGGTAAGA GCGGGGTGAC GCGGCTAAT TTTTATTATT

```

FIG. 48A

[illegible]

FIG. 48B

601 CAAAGCATGA CCACAACCTC TTCAGTGGAA GGTAACAGA ATCTGGTGAT TATGGGTAGG AARACCTGGT TCTCCATTCC TGAGAAGAAAT CGACCTTTAA
 GTTTCTTACT GGTGTGGAG AGTCACCTT CCAATTGTCT TAGACCACTA ATACCACATCC TTTTGACCA AGAGGTAAGG ACTCTTCTTA GCTGGAAATT
 701 AGGACAGAAAT TAATATAGTT CTCAGTAGAG AACTCAAGA ACCACCACGA GGAGCTCAT TTTTGGCCAA AGTTTGGAT GATGCCCTAA GACTTATTGA
 TCTGTCTTA ATTATATCA GAGTCATCTC TTGAGTTTCT TGGTGTGCT CCTGGAGTAA AAGAAGCGTT TTCAAACCTA CTACGGAAAT CTGAATAACT
 801 ACNACCGGAA TTGGCAAGTA AGTAGACAT GGTGTGGATA GTCGGAGGCA GTTGTGTTA CCAGGAGGCC ATGANTCAAC CAGGCCACCT TAGACTCTTT
 TGTGGCCTT AACCGTTCAT TTCATCTGTA CCNACCTAT CAGCCTCGT CAAGACAAAT GGTCCTCGG TACTTAGTTG GTCCGGTGA ATCTGAGAAA

scrFI mvaI
 mvaI ecorII
 dsav bstNI
 apyI(dcm+) ddeI mboII taqI ahaIII/draI
 sexAI
 tfII hinfI
 tru9I mseI
 sstI
 sacI
 hgiIII
 hgiAI/asplI
 eciI36II
 bspI286
 bstHKAI
 bmyI
 mnlI aluI
 bssI banII
 balI bseRI
 ddeI
 tru9I mseI
 aseI/asnI/vspI
 foki sfaNI mseI
 bstXI
 haeIII/palI
 haeI
 scrFI mvaI
 mvaI ecorII
 dsav tfII dsav
 bstNI nlaIII
 bstNI ddeI pleI
 apyI(dcm+) hinfI apyI(dcm+) hinfI
 accI nlaIII
 mnlI
 mspI
 hpaII
 bsaHI

FIG. 48C

```

          hgaI
          hinfI/acyI
          shaII/bseII
scrFI
mvaI      mnlI
ecorII
dsav
bstNI     econI
          apyI(dcm+) mnlI
          bsaJI     balI ddeI
          mnlI
          901 GTGACAGGA TCATGCAGGA ATTGGAAGT GACACGTTT TCCAGAAAT TGATTGGGG AAATATAAAC CTCTCCAGA ATACCCAGGC GTCTCTCTG
             CACTGTTCCT AGTACGTCCT TAACTTTCA CTGTGCAAAA AGGTCTTTA ACTAAACCCC TTTATATTG GAGAGGTCT TATGGGTCCG CAGGAGAGAC

          scrFI
          mvaI
          ecorII
          dsav
          bstNI
          apyI(dcm+)
          sau96I
          avaiI
          asuI
          mnlI      sfaNI      mboII      mnlI      aluI
          1001 AGGTCCAGGA GGAAGAAGC ATCAAGTATA AGTTGAAGT CTACGAGAAG AAAGACTAAC AGGAAGATGC TTTCAGATTC TCTGCTCCC TCTTAJAGCT
             TCCAGGTCTC CTTTTTCCG TAGTTCATAT TCAAACTTCA GATGCTCTC TTTCTGATG TTTCTTACG AAAGTTCAG AGACGAGGG AGGATTTCGA
             *END DHER

          styI
          bsaJI
          sau3AI
          mboI/ndeII(dam-)
          dpnI(dam+)
          dpnII(dam-)
          nlaIII
          styI
          ncoI
          dsaI bamFI
          ppulOI
          nsiI/avaII
          1101 ATGCATTTT ATAGACCAT GGGACTTTT CTGCTTTAG ATCCCTTGG CTTCGTTAGA ACCGAGCTAC AATTAATACA TAACCTTATG TATCAATAC
             TAGTAATAA TATTCTGGTA CCTGTAAAC GACCGAATC TAGGGAACC GAAGCAATCT TCGCTCGATG TTAATTATGT ATTGGAATAC ATAGTATGTG

          sau96I
          avaiI
          asuI
          scrFI
          mvaI
          ecorII

```

FIG. 48D

1201 ATACGATTTA GGTGACACTA TAGATACAT CCACTTTGCC TTTCTCTCCA CAGGTGTCCA CAGGTGCACC CACTGCAGTC TCGGTTCTAT CGATTGAAAT
 TATGCTAAAT CCACTGTAT ATCTATTGTA GGTAACCG ANAGAGAGT GTCCACAGT GAGGTCCAG GTTGACCTGG AGCCAGATA GCTACTTAA
 seq from PRK6G425VH: Cla-AvrII-
 1301 CCACCATGGG ATGCTCATGT ATCATCCTTT TTCTAGTAGC AACTGCACT CGACTACAT CAGAGTTCA GCTAGTCCAG TCTGGCGTG GCTGTGTCCA
 GGTGTACCC TACCAGTACA TAGTAGGAA AGATCATCG TTGAGTTGA GCTCATGTAA GTTCTCAAGT CGATCAGTC AGACGGCAC CGACCCACCT
 1401 GCCAGGGGC TCACTCCGTT TGCTCTGTC AGCTTCTGCG TACTCTCTCT CGAGTCACTA TATCAGTGG GTCCGTACG CCGCGGTAA GGGCTGGAA
 CGGTCCCGG AGTAGGCAA ACAGGACAG TCGAGACCG ATGAGGAGAG GCTCAGTGAT ATACGTACC CAGGCAGTCC GGGGCCAT CCGGACCTT
 14 P G G S L R L S C A A S G Y S F S S H Y M H W V R Q A P G K G L E

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782
 783
 784
 785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796
 797
 798
 799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000

FIG. 48E

```

81  L Q M N S L R A E D T A V Y Y C A R G D Y R Y N G D W F F D V W G
      scfI      petI      bspI      bspHI      cac8I      mnlI      ddeI      drdI      cac8I      ddeI      drdI
      mboI/ndeII[dam-]      dpnI[dam+]      dpnII[dam-]      alwI[dam-]      hphI      maeII      snaBI      bsaAI      haeIII/pali      sau96I      asuI      nruI      thalI      fnuDII/mvnI      bstUI      bsh1236I
1501 TGGTTGGAT ATATTGATCC TTCCNATGGT GAACTACGT ATATCAAAA GTTCNAGGCG CGTTACCTT TATCTGCGA CAATCCAAA AACACAGCAT
      ACCNACCTA TATRACTAGG AGGTTACCA CTTGATGCA TATTAGTTT CAAGTTCCCG GCNAGTGA ATAGAGGCT GTGAGGTTT TTGTCGCTA
47 W V G Y I D P S N G E T T Y N Q K F K G R F T L S R D N S K N T A Y
      hnlI/acyI      ahaII/bsaHI      aatII      bsrI      maeIII      taqI      hphI      mboII      maeII
1601 ACTGCAGAT GAACAGCTG CGTCTGAGG ACAGTCCGT CTATTACTGT GCNAGGGG ATTATCGCTA CAATGGTAC TGGTTCTTCG ACGTCGGGG
      TGCAGCTCTA CTTGTCGAC GCAGACTCC TGTGACGGCA GATATGACA CGTCTCCCG TANTAGGAT GTTACCACTG ACCAAGAGC TGCAGACCCC
      scrFI      mvaI      ecorII      dsav      bstNI      sau96I      haeIII/pali      asuI      fnu4HI      bsoFI      bsp1286      acII      bsaJI
      nlaIV      hglJII      bsp1286      bsp120I      bmyI      banII      asuI      mboII      ecorII      mvaI      scrFI      hglAI/aspHI      bseRI      bsp1286      bsaIHKAI      bmyI      mnlI      bmyI      nspBII      apyI[dcm+]
1701 TCAGGAGACC CTGGTACCG TCTCTCGG CTTCCACGAG GCCCATCGG TCTTCCCTCC GCACCTCC TCCNAGGCA CCTCTGGGG CACAGCGCC
      AGTTCCTGG GACCACTGGC AGAGAGCCG GAGGTGTTT CCGGTAGCC AGNAGGGGA CCGTGGAGG AGGTTCTCGT GGAGACCCC GTGTCGCGG
114 Q G T L V T V S S A S T K G P S V F P L A P S S K S T S G G T A A

```

FIG. 48F

scrFI hgiAI/aepHI hlnPI nlaIV nari kasi hlnII/acyI mspI hpaII
 mvaI bsp1286 bslHKAII bmyI cac8I scrFI hpaII
 ecorII fnu4HI nciI
 daav bstNI bsoFI
 belI apyI(dcm+) acII apaII/enoiI deav scfI
 fnu4HI bsoFI ahaII/bseHI ddei hhai/cfoI nspBII alw44I/enoiI cauli
 bsoFI bmyI
 1801 CTGGGCTGCC TGGTCAAGGA CTACTTCCC GAAACCGTGA CGGTCTCGTG GAACTCAGGC GCCCTGACCA CGCGGTGCA CACCTTCCG CTTGCTCTAC
 GACCCGACCG ACCAGTTCCT GATGAAGGG CTGGCCACT GCCACAGCAC CTGAGTCCG CGGACTGGT CGCCGACGT GTGGAAGGCC CGACAGGATG
 147 L G C L V K D Y F P E P V T V S W N S G A L T S G V H T F P A V L Q

fnu4HI nlaIV hgiAI hlnPI
 bsoFI bmyI
 bsp1286 maeI aluI bsp1286
 maeIII hphI bmyI mnlI bbvI bmyI
 hphI bmyI mnlI bbvI
 1901 AGTCCTCAGG ACTCTACTCC CTCAGCAGCG TGGTCACTGT GCCCTCTAGC AGCTTGGCA CCCAGACCTA CATCTGCAAC GTGAATCACA AGCCGACAA
 TCAGGAGTCC TGAGATGAGG GATCGTCCG ACCACTGACA CGGGAGATCG TCGACCCGT GGTCTGGAT GTAGACGTG CACTAGTGT TCGGTGCTT
 181 S S G L Y S L S S V T V P S S S L G T Q T Y I C N V N H K P S N
 ahdI/eam1105I
 sau96I
 avari
 scrFI
 mvaI
 ecorII
 daav
 bstNI nlaIV mboII
 bseJI bseJI bseJI
 2001 CACCAAGTG GACAGAAG TTGAGCCCA ATCTTGTGAC AACTCTACA CATGCCACC GTGCCAGCA CCTGNACTCC TGGGGGACC GTCACTCTTC
 CTGTTCCAC CTGTTCTTC AACTCGGTT TAGACACTG TTTTCACTGT GTACGGGTG CACGGGTCTG GACTTACG ACCCCCTGG CAGTCAGAG
 214 T K V D K K V E P K S C D K T H T C P P C P A P E L L C G P S V F

FIG. 48G

FIG. 48H

[illegible]

FIG. 48I

[illegible]

FIG. 48J

3501 CCACCTTTTC TTTTCTCCA CAGGCTCCA CTCCCAGGTC CAACTGACCC TGGTTCGGC AAGCTAGCTT GGGCTGCATC GATTGAATTC CACCATGGGA
GGTGAAGAAG AAAAGAGGT GTCCACAGGT GAGGTCCAG GTTGACGTGC ACCAAGCGC TTGCATCGAA CCCGACCTAG CTAACCTAAG GTGCTACCTT

3701 ATAGGCTCAC CATCACTGC AGGCTAAGTC AAGCTTAGT ACATGGTATA GGTGCTACGT ATTTACACTG GTATCAACAG AAACAGGAA AGCTCCGAA
TATCCAGTG GTAGTGACG TCCAGTTCAG TTTCGNATCA TGTACCTAT CCACGNTGCA TAAATGTGAC CATAGTTGC TTTGCTCCTT TTCGAGGCTT

18 R V T I T C R S S Q

3601 TGGTCATGTA TCATCCTTTT TCTAGTAGCA ACTGCAACTG GAGTACATTC AGATATCCAG TCTATAGGTC TACTGGGTCA GGGGCTCAG GGACAGCGG AGACACCGC
ACCAGTACAT AGTAGGAAA AGATCATCGT TGACGTTGAC CTCATGTNAG D I Q M T Q S P S L S A S V G D

1

FIG. 48M

FIG. 48N

[illegible]

sau3AI
 mboI/ndeII(dam-)
 dpnI(dam+)
 dpnII(dam-)
 pvuI/bspCI
 mcrI
 bslEI
 taqI(dam-)
 clal/bsp106(dam-)
 bspDI(dam-) tru9I
 sau3AI maeI
 mboI/ndeII(dam-)
 dpnI(dam+) xmnI
 dpnII(dam-) hlnPI dsal haeIII/palI
 dpnII(dam-) asel/asnI/vspI bsaJI
 nlaIII alwI(dam-) asp700 hhaI/cfoI nlaIII
 aluI
 pvuII
 nspBII
 rsal
 csp6I
 nlaIV
 kpnI
 hgiCI
 bani
 asp718 mnII
 acc65I ddeI acII
 mnII
 mnlI
 bsmI bfaI
 rmaI
 maeI
 bsmI
 bfaI
 4401 TTTTCACTG CATCTAGTT GTGGTTGTC CAAACTCATC ANTGTATCTT ATCATGTCTG GATCGATCGG GAATTAATTC GCGCAGCAC CATGGCCTGA
 AAAAGTGAC GTAGATCAA CACCAACAG GTTGAGTAG TTACATAGTA TAGTACAGAC CTAGCTAGCC CTTTATTAG CCGCGTCTG GTACCGGACT
 ~sv40 origin
 scrFI
 mvaI
 ecorII
 dsav
 bstNI
 apyI(dcm+)
 bsaJI
 bsmFI nlaIV
 4501 AATAACCTCT GAAGAGGAA CTTGGTTAGG TACCTTCTGA GCGGGAAGA ACCAGCTGTG GAATGTGTGT CAGTTAGGT GTGGAACTC CCCAGGCTCC
 TTATTGGACA CTTTCTCCTT GAACCAATCC ATGGAAGACT CCGCCTTTCT TGGTCGACAC CTACACACA GTCAATCCCA CACCTTTCAG GGGTCCGAGG
 scrFI
 mvaI
 ecorII
 dsav
 bstNI
 apyI(dcm+)
 bsaJI
 bsmFI nlaIV
 4601 CCAGCAGGCA GAAGTATGCA AAGCATGCAT CTCATTAGT CAGCAACCGG GTGTGGAAAG TCCCGAGGCT CCCAGCAGG CAGAGTATG CAAGCATGC
 GGTGTCCTCGT CTTTCATACGT TTCGTACGTA GAGTTAATCA GTCGTTGTC CACACCTTC AGGGTCCGA GGGGTCTGTC GTCTTCATAC GTTTCGTACG
 scrFI
 mvaI
 ecorII
 dsav
 bstNI
 apyI(dcm+)
 bsaJI
 bsmFI nlaIV
 4601 CCAGCAGGCA GAAGTATGCA AAGCATGCAT CTCATTAGT CAGCAACCGG GTGTGGAAAG TCCCGAGGCT CCCAGCAGG CAGAGTATG CAAGCATGC
 GGTGTCCTCGT CTTTCATACGT TTCGTACGTA GAGTTAATCA GTCGTTGTC CACACCTTC AGGGTCCGA GGGGTCTGTC GTCTTCATAC GTTTCGTACG

FIG. 48P

4701 ATCTCATTTA GTCAGCAACC ATAGTCCGCG CCTACTCCG GCCCATCCG CCGGTAACTC GCGCCAGTTC CGCCCATTTCT CCGCCCATG GCTGACTAAT
TAGAGTTANT CAGTCGTTGG TATCAGGCGC GGGATTGAGC CCGGTAGGCG GGGGATTGNG GCGGCTCAG GCGGGTACG GCGGGGTAC CGACTGATTA

4801 TTTTITTTAT TATGCAGAGC CCGAGGCGCG CTCGGCTCT GAGCTATTCC AGNAGTAGTG AGGAGGCTTT TTGGAGGCC TAGGCTTTTG CAAAGAGCTG
AAAAAATAA ATACGTCCTC GGCTCCGGCG GAGCCGGAGA CTCGATAGG TCTTCATCAC TCCTCCGAAA AACCTCCGG ATCCGAAAC GTTTTCGAC
start pUC118

4901 TTACTCTCAG CGGCCGCTTA ATTAAGGCGC GCCATTTTAA TCCTGCAGGT AACAGCTTGG CACTGGCCGT CGTTTACAA CGTCTGACT GGGAAACCC
ANTGAGCTC GCCGGGAAT TAATCCGCG CGGTAAATTT AGGAGTCCA TTGTCGNAAC GTGACCGCA GCAAAATGTT GCAGCACTGA CCCTTTTGGG
linearization linker inserted into HpaI site

FIG. 48Q

mepI nlaIV hgiII hgiII nlaIV hgiII taqI
 hpaII hpaII hpaII hpaII hpaII hpaII
 naeI bmyI bmyI bmyI bmyI bmyI
 cfr101/bsrFI aluI aluI aluI aluI aluI
 maeII cac8I maeII cac8I maeII cac8I maeII cac8I maeII cac8I
 5301 TTCTCGCCA CGTTCGCCG CTTTCCCGT CAAGCTCTAA ATCGGGGGCT CCCTTTAGG TTCCGATTGA GTGCTTTACG GCACCTCGAC CCCAAAAAC
 AAGAGCGGT GCAAGCGCC GNAAGGGCA GTTCGAGATT TAGCCCCCGA GGGAAATCCC AAGCTAAT CACGAATGC CGTGAGCTG GGGTTTTG
 hphI hphI hphI hphI hphI hphI
 5401 TTGATTTGG TGATGTTCA CGTAGTGGC CATCGCCCTG ATAGACGGT TTTCGCCCTT TGACGTTGA GTCACGTTT TTAATAGT GACTCTTGT
 AACTAAACCC ACTACCAAGT GCATCACCCG GTAGCGGAC TATCTGCCNA AANGCGGNA ACTGCNACCT CAGGTSCAG AAATTATCAC CTGAGAACAA
 bsrI bsrI bsrI bsrI bsrI bsrI
 5501 CCAACTGGA ACAACACTCA ACCCTATCTC GGGCTATTCT TTGATTAT AAGGGATTTC GCCGATTTCG GCCTATTGTT TAAAAATGA GCTGATTAA
 GGTGACCT TGTGTGAGT TGGATAGAG CCCGATNAGA AACTAATA TTCCCTNAA CGGCTAAAGC CGGATAACCA ATTTTACT CGACTAAAT
 hgiII/aspHI hgiII/aspHI hgiII/aspHI hgiII/aspHI hgiII/aspHI hgiII/aspHI
 bsp1286 bsp1286 bsp1286 bsp1286 bsp1286 bsp1286
 bslIKAI bslIKAI bslIKAI bslIKAI bslIKAI bslIKAI
 bmyI ddel bmyI ddel bmyI ddel bmyI ddel bmyI ddel
 apaLI/enoI rsaI apaLI/enoI rsaI apaLI/enoI rsaI apaLI/enoI rsaI
 alw44I/enoI csp6I alw44I/enoI csp6I alw44I/enoI csp6I alw44I/enoI csp6I
 5601 CAAAATTTA ACGCAATTT TAACAAATA TTAACGTTA CAATTTTATG GTGCACTCTC AGTACAACTCT GCTCTGATGC CGCATAGTTA AGCCAACCTC
 GTTTTTAAAT TCGCGCTTAA ATTGTTTTAT AATTGCAAT GTTAAANTAC CACGTGAGAG TCATGTTAGA CGAGACTAGC CGGTATCAAT TCGGTGAGG
 hlnPI hlnPI hlnPI hlnPI hlnPI hlnPI
 hhaI/cfoI hhaI/cfoI hhaI/cfoI hhaI/cfoI hhaI/cfoI hhaI/cfoI
 thal thal thal thal thal thal
 fnuDII/mvni fnuDII/mvni fnuDII/mvni fnuDII/mvni fnuDII/mvni fnuDII/mvni
 bsp1286 bsp1286 bsp1286 bsp1286 bsp1286 bsp1286
 bmyI ddel bmyI ddel bmyI ddel bmyI ddel bmyI ddel
 apaLI/enoI rsaI apaLI/enoI rsaI apaLI/enoI rsaI apaLI/enoI rsaI
 alw44I/enoI csp6I alw44I/enoI csp6I alw44I/enoI csp6I alw44I/enoI csp6I
 5701 GCTATCGCTA CGTACCTGGG TCATGCTGC GCCCGACAC CGCCNACAC CGGCTGACGC GCGCTGACGC GCTTGTCTGC TCCCGGCATC CGCTTACAGA
 CGATAGCGAT GCACTGACCC AGTACCGAG CGGGCTGTG GCGGTTGTG GCGACTGCG GCGGACTGCC CGAACAGAG AGGCCGCTAG CGGATGTCT

FIG. 48S

[illegible]

5201 GAGGAGTGGG TTACATCGAA CTGGATCTCA ACAGCGGTAA GATCCTTGAG AGTTTCGCC CCGNAGNACG TTTTCCATNG ATGAGCACCTT TTAAAGTTCT
GTCCTCACCC AATGTAGCTT GACCTAGAGT TGTGCCCAT TGTGCCCAT CTAGGNACTC TCNANAGCGG GGCCTCTTGC AAAGGTTAC TACTCGTGAA AATTTCAGAA
bsfI maeIII taqI alwI[dam-] acII bstYI/xhoII
sau3AI nspBI mboI/ndeII[dam-] dpnI[dam+] dpnII[dam-] alwI[dam-] bstYI/xhoII
bsrI dpnII[dam-] alwI[dam-] bstYI/xhoII
maeII p  p1406I xnnI asp700 mboII
hglAI/aspHI bspI286 tru9I bslHKA I mseI bmyI ahaIII/draI

5301 GCTATGTGGC GCGGTATTAT CCGGTGATGA CCGCGGGCAA GAGCAACTCG GTCGCCGCAT ACACATNTCT CAGAAATGACT TGGTTGAGTA CTCACCAAGTC
GGATACACCG CGCCATTAATA GGGCACTACT GCGGCCCGT CTCGTTGAGC CAGCGGGGTA TGTGATNAGA GTCTTACTGA ACCNACTCAT GAGTGGTCAG
rsal csp6I bsrI scaI hphI maeIII
acII fnu4HI mcrI bseI bsofI ddeI
hnlI/acyI hgaI caulI bcoI bseI bsofI
ahaII/bsaHI hhaI/cfoI

5401 ACAGAAAGC ATCTTACGGA TGGCATGACA GTAAGAGAAAT TATGCAGTGC TGGCATACC ATGATGATA ACACGTGGGC CAACTTACTT CTGACAACGA
TGCTTTTCG TAGAATGCCT ACCGACTGT CATCTCTTA ATACGTCAGC ACGTATTGG TACTCACTAT TGTGACGCCG GTTGAATGAA GACTGTTGCT
sfanI foki nlaIII
fnu4HI bsofI bbvI mslI nlaIII
haeIII/palI eaeI cfrI fnu4HI bsofI acII
sau3AI mboI/ndeII[dam-] dpnI[dam+] dpnII[dam-] pvuI/bspCI mcrI bseI

5501 TCGGAGGACC GAAGGAGCTA ACGGCTTTT TGCACAACAT GGGGATCAT GTACTCGCC CATTCGCTGTA CCCCCTAGTA CATTGAGCGG AACTAGCAAC CCTTGGCCTC GACTTACTTC GGTATGTTTT
AGCCTCCTGG CTTCCTCGAT TGCGGNAAAA
sau96I avaiI asuI mnlI
sau3AI maeIII
mboI/ndeII[dam-] dpnI[dam+] dpnII[dam-] alwI[dam-]
nlaIII
mspI nlaIV
sau3AI nlaIV
mboI/ndeII[dam-] aluI
dpnI[dam+] hpaII
dpnII[dam-] bsaVI

FIG. 48U

hinPI mspI
 hhai/cfoI hpaII
 mstI scrFI
 avIII/fspI bsrI tru9I
 maeII pspl406I maeI dsav mseI
 cac8I bsrDI bsaI asei/asnl/vspI
 maeIII sfaNI bsvI
 6601 CGACGAGCGT GACACCAGCA TGGCAGCAGC AATGGCAACA ACGTTGGCA AACTATTAC TGGCGNACTA CTTACTCTAG CTTCGCCGA ACAATTATA
 GCTGCTCGCA CTGTGGTGT ACGGTGCTG TTACCGTTGT TGCNACGCGT TGCNACGCGT TGCNACGCGT TGCNACGCGT TGCNACGCGT TGCNACGCGT
 bglI bspI
 sau96I hpaII
 haeIII/palI cfr10I/bsrFI
 sau96I hpaII
 avall mspI
 hinPI hpaII
 hhai/cfoI hpaII
 sau96I bsrDI bsaI
 bsh1236I bsrDI bsaI
 6701 GACTGGATGG AGCGGATAA AGTTGAGCA CCACTTCTGC GCTGGCCCT TCCGGCTGGC TGGTTATTG CTGATANAATC TGGAGCGGT GAGCGTGGGT
 CTGACCTACC TCGGCTATT TCNACGCTCT GTGAAGAGC CGAGCCGGA AGCGGACCG ACCANNTAC GACTATTAG ACCTCGCCA CTGCGACCCA
 acli
 fnu4HI haeIII/palI
 fnuDII/mvni bsoFI sau96I
 batUI bsvI nlaIV
 bsh1236I bsrDI bsaI
 6801 CTGCGGTAT CATTGAGCA CTGGGCCAG ATGGTAAGC CTCCGCTATC GTAGTTATCT ACACGAGCGG GAGTACGGA ACTATGGATG AACGAATAG
 GAGGCCATA GTAACGCTCT GACCCGCTC TACCATTGCG GAGGCTAG CATCATAGA TGTGCTGCC CTCAGTCCCT TGATACCTAC TTGCTTTATC
 ddeI
 sau3AI nlaIV
 mboI/ndeII(dam-) hglCI tru9I
 dpnII(dam+) hglCI maeII
 dpnII(dam-) bsnI mnlI maeII
 6901 ACAGATGCT GAGATAGTG CCTACTGAT TAAGCATGG TAACTGTGAG ACCAGTTTA CTCATATATA CTTAGATTG ATTTAAACT TCATTTTAA
 TGCTAGGCA CTCTATCCAC GGAGTACTA ATTCGTAACC ATTGACAGTC TGGTTCAAT GAGTATATAT GAAATCTAC TAAATTTGA AGTAAATTT
 rmaI sau3AI
 maeI mboI/ndeII(dam-) tru9I
 sau3AI hpaI mboI/ndeII(dam-) maeI
 mboI/ndeII(dam-) dpnII(dam+) hglCI
 dpnII(dam-) dpnII(dam-) dpnII(dam-) hglCI
 tru9I bsnI/xhoII alwi(dam-) batYI/xhoII
 maeI alwi(dam-) bsnI mboI/ndeII(dam-) sau3AI
 haeIII/draI bsaI mboI/ndeII(dam-) dpnII(dam+) maeI
 7001 TTTAAAGGA TCTAGGTGAA GATCCCTTTT GATATCTCA TGACCAAAAT CCCTTACGT GAGTTTCTG TCCACTGAGC GTCAGACCCC GTAGAAAGA
 AAATTTCTCT AGATCCACTT CTAGGAAA CTATTAGAT ACTGCTTTA GGGATTGCA CTCMAAGCA AGGTGACTCG CAGTCTGGG CATCTTTCT

FIG. 48V

FIG. 48W


```

scrFI      mvaI      ecorII      deav      bstNI      bsaJI      bsaSI      hinPI      mnlI      hhaI/cfoI      aluI apyI(dcm+)
7501 TGAGCATGGA GAAAGCGCA CGCTTCCCGA AGGAGAAAG GCGACAGGT ATCCGTAAG CGCAGGTC GGAACAGGAG AGCGACGAG GGAGCTTCCA
ACTCGTAAT CTTCCGGGT GGAAGGGCT TCCCTCTTC 'CGCTGTCCA TAGGCATTC GCGTCCCAG CCTGTCTC TCGGTCTC CCTCGAAGGT

scrFI      mvaI      ecorII      deav      bstNI      apyI(dcm+)      nlaIV      acII
7601 GGGGAAAG CCTGTATCT TTATATCTCT GTCGGGTTT GCCACTCTG ACTTGAGGT CGATTTTGT GATGCTGTC AGGGGCGCG AGCTATGGA
CCCCCTTTC GGACCATAGA AATATCAGGA CAGCCCAAG CGGTGGAGAC TGAATCCCA GCTAAACA CTACGAGCAG TCCCCCGCC TCGATACCT

haeIII/palI      haeIII/palI      scrFI      mvaI      bsaI      ecorII      dsaV      bstNI      nlaIII      haeIII/palI      nspI      aflIII
7701 AAAAGCGCAG CAACGGGCG TTTTACGGT TCTGGCCCT TTGCTGCGCT TTGCTGCGCT TTTGCTGCGCT TTTGCTGCGCT TTTGCTGCGCT TTTGCTGCGCT TTTGCTGCGCT TTTGCTGCGCT
TTTTGGGTC GTTGGGCGG AAAATGCCA AGACCGGAA AACGACCGA AACGAGTGT ACAAGAAAG ACCTAAGAG ACCTAATGGC

cacBI      bsh1236I      nlaIV      haeI      cacBI      haeI      haeIII/palI      nspI      aflIII      tfII      hinFI
7801 TATTACGCG TTTAGTGAG CTGATACCG TCGCGGCGC GGAACGCGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC
ATAATGGCG AACTCACTC GACTATGCG AGCGGCGTC GCTGTGTC TCGGTGCT CAGTCACTC TCTCTGCG TCTCTGCG TCTCTGCG TCTCTGCG TCTCTGCG

scrFI      mvaI      ecorII      deav      bstNI      apyI(dcm+)      nlaIV      acII      hinPI      mnlI      hhaI/cfoI      earI/asp632I      haeII
7801 TATTACGCG TTTAGTGAG CTGATACCG TCGCGGCGC GGAACGCGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC GTGAGTGGC
ATAATGGCG AACTCACTC GACTATGCG AGCGGCGTC GCTGTGTC TCGGTGCT CAGTCACTC TCTCTGCG TCTCTGCG TCTCTGCG TCTCTGCG TCTCTGCG

```

FIG. 48X


```

tru9I
mseI
aseI/aseI/vspI
xmni
nleIII asp700
8101 TGACCATGAT TACGAATTAA (SEQ ID NO:68)
ACTGGTACTA ATGCTTAATT

>length: 8120

aatII(GACGTC): 1690 5947
acc65I(GGTACC): 2969 3967 4529
823 1039 2738 4237
217 229 238 250 260 271 317 422 454 485 574 1385 1795 1871 2248 2250 2758 2982
3167 3179 3188 3200 3210 3221 3267 3372 3404 3449 3686 3949 4021 4318 4542 4727
4739 4748 4760 4770 4781 4827 4910 4914 5070 5127 5153 5166 5203 5217 5220 5248
5275 5680 5699 5741 5751 5790 5979 6026 6125 6234 6311 6355 6476 6522 6713 6804
7166 7175 7310 7420 7541 7560 7687 7715 7806 7827 7834 7877 7901 7911 7967 8070
see hlnI
acyI
afIII/bfI(CTTAAG): 786
afIII(ACRYGT): 932 7758
ageI(ACCGGT): 1833
ahaII/baaHI(GRCGYC): 988 1690 1858 5117 5947 6329
ahaII/draI(TTTAA): 696 4935 6290 6982 7001
ahdI/eam1105I(GACNNNNNGTC): 2087 6865
5 44 332 386 390 753 1097 1165 1370 1431 1951 2603 2751 2784 3282 3336 3340
3562 3566 3676 3733 3792 4270 4288 4311 4344 4554 4842 4896 4954 5047 5333 5590
5803 5822 6516 6579 6679 7200 7457 7593 7819 7937 8096
aluI(AGCT):
alw44I/anoI(GTGCAC): 1876 5651 6198 7444

```

FIG. 48Z




FIG. 49A




FIG. 49B




FIG. 49C




FIG. 49D




FIG. 50A





FIG. 50B



Representative Conc versus Time Plot. Shown is the kinetic data for 6G4V11N35A.IgG1

SAMPLE	k_a	k_d	K_d
Murine 6G4.2.5 IgG2a	8.3×10^5	2.9×10^{-4}	350pM
6G4V11N35A-IgG1	8.7×10^5	7.7×10^{-5}	88pM
6G4V11N35E-IgG1	3.0×10^6	1.4×10^{-4}	49pM

FIG. 51




FIG. 52A




FIG. 52B

781 AAAAGGGTAT CTAGAGGTTG AGGTGATTTT ATGAAAAAGA ATATCGCATT TCTTCTTGCA
 TTTTCCCATA GATCTCCAAC TCCACTAAAA TACTTTTCT TATAGCGTAA AGAAGAACGT
 -1 M K K N I A F L L A

841 TCTATGTTCTG TTTTCTCTAT TGCTACAAAC GCGTACGCTG AGGTTTCAGCT AGTGCAGTCT
 AGATACAAGC AAAAAAGATA ACGATGTTTG CGCATGCGAC TCCAAGTCGA TCACGTCAGA
 -11 S M F V F S I A T N A Y A E V Q L V Q S

901 GGCGGTGGCC TGGTGCAGCC AGGGGGCTCA CTCCGTTTGT CCTGTGCAGC TTCTGGCTAC
 CCGCCACCGG ACCACGTCGG TCCCCGAGT GAGGCAAACA GGACACGTCG AAGACCGATG
 8 G G G L V Q P G G S L R L S C A A S G Y

961 TCCTTCTCGA GTCACTATAT GCACTGGGTC CGTCAGGCCC CGGGTAAGGG CCTGGAATGG
 AGGAAGAGCT CAGTGATATA CGTGACCCAG GCAGTCCGGG GCCCATTCCTC GGACCTTACC
 28 S F S S H Y M H W V R Q A P G K G L E W

1021 GTTGATATA TTGATCCTTC CAATGGTGAA ACTACGTATA ATCAAAAGTT CAAGGGCCGT
 CAACCTATAT AACTAGGAAG GTTACCACTT TGATGCATAT TAGTTTTCAA GTTCCCGGCA
 48 V G Y I D P S N G E T T Y N O K F K G R

1081 TTCACTTTAT CTCGCGACAA CTCCAAAAAC ACAGCATACC TGCAGATGAA CAGCCTGCGT
 AAGTGAATA GAGCGCTGTT GAGGTTTTTG TGTCGTATGG ACGTCTACTT GTCGGACGCA
 68 F T L S R D N S K N T A Y L Q M N S L R

1141 GCTGAGGACA CTGCCGTCTA TTACTGTGCA AGAGGGGATT ATCGCTACAA TGGTGACTGG
 CGACTCCTGT GACGGCAGAT AATGACACGT TCTCCCTTAA TAGCGATGTT ACCACTGACC
 88 A E D T A V Y Y C A R G D Y R Y N G D W

1201 TTCTTCGACG TCTGGGGTCA AGGAACCCTG GTCACCGTCT CCTCGGCCTC CACCAAGGGC
 AAGAAGCTGC AGACCCCACT TCCTTGGGAC CAGTGGCAGA GGAGCCGGAG GTGGTTCCCG
 108 F F D V W G Q G T L V T V S S A S T K G

1261 CCATCGGTCT TCCCCCTGGC ACCCTCCTCC AAGAGCACCT CTGGGGGCAC AGCGGCCCTG
 GGTAGCCAGA AGGGGGACCG TGGGAGGAGG TTCTCGTGGA GACCCCGTG TCGCCGGGAC
 128 P S V F P L A P S S K S T S G G T A A L

1321 GGCTGCCTGG TCAAGGACTA CTTCCCCGAA CCGGTGACGG TGTCGTGGAA CTCAGGCGCC
 CCGACGGACC AGTTCCTGAT GAAGGGGCTT GGCCACTGCC ACAGCACCTT GAGTCCGCGG
 148 G C L V K D Y F P E P V T V S W N S G A

1381 CTGACCAGCG GCGTGCACAC CTTCCCGGCT GTCCTACAGT CCTCAGGACT CTACTCCCTC
 GACTGGTTCG CGCACGTGTG GAAGGGCCGA CAGGATGTCA GGAGTCCTGA GATGAGGGAG
 168 L T S G V H T F P A V L Q S S G L Y S L

1441 AGCAGCGTGG TGACCGTGCC CTCCAGCAGC TTGGGCACCC AGACCTACAT CTGCAACGTG
 TCGTCGCACC ACTGGCACGG GAGGTGCTCG AACCCGTGGG TCTGGATGTA GACGTTGCAC
 188 S S V V T V P S S S L G T Q T Y I C N V

1501 AATCACAAGC CCAGCAACAC CAAGGTGCGAC AAGAAAGTTG AGCCCAAATC TTGTGACAAA
 TTAGTGTTCTG GGTGCTGTG GTTCCAGCTG TTCTTTCAAC TCGGGTTTAG AACACTGTTT
 208 N H K P S N T K V D K K V E P K S C D K

1561 ACTCACACAT GCCCGCCGTGA (SEQ ID NO: 69)
 TGAGTGTGTA CCGGCGGCACT
 228 T H T C P P O
 (SEQ ID NO: 70)

FIG. 53






FIG. 54A



Antibody Competitor (nM)

FIG. 54B



Antibody Competitor (nM)

FIG. 54C




FIG. 55A




FIG. 55B





FIG. 55C






FIG. 56B




FIG. 56C


Pegylated F(ab')₂ (nM)

FIG. 57A


Pegylated F(ab')₂ (nM)

FIG. 57B




FIG. 58A




FIG. 58B




FIG. 59A





FIG. 59B






FIG. 61A




FIG. 61B




FIG. 62




FIG. 63






FIG. 64






FIG. 66




FIG. 67




FIG. 68




FIG. 69




FIG. 70A




FIG. 70B




FIG. 70C




FIG. 70D





FIG. 70E

The Effect of Pegylated Anti-IL-8 in the Rabbit Ear model of Ischemia-Reperfusion Injury



Anti-IL-8 formulations :
Single Dose (5 mg/kg)
administered IV at time
of reperfusion

FIG. 71

Sequence Listing

<110> Genentech, Inc.,
Hsei, Vanessa
Koumenis, Iphigenia
5 Leong, Steven R.
Presta, Leonard G.
Shahrokh, Zahra
Zapata, Gerardo A.

<120> ANTIBODY FRAGMENT-POLYMER CONJUGATES AND HUMANIZED
10 ANTI-IL-8 MONOCLONAL ANTIBODIES AND USES OF SAME

<130> P1085R5PCT

<150> US 09/012,116
<151> 1998-01-22

<150> PCT/US98/03337
15 <151> 1998-02-20

<150> US 09/121,952
<151> 1998-07-24

<150> US 09/122,513
<151> 1998-07-24

20 <160> 72

<210> 1
<211> 22
<212> DNA
<213> Mus musculus

25 <400> 1
cagtccaact gttcaggacg cc 22

<210> 2
<211> 22
<212> DNA
30 <213> Mus musculus

<400> 2
gtgctgctca tgctgtaggt gc 22

<210> 3
<211> 23
35 <212> DNA
<213> Mus musculus

<400> 3
gaagttgatg tcttgtgagt ggc 23

<210> 4
40 <211> 24

<212> DNA
<213> Mus musculus

<400> 4
gcacccctaga gtcaccgagg agcc 24

5 <210> 5
<211> 22
<212> DNA
<213> Mus musculus

<400> 5
10 cactggctca gggaaataac cc 22

<210> 6
<211> 22
<212> DNA
<213> Mus musculus

15 <400> 6
ggagagctgg gaaggtgtgc ac 22

<210> 7
<211> 35
<212> DNA
20 <213> Mus musculus

<400> 7
acaaacgcgt acgctgacat cgatcatgacc cagtc 35

<210> 8
<211> 35
25 <212> DNA
<213> Mus musculus

<400> 8
acaaacgcgt acgctgatat tgatcatgact cagtc 35

<210> 9
30 <211> 35
<212> DNA
<213> Mus musculus

<400> 9
acaaacgcgt acgctgacat cgatcatgaca cagtc 35

35 <210> 10
<211> 37
<212> DNA
<213> Mus musculus

<400> 10
40 gctcttcgaa tgggtgggaag atggatacag ttggtgc 37

<210> 11
<211> 39

<212> DNA
<213> Mus musculus

<400> 11
cgatgggccc ggatagaccg atggggctgt tgttttggc 39

5 <210> 12
<211> 39
<212> DNA
<213> Mus musculus

<400> 12
10 cgatgggccc ggatagactg atggggctgt cgttttggc 39

<210> 13
<211> 39
<212> DNA
<213> Mus musculus

15 <400> 13
cgatgggccc ggatagacgg atggggctgt tgttttggc 39

<210> 14
<211> 39
<212> DNA
20 <213> Mus musculus

<400> 14
cgatgggccc ggatagacag atggggctgt tgttttggc 39

<210> 15
<211> 39
25 <212> DNA
<213> Mus musculus

<400> 15
cgatgggccc ggatagactg atggggctgt tgttttggc 39

<210> 16
30 <211> 369
<212> DNA
<213> Mus musculus

<400> 16
gacattgtca tgacacagtc tcaaaaattc atgtccacat cagtaggaga 50

35 cagggtcagc gtcacctgca aggccagtca gaatgtgggt actaatgtag 100

cctggatatca acagaaacca gggcaatctc ctaaagcact gatttactcg 150

tcatectacc ggtacagtgg agtcctgat cgcttcacag gcagtggatc 200

tgggacagat ttcaactctca ccatcagcca tgtgcagtct gaagacttgg 250

cagactattt ctgtcagcaa tataacatct atcctctcac gttcggctct 300

gggaccaagc tggagttgaa acgggctgat gctgcaccac caactgtatc 350

catcttccca ccattcgaa 369

<210> 17

<211> 123

5 <212> PRT

<213> Mus musculus

<400> 17

	Asp	Ile	Val	Met	Thr	Gln	Ser	Gln	Lys	Phe	Met	Ser	Thr	Ser	Val	
	1				5					10					15	
10	Gly	Asp	Arg	Val	Ser	Val	Thr	Cys	Lys	Ala	Ser	Gln	Asn	Val	Gly	
					20					25					30	
	Thr	Asn	Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	
					35					40					45	
	Ala	Leu	Ile	Tyr	Ser	Ser	Ser	Tyr	Arg	Tyr	Ser	Gly	Val	Pro	Asp	
15					50					55					60	
	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	
					65					70					75	
	Ser	His	Val	Gln	Ser	Glu	Asp	Leu	Ala	Asp	Tyr	Phe	Cys	Gln	Gln	
					80					85					90	
20	Tyr	Asn	Ile	Tyr	Pro	Leu	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Leu	Glu	
					95					100					105	
	Leu	Lys	Arg	Ala	Asp	Ala	Ala	Pro	Pro	Thr	Val	Ser	Ile	Phe	Pro	
					110					115					120	
	Pro	Phe	Glu													
25																

<210> 18

<211> 417

<212> DNA

<213> Mus musculus

30 <400> 18

ttctattgct acaaacgcgt acgctgaggt gcagctggtg gactctgggg 50

gaggcttagt gccgcctgga gggccctga aactctcctg tgcagcctct 100

ggattcatat tcagtagtta tggcatgtct tgggttcgcc agactccagg 150

caagagcctg gaggtagtgc caaccattaa taataatggt gatagcacct 200

35 attatccaga cagtgtgaag ggccgattca ccattctccg agacaatgcc 250

agaacacccc tgtacctgca aatgagcagt ctgaagtctg aggacacagc 300

catgttttac tgtgcaagag cctcattag ttcggctact tgggttggtt 350

actggggcca agggactctg gtcactgtct ctgcagccaa aacaacagcc 400

ccatctgtct atccggg 417

<210> 19

<211> 130

5 <212> PRT

<213> Mus musculus

<400> 19

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Pro Pro Gly
1 5 10 15

10 Gly Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ile Phe Ser
20 25 30

Ser Tyr Gly Met Ser Trp Val Arg Gln Thr Pro Gly Lys Ser Leu
35 40 45

15 Glu Leu Val Ala Thr Ile Asn Asn Asn Gly Asp Ser Thr Tyr Tyr
50 55 60

Pro Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala
65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Ser Ser Leu Lys Ser Glu Asp
80 85 90

20 Thr Ala Met Phe Tyr Cys Ala Arg Ala Leu Ile Ser Ser Ala Thr
95 100 105

Trp Phe Gly Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
110 115 120

25 Ala Lys Thr Thr Ala Pro Ser Val Tyr Pro
125 130

<210> 20

<211> 31

<212> DNA

<213> Artificial Sequence

30 <220>

<221> Artificial Sequence

<222> 1-31

<223> recombinant immunoglobulin

<400> 20

35 acaaacgcgt acgctgatat cgatcatgaca g 31

<210> 21

<211> 31

<212> DNA

<213> Artificial Sequence

40 <220>

<221> Artificial Sequence
 <222> 1-31
 <223> recombinant immunoglobulin

<400> 21
 5 gcagcatcag ctcttcgaag ctccagcttg g 31

<210> 22
 <211> 21
 <212> DNA
 <213> Escherichia coli

10 <400> 22
 ccactagtac gcaagttcac g 21

<210> 23
 <211> 33
 <212> DNA
 15 <213> Artificial Sequence

<220>
 <221> Artificial Sequence
 <222> 1-33
 <223> recombinantimmunoglobulin

20 <400> 23
 gatggggccct tgggtggaggc tgcagagaca gtg 33

<210> 24
~~<211> 714~~
 <212> DNA
 25 <213> Artificial Sequence

<220>
 <221> Artificial Sequence
 <222> 1-714
 <223> recombinant immunoglobulin

30 <400> 24
 atgaagaaga atatgcatt tcttcttgca tctatgttcg ttttttctat 50
 tgctacaaac gcgtagctg atategtcat gacacagtct caaaaattca 100
 tgtccacatc agtaggagac agggtcagcg tcacctgcaa ggccagtcag 150
 aatgtgggta ctaatgtagc ctgggtatcaa cagaaaccag ggcaatctcc 200
 35 taaagcactg atttactcgt catcctaccg gtacagtgga gtccctgac 250
 gcttcacagg cagtggatct gggacagatt tcactctcac catcagccat 300
 gtgcagtctg aagacttggc agactatttc tgcagcaat ataacatcta 350
 tcctctcagc ttcggtcctg ggaccaagct ggagcttcga agagctgtgg 400
 ctgcaccatc tgtcttcac ttcctgccat ctgatgagca gttgaaatct 450

ggaactgctt ctgttggttg cctgctgaat aacttctatc ccagagagggc 500

caaagtacag tggaaggtgg ataacgcctt ccaatcgggt aactcccagg 550

agagtgtcac agagcaggac agcaaggaca gcacctacag cctcagcagc 600

accctgacgc tgagcaaagc agactacgag aaacacaaag tctacgcctg 650

5 cgaagtcacc catcagggcc tgagctcgcc cgtcacaaag agcttcaaca 700

ggggagagtg ttaa 714

<210> 25

<211> 237

<212> PRT

10 <213> Artificial Sequence

<220>

<221> Artificial Sequence

<222> 1-237

<223> recombinant immunoglobulin

15 <400> 25

Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15

Ser Ile Ala Thr Asn Ala Tyr Ala Asp Ile Val Met Thr Gln Ser
20 25 30

20 Gln Lys Phe Met Ser Thr Ser Val Gly Asp Arg Val Ser Val Thr
35 40 45

Cys Lys Ala Ser Gln Asn Val Gly Thr Asn Val Ala Trp Tyr Gln
50 55 60

25 Gln Lys Pro Gly Gln Ser Pro Lys Ala Leu Ile Tyr Ser Ser Ser
65 70 75

Tyr Arg Tyr Ser Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser
80 85 90

Gly Thr Asp Phe Thr Leu Thr Ile Ser His Val Gln Ser Glu Asp
95 100 105

30 Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Asn Ile Tyr Pro Leu Thr
110 115 120

Phe Gly Pro Gly Thr Lys Leu Glu Leu Arg Arg Ala Val Ala Ala
125 130 135

35 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
140 145 150

Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
155 160 165

Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
170 175 180

Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
185 190 195

5 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
200 205 210

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
215 220 225

10 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
230 235 237

<210> 26

<211> 756

<212> DNA

<213> Artificial Sequence

15 <220>

<221> Artificial Sequence

<222> 1-756

<223> recombinant immunoglobulin

<400> 26

20 atgaaaaaga atatcgcat tcttcttgca tctatgttcg tttttctat 50
tgctacaaac gcgtacgctg aggtgcagct ggtggagtct gggggaggct 100
tagtgccgcc tggagggtcc ctgaaactct cctgtgcagc ctctggattc 150
atattcagta gttatggcat gtcttgggtt cgccagactc caggcaagag 200
cctggagtgt gtcgcaacca ttaataataa tggatagc acctattatc 250
25 cagacagtgt gaagggccga ttcacatct cccgagacaa tgccaagaac 300
accctgtacc tgcaaatgag cagtctgaag tctgaggaca cagccatgtt 350
ttactgtgca agagccctca ttagttcggc tacttggttt ggttactggg 400
gccaaggac tctggtcact gtctctgcag cctccaccaa gggcccatcg 450
gtcttcccc tggcaccctc ctccaagagc acctctgggg gcacagcggc 500
30 cctgggctgc ctggtcaagg actacttccc cgaaccggtg acggtgtcgt 550
ggaactcagg cgccctgacc agcggtgtgc acaccttccc ggctgtccta 600
cagtcctcag gactctactc cctcagcagc gtggtgaccg tgccctccag 650
cagcttgggc acccagacct acatctgcaa cgtgaatcac aagcccagca 700
acaccaaggt ggacaagaaa gttgagccca aatcttgtga caaaactcac 750

acatga 756

<210> 27

<211> 251

<212> PRT

5 <213> Artificial Sequence

<220>

<221> Artificial Sequence

<222> 1-251

<223> recombinant immunoglobulin

10 <400> 27

Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15

Ser Ile Ala Thr Asn Ala Tyr Ala Glu Val Gln Leu Val Glu Ser
20 25 30

15 Gly Gly Gly Leu Val Pro Pro Gly Gly Ser Leu Lys Leu Ser Cys
35 40 45

Ala Ala Ser Gly Phe Ile Phe Ser Ser Tyr Gly Met Ser Trp Val
50 55 60

20 Arg Gln Thr Pro Gly Lys Ser Leu Glu Leu Val Ala Thr Ile Asn
65 70 75

Asn Asn Gly Asp Ser Thr Tyr Tyr Pro Asp Ser Val Lys Gly Arg
80 85 90

Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr Leu Gln
95 100 105

25 Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Phe Tyr Cys Ala
110 115 120

Arg Ala Leu Ile Ser Ser Ala Thr Trp Phe Gly Tyr Trp Gly Gln
125 130 135

30 Gly Thr Leu Val Thr Val Ser Ala Ala Ser Thr Lys Gly Pro Ser
140 145 150

Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
155 160 165

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
170 175 180

35 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
185 190 195

Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser
200 205 210

Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile

[illegible]

```

<210> 28
<211> 37
<212> DNA
<213> Mus musculus

10 <400> 28
    ccaatgcata cgctgacatc gtgatgaccc agacccc 37

    <210> 29
    <211> 37
    <212> DNA
15 <213> Mus musculus

    <400> 29
    ccaatgcata cgctgatatt gtgatgactc agactcc 37

    <210> 30
    <211> 37
20 <212> DNA
    <213> Mus musculus

    <400> 30
    ccaatgcata cgctgacatc gtgatgacac agacacc 37

    <210> 31
25 <211> 35
    <212> DNA
    <213> Mus musculus

    <400> 31
    agatgtcaat tgctcaactgg atggtgggaa gatgg 35

30 <210> 32
    <211> 32
    <212> DNA
    <213> Mus musculus

    <400> 32
35 caaacgcgta cgctgagatc cagctgcagc ag 32

    <210> 33
    <211> 32
    <212> DNA
    <213> Mus musculus

40 <400> 33
    caaacgcgta cgctgagatt cagctccagc ag 32

```

<210> 34
 <211> 391
 <212> DNA
 <213> Mus musculus

5 <400> 34
 gatatcgtga tgacacagac accactctcc ctgcctgtca gtcttggaga 50
 tcaggcctcc atctcttgca gatctagtca gagccttgta cacgggtattg 100
 gaaacaccta ttacattgg tacctgcaga agccaggcca gtctccaaag 150
 ctctgatct acaaagtttc caaccgattt tctgggggtcc cagacagggt 200
 10 cagtggcagt ggatcaggga cagatttcac actcaggatc agcagagtgg 250
 aggctgagga tctgggactt tatttctgct ctcaaagtac acatgttccg 300
 ctcacgttcg gtgctgggac caagctggag ctgaaacggg ctgatgctgc 350
 accaactgta tccatcttcc caccatccag tgagcaattg a 391

<210> 35
 15 <211> 131
 <212> PRT
 <213> Mus musculus

<400> 35
 Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu
 20 1 5 10 15
 Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val
 20 25 30
 His Gly Ile Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro
 35 40 45
 25 Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe
 50 55 60
 Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp
 65 70 75
 Phe Thr Leu Arg Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Leu
 30 80 85 90
 Tyr Phe Cys Ser Gln Ser Thr His Val Pro Leu Thr Phe Gly Ala
 95 100 105
 Gly Thr Lys Leu Glu Leu Lys Arg Ala Asp Ala Ala Pro Thr Val
 110 115 120
 35 Ser Ile Phe Pro Pro Ser Ser Glu Gln Leu Lys
 125 130 131

<210> 36

<211> 405
 <212> DNA
 <213> Mus musculus

<400> 36

5 gagattcagc tgcagcagtc tggacctgag ctgatgaagc ctggggcttc 50
 agtgaagata tcttgcaagg cttctgggta ttcattcagt agccactaca 100
 tgcactgggt gaagcagagc catggaaaga gccttgagt gattggctac 150
 attgatcctt ccaatgggtga aactacttac aaccagaaat tcaagggcaa 200
 ggccacattg actgtagaca catcttcag cacagccaac gtgcatctca 250
 10 gcagcctgac atctgatgac tctgcagtct atttctgtgc aagaggggac 300
 tatagataca acggcgactg gtttttcgat gtctggggcg cagggaccac 350
 ggtcaccgtc tctccgcca aaaccgacag ccccatcggt ctatccgggc 400
 ccatac 405

15 <210> 37
 <211> 135
 <212> PRT
 <213> Mus musculus

<400> 37

20 ~~Glu Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly~~
 1 5 10 15
 Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser
 20 25 30
 Ser His Tyr Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu
 35 40 45
 25 Glu Trp Ile Gly Tyr Ile Asp Pro Ser Asn Gly Glu Thr Thr Tyr
 50 55 60
 Asn Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser
 65 70 75
 30 Ser Ser Thr Ala Asn Val His Leu Ser Ser Leu Thr Ser Asp Asp
 80 85 90
 Ser Ala Val Tyr Phe Cys Ala Arg Gly Asp Tyr Arg Tyr Asn Gly
 95 100 105
 Asp Trp Phe Phe Asp Val Trp Gly Ala Gly Thr Thr Val Thr Val
 110 115 120
 35 Ser Ser Ala Lys Thr Asp Ser Pro Ile Gly Leu Ser Gly Pro Ile
 125 130 135

<210> 38
<211> 22
<212> DNA
<213> Artificial Sequence

5 <220>
<221> Artificial Sequence
<222> 1-22
<223> recombinant immunoglobulin

<400> 38
10 cttggtggag gcggaggaga cg 22

<210> 39
<211> 38
<212> DNA
<213> Artificial Sequence

15 <220>
<221> Artificial Sequence
<222> 1-38
<223> recombinant immunoglobulin

<400> 39
20 gaaacgggct gttgctgcac caactgtatt catcttcc 38

<210> 40
<211> 31
<212> DNA
<213> Artificial Sequence

25 <220>
<221> Artificial Sequence
<222> 1-31
<223> recombinant immunoglobulin

<400> 40
30 gtcacgtct cctccgctc caccaagggc c 31

<210> 41
<211> 729
<212> DNA
<213> Artificial Sequence

35 <220>
<221> Artificial Sequence
<222> 1-729
<223> recombinant immunoglobulin

<400> 41
40 atgaagaaga atatcgatt tcttcttgca tctatgttcg tttttctat 50

tgctacaaat gcatacgctg atatcgatgac gacacagaca ccactctccc 100

tgctgtcag tcttgagat caggcctcca tctcttgag atctagtcag 150

- 14 -

Val Glu Ala Glu Asp Leu Gly Leu Tyr Phe Cys Ser Gln Ser Thr
 110 115 120
 His Val Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 125 130 135
 5 Arg Ala Val Ala Ala Pro Thr Val Phe Ile Phe Pro Pro Ser Ser
 140 145 150
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn
 155 160 165
 10 Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn
 170 175 180
 Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp
 185 190 195
 Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser
 200 205 210
 15 Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr
 215 220 225
 His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly
 230 235 240

20 Glu Cys
 242

<210> 43

<211> 762

<212> DNA

<213> Artificial Sequence

25 <220>

<221> Artificial Sequence

<222> 1-762

<223> recombinant immunoglobulin

<400> 43

30 atgaaaaaga atatcgcat tcttcttgca tctatgttcg ttttttctat 50
 tgctacaaac gcgtacgctg agattcagct gcagcagtct ggacctgagc 100
 tgatgaagcc tggggcttca gtgaagatat cctgcaagcc ttctgggttat 150
 tcattcagta gccactacat gcactgggtg aagcagagcc atggaaagag 200
 ccttgagtgg attggctaca ttgatccttc caatgggtgaa actacttaca 250
 35 accagaaatt caagggcaag gccacattga ctgtagacac atcttccagc 300
 acagccaacg tgcattctcag cagcctgaca tctgatgact ctgcagtcta 350
 tttctgtgca agaggggact atagatacaa cggcgactgg tttttcgatg 400

tctggggcgc agggaccacg gtcaccgtct cctccgcctc caccaagggc 450
 ccacggtct tccccctggc accctcctcc aagagcacct ctgggggcac 500
 agcgccctg ggctgcctgg tcaaggacta cttccccgaa ccggtgacgg 550
 tgctgtggaa ctcaggcgcc ctgaccagcg gcgtgcacac cttcccggt 600
 5 gtctacagt cctcaggact ctactccctc agcagcgtgg tgaccgtgcc 650
 ctccagcagc ttgggcaccc agacctacat ctgcaacgtg aatcacaagc 700
 ccagcaacac caaggtggac aagaaagttg agcccaaata ttgtgacaaa 750
 actcacacat ga 762
 <210> 44
 10 <211> 253
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> Artificial Sequence
 15 <222> 1-253
 <223> recombinant immunoglobulin
 <400> 44
 Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
 1 5 10 15
 20 Ser Ile Ala Thr Asn Ala Tyr Ala Glu Ile Gln Leu Gln Gln Ser
 20 25 30
 Gly Pro Glu Leu Met Lys Pro Gly Ala Ser Val Lys Ile Ser Lys
 35 40 45
 Lys Ala Ser Gly Tyr Ser Phe Ser Ser His Tyr Met His Trp Val
 50 55 60
 25 Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile Gly Tyr Ile Asp
 65 70 75
 Pro Ser Asn Gly Glu Thr Thr Tyr Asn Gln Lys Phe Lys Gly Lys
 80 85 90
 30 Ala Thr Leu Thr Val Asp Thr Ser Ser Ser Thr Ala Asn Val His
 95 100 105
 Leu Ser Ser Leu Thr Ser Asp Asp Ser Ala Val Tyr Phe Cys Ala
 110 115 120
 Arg Gly Asp Tyr Arg Tyr Asn Gly Asp Trp Phe Phe Asp Val Trp
 125 130 135
 35 Gly Ala Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly
 140 145 150

	Pro Ser Val Phe	Pro Leu Ala	Pro Ser Ser Lys	Ser Thr Ser Gly	
	155		160	165	
	Gly Thr Ala Ala	Leu Gly Cys Leu	Val Lys Asp Tyr	Phe Pro Glu	
	170		175	180	
5	Pro Val Thr Val	Ser Trp Asn Ser	Gly Ala Leu Thr	Ser Gly Val	
	185		190	195	
	His Thr Phe Pro	Ala Val' Leu Gln	Ser Ser Gly Leu	Tyr Ser Leu	
	200		205	210	
10	Ser Ser Val Val	Thr Val Pro Ser	Ser Ser Leu Gly	Thr Gln Thr	
	215		220	225	
	Tyr Ile Cys Asn	Val Asn His Lys	Pro Ser Asn Thr	Lys Val Asp	
	230		235	240	
	Lys Lys Val Glu	Pro Lys Ser Cys	Asp Lys Thr His	Thr	
	245		250	253	
15	<210> 45				
	<211> 114				
	<212> PRT				
	<213> Mus musculus				
	<400> 45				
20	Asp Ile Val Met	Thr Gln Thr Pro	Leu Ser Leu Pro	Val Ser Leu	
	1	5	10	15	
	Gly Asp Gln Ala	Ser Ile Ser Cys	Arg Ser Ser Gln	Ser Leu Val	
	20		25	30	
25	His Gly Ile Gly	Asn Thr Tyr Leu	His Trp Tyr Leu	Gln Lys Pro	
	35		40	45	
	Gly Gln Ser Pro	Lys Leu Leu Ile	Tyr Tyr Lys Val	Ser Asn Arg	
	50		55	60	
	Phe Ser Gly Val	Pro Asp Arg Phe	Ser Asp Ser Gly	Ser Gly Thr	
	65		70	75	
30	Asp Phe Thr Leu	Arg Ile Ser Arg	Val Glu Ala Glu	Asp Leu Gly	
	80		85	90	
	Leu Tyr Phe Cys	Ser Gln Ser Thr	His Val Pro Leu	Thr Phe Gly	
	95		100	105	
35	Ala Gly Thr Lys	Leu Glu Leu Lys	Arg		
	110		114		
	<210> 46				
	<211> 114				
	<212> PRT				
	<213> Artificial Sequence				

<220>
 <221> Artificial Sequence
 <222> 1-114
 <223> recombinant immunoglobulin

5 <400> 46
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile' Thr Cys Arg Ser Ser Gln Ser Leu Val
 20 25 30
 10 His Gly Ile Gly Asn Thr Tyr Leu His Trp Tyr Gln Gln Lys Pro
 35 40 45
 Gly Lys Ala Pro Lys Leu Leu Ile Tyr Tyr Lys Val Ser Asn Arg
 50 55 60
 15 Phe Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr
 65 70 75
 Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala
 80 85 90
 Thr Tyr Tyr Cys Ser Gln Ser Thr His Val Pro Leu Thr Phe Gly
 95 100 105
 20 Gln Gly Thr Lys Val Glu Ile Lys Arg
 110 114
 <210> 47
 <211> 109
 <212> PRT
 25 <213> Homo sapiens
 <400> 47
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Lys Thr Ile Ser
 20 25 30
 30 Lys Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
 35 40 45
 Leu Leu Ile Tyr Tyr Ser Gly Ser Thr Leu Glu Ser Gly Val Pro
 50 55 60
 35 Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln
 80 85 90
 Gln His Asn Glu Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val
 95 100 105
 40

Glu Ile Lys Arg
109

<210> 48

<211> 117

5 <212> PRT

<213> Mus Musculus

<400> 48

Glu Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly
1 5 10 15

10 Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser
20 25 30

Ser His Tyr Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu
35 40 45

15 Glu Trp Ile Gly Tyr Ile Asp Pro Ser Asn Gly Glu Thr Thr Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser
65 70 75

Ser Ser Thr Ala Asn Val His Leu Ser Ser Leu Thr Ser Asp Asp
80 85 90

20 Ser Ala Val Tyr Phe Cys Ala Ala Arg Gly Asp Tyr Arg Tyr Asn
95 100 105

Gly Asp Trp Phe Phe Asp Val Trp Gly Ala Gly Thr
110 115 117

<210> 49

25 <211> 117

<212> PRT

<213> Artificial Sequence

<220>

<221> Artificial Sequence

30 <222> 1-117

<223> recombinant immunoglobulin

<400> 49

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
1 5 10 15

35 Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Ser Phe Ser
20 25 30

Ser His Tyr Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

40 Glu Trp Val Gly Tyr Ile Asp Pro Ser Asn Gly Glu Thr Thr Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser
 65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90

5 Thr Ala Val Tyr Tyr Cys Ala Ala Arg Gly Asp Tyr Arg Tyr Asn
 95 100 105

Gly Asp Trp Phe Phe Asp Val Trp Gly Gln Gly Thr
 110 115 117

10 <210> 50
 <211> 116
 <212> PRT
 <213> Homo sapiens

15 <400> 50
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Phe Thr
 20 25 30

Gly His Trp Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 35 40 45

20 Glu Trp Val Gly Met Ile His Pro Ser Asp Ser Glu Thr Arg Tyr
 50 55 60

Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser
 65 70 75

25 Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90

Thr Ala Val Tyr Tyr Cys Ala Ala Arg Gly Ile Tyr Phe Tyr Gly
 95 100 105

Thr Thr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr
 110 115 116

30 <210> 51
 <211> 242
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <221> Artificial Sequence
 <222> 1-242
 <223> recombinant immunoglobulin

40 <400> 51
 Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
 1 5 10 15

	Ser	Ile	Ala	Thr	Asn	Ala	Tyr	Ala	Asp	Ile	Gln	Met	Thr	Gln	Ser	
					20					25					30	
	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	
					35					40					45	
5	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Gly	Ile	Gly	Asn	Thr	Tyr	
					50					55					60	
	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	
					65					70					75	
10	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	Ser	Arg	Phe	
					80					85					90	
	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	
					95					100					105	
	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Ser	Gln	Ser	Thr	
					110					115					120	
15	His	Val	Pro	Leu	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	
					125					130					135	
	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	
					140					145					150	
20	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	
					155					160					165	
	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	
					170					175					180	
	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	
					185					190					195	
25	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	
					200					205					210	
	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	
					215					220					225	
30	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	
					230					235					240	
	Glu	Cys														
		242														
	<210>	52														
	<211>	253														
35	<212>	PRT														
	<213>	Artificial Sequence														
	<220>															
	<221>	Artificial Sequence														
	<222>	1-253														

<223> recombinant immunoglobulin

<400> 52

	Met	Lys	Lys	Asn	Ile	Ala	Phe	Leu	Leu	Ala	Ser	Met	Phe	Val	Phe	
	1				5						10				15	
5	Ser	Ile	Ala	Thr	Asn	Ala	Tyr	Ala	Glu	Val	Gln	Leu	Val	Gln	Ser	
					20					25					30	
	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	
					35					40					45	
10	Ala	Ala	Ser	Gly	Tyr	Ser	Phe	Ser	Ser	His	Tyr	Met	His	Trp	Val	
					50					55					60	
	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Gly	Tyr	Ile	Asp	
					65					70					75	
	Pro	Ser	Asn	Gly	Glu	Thr	Thr	Tyr	Asn	Gln	Lys	Phe	Lys	Gly	Arg	
					80					85					90	
15	Phe	Thr	Leu	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Ala	Tyr	Leu	Gln	
					95					100					105	
	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	
					110					115					120	
20	Arg	Gly	Asp	Tyr	Arg	Tyr	Asn	Gly	Asp	Trp	Phe	Phe	Asp	Val	Trp	
					125					130					135	
	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	
					140					145					150	
	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	
					155					160					165	
25	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	
					170					175					180	
	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	
					185					190					195	
30	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	
					200					205					210	
	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	
					215					220					225	
	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	
					230					235					240	
35	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr			
					245					250					255	

<210> 53

<211> 159

<212> PRT

<213> Artificial Sequence

<220>

<221> Artificial Sequence

5 <222> 1-159

<223> recombinant phage protein

<400> 53

Ser	Gly	Gly	Gly	Ser	Gly	Ser	Gly	Asp	Phe	Asp	Tyr	Glu	Lys	Met
1				5				10					15	

10	Ala	Asn	Ala	Asn	Lys	Gly	Ala	Met	Thr	Glu	Asn	Ala	Asp	Glu	Asn
				20				25						30	

	Ala	Leu	Gln	Ser	Asp	Ala	Lys	Gly	Lys	Leu	Asp	Ser	Val	Ala	Thr
				35				40						45	

15	Asp	Tyr	Gly	Ala	Ala	Ile	Asp	Gly	Phe	Ile	Gly	Asp	Val	Ser	Gly
				50				55						60	

	Leu	Ala	Asn	Gly	Asn	Gly	Ala	Thr	Gly	Asp	Phe	Ala	Gly	Ser	Ser
				65				70						75	

	Asn	Ser	Gln	Met	Ala	Gln	Val	Gly	Asp	Gly	Asp	Asn	Ser	Pro	Leu
				80				85						90	

20	Met	Asn	Asn	Phe	Arg	Gln	Tyr	Leu	Pro	Ser	Leu	Pro	Gln	Ser	Val
				95				100						105	

	Glu	Cys	Arg	Pro	Phe	Val	Phe	Ser	Ala	Gly	Lys	Pro	Tyr	Glu	Phe
				110				115						120	

25	Ser	Ile	Asp	Cys	Asp	Lys	Ile	Asn	Leu	Phe	Arg	Gly	Val	Phe	Ala
				125				130						135	

	Phe	Leu	Leu	Tyr	Val	Ala	Thr	Phe	Met	Tyr	Val	Phe	Ser	Thr	Phe
				140				145						150	

	Ala	Asn	Ile	Leu	Arg	Asn	Lys	Glu	Ser
				155				159	

30 <210> 54

<211> 780

<212> DNA

<213> Artificial Sequence

<220>

35 <221> Artificial Sequence

<222> 1-780

<223> recombinant immunoglobulin

<400> 54

atgaaaaaga atatcgcat tcttcttgca tctatgttcg tttttctat 50

40 tgctacaaac gcatacgctg atatccagat gaccagtc cccagctccc 100

tgtccgcctc tgtgggcat agggtcacca tcacctgcag gtcaagtcaa 150
 agcttagtac atggtatagg taacacgtat ttacactggg atcaacagaa 200
 accaggaaaa gctccgaaac tactgattta caaagtatcc aatcgattct 250
 ctggagtcctt ttctcgcttc tctggatccg gttctgggac ggatttcact 300
 5 ctgaccatca gcagtctgca gccagaagac ttcgcaactt attactgttc 350
 acagagtact catgtccgcg tcacgtttgg acaggggtacc aagggtggaga 400
 tcaaacgaac tgtggctgca ccatctgtct tcattctccc gccatctgat 450
 gagcagttga aatctggaac tgcttctgtt gtgtgctgc tgaataactt 500
 ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 550
 10 cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc 600
 tacagcctca gcagcaccct gacgctgagc aaagcagact acgagaaaca 650
 caaagtctac gcctgcgaag tcacccatca gggcctgagc tcgcccgtca 700
 caaagagctt caacagggga gagtggttaag ctgatcctct acgccggacg 750
 catcgtggcc ctagtacgca actagtcgta 780
 15 <210> 55
 <211> 253
 <212> PRT
 <213> Artificial Sequence
 <220>
 20 <221> Artificial Sequence
 <222> 1-253
 <223> recombinant immunoglobulin
 <400> 55
 Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
 25 1 5 10 15
 Ser Ile Ala Thr Asn Ala Tyr Ala Glu Val Gln Leu Val Glu Ser
 20 25 30
 Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys
 35 40 45
 30 Ala Ala Ser Gly Tyr Ser Phe Ser Ser His Tyr Met His Trp Val
 50 55 60
 Lys Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Tyr Ile Asp
 65 70 75
 Pro Ser Asn Gly Glu Thr Thr Tyr Asn Gln Lys Phe Lys Gly Arg
 80 85 90
 35

	Phe Thr Leu Ser Arg Asp Asn Ser Lys Asn Thr Ala Tyr Leu Gln	
	95	100 105
	Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr-Tyr Cys Ala	
	110	115 120
5	Arg Gly Asp Tyr Arg Tyr Asn Gly Asp Trp Phe Phe Asp Val Trp	
	125	130 135
	Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly	
	140	145 150
10	Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly	
	155	160 165
	Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu	
	170	175 180
	Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val	
	185	190 195
15	His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu	
	200	205 210
	Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr	
	215	220 225
20	Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp	
	230	235 240
	Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr	
	245	250 253
	<210> 56	
	<211> 242	
25	<212> PRT	
	<213> Artificial Sequence	
	<220>	
	<221> Artificial Sequence	
	<222> 1-242	
30	<223> recombinant immunoglobulin	
	<400> 56	
	Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe	
	1 5 10 15	
35	Ser Ile Ala Thr Asn Ala Tyr Ala Asp Ile Gln Met Thr Gln Ser	
	20 25 30	
	Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr	
	35 40 45	
	Cys Arg Ser Ser Gln Ser Leu Val His Gly Ile Gly Ala Thr Tyr	
	50 55 60	

Leu His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu
 65 70 75
 Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro-Ser Arg Phe
 80 85 90

5 Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
 95 100 105
 Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Ser Gln Ser Thr
 110 115 120
 10 His Val Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 125 130 135
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
 140 145 150
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn
 155 160 165
 15 Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn
 170 175 180
 Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp
 185 190 195
 20 Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser
 200 205 210
 Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr
 215 220 225
 His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly
 230 235 240
 25 Glu Cys
 242
 <210> 57
 <211> 45
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <221> Artificial Sequence
 <222> 1-45
 <223> recombinant leucine zipper peptide
 35 <400> 57
 Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Arg Met Lys
 1 5 10 15
 Gln Leu Glu Asp Lys Val Glu Glu Leu Leu Ser Lys Asn Tyr His
 20 25 30

Leu Glu Asn Glu Val Ala Arg Leu Lys Lys Leu Val Gly Glu Arg
 35 40 45

<210> 58

<211> 780

5 <212> DNA

<213> Artificial Sequence

<220>

<221> Artificial Sequence

<222> 1-780

10 <223> recombinant immunoglobulin

<400> 58

atgaaaaaga atatcgcat ttttcttgca tctatgttcg tttttctat 50

tgctacaaac gcatacgtg atatccagat gaccagtc ccgagctccc 100

tgtccgctc tgtggcgat agggcacca tcacctgcag gtcaagtcaa 150

15 agcttagtac atggtatagg tgctacgtat ttacactggg atcaacagaa 200

accaggaaaa gctccgaaac tactgattta caaagtatcc aatcgattct 250

ctggagtc cc ttctcgctt tctggatccg gttctgggac ggatttcact 300

ctgaccatca gcagtctgca gccagaagac ttcgcaactt attactgttc 350

acagagtact catgtccgc tcacgtttgg acaggggtacc aagggtggaga 400

20 tcaaacgaac tgtggctgca ccatctgtct tcatcttccc gccatctgat 450

gagcagttga aatctggaac tgcttctgtt gtgtgctgc tgaataactt 500

ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 550

cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc 600

tacagcctca gcagcaccct gacgctgagc aaagcagact acgagaaaca 650

25 caaagtctac gcctgcgaag tcacccatca gggcctgagc tcgcccgtca 700

caaagagctt caacagggga gagtgttaag ctgatectct acgccggacg 750

catcgtggcc ctagtacgca actagtcgta 780

<210> 59

<211> 927

30 <212> DNA

<213> Artificial Sequence

<220>

<221> Artificial Sequence

<222> 1-927

35 <223> recombinant immunoglobulin

<400> 59

aaaagggtat ctagagggtg aggtgatttt atgaaaaaga atatcgatt 50

tcttcttgca tctatgttcg ttttttctat tgctacaaac gcgtacgctg 100

aggttcagct agtgcagtct ggcgggtggc tgggtgcagcc agggggctca 150

5 ctcctgtttgt cctgtgcagc ttctggctac tccttctcga gtcactatat 200

gcactgggtc cgtcaggccc cgggtaaggg cctggaatgg gttggatata 250

ttgatccttc caatggtgaa actacgtata atcaaaaagt caagggccgt 300

ttcactttat ctgcgcgaaa ctccaaaaac acagcatacc tgcagatgaa 350

cagcctgcgt gctgaggaca ctgccgtcta ttactgtgca agaggggatt 400

10 atcgctacaa tgggtgactgg ttcttcgacg tctgggggtca aggaaccctg 450

gtcacctctt cctcggcctc caccaagggc ccatcggtct tccccctggc 500

accctcctcc aagagcacct ctgggggcac agcggccctg ggctgcctgg 550

tcaaggacta cttccccgaa ccggtgacgg tgtcgtggaa ctcaggcgcc 600

ctgaccagcg gcgtgcacac cttccccggt gtcttacag cctcaggact 650

15 ctactccctc agcagcgtgg tgaccgtgcc ctccagcagc ttgggcaccc 700

agacctacat ctgcaacgtg aatcacaagc ccagcaacac caaggtcgac 750

aagaaagttg agcccaaate ttgtgacaaa actcacacat gcccgcctg 800

cccagcacca gaactgctgg gcggccgcat gaaacagcta gaggacaagg 850

tcgaagagct actctccaag aactaccacc tagagaatga agtggcaaga 900

20 ctcaaaaagc ttgtcgggga gcgctaa 927

<210> 60

<211> 298

<212> PRT

<213> Artificial Sequence

25 <220>

<221> Artificial Sequence

<222> 1-298

<223> recombinant immunoglobulin

<400> 60

30 Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
1 5 10 15Ser Ile Ala Thr Asn Ala Tyr Ala Glu Val Gln Leu Val Gln Ser
20 25 30

	Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys	35	40	45
	Ala Ala Ser Gly Tyr Ser Phe Ser Ser His Tyr Met His Trp Val	50	55	60
5	Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Tyr Ile Asp	65	70	75
	Pro Ser Asn Gly Glu Thr Thr Tyr Asn Gln Lys Phe Lys Gly Arg	80	85	90
10	Phe Thr Leu Ser Arg Asp Asn Ser Lys Asn Thr Ala Tyr Leu Gln	95	100	105
	Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala	110	115	120
	Arg Gly Asp Tyr Arg Tyr Asn Gly Asp Trp Phe Phe Asp Val Trp	125	130	135
15	Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly	140	145	150
	Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly	155	160	165
20	Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu	170	175	180
	Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val	185	190	195
	His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu	200	205	210
25	Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr	215	220	225
	Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp	230	235	240
30	Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro	245	250	255
	Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Arg Met Lys Gln Leu	260	265	270
	Glu Asp Lys Val Glu Glu Leu Leu Ser Lys Asn Tyr His Leu Glu	275	280	285
35	Asn Glu Val Ala Arg Leu Lys Lys Leu Val Gly Glu Arg	290	295	298

<210> 61

<211> 6563

<212> DNA

<213> Artificial Sequence

<220>

<221> Artificial Sequence

5 <222> 1-6563

<223> recombinant immunoglobulin

<400> 61

gaattcaact tctccatact ttggataagg aaatacagac atgaaaaatc 50
 tcattgctga gttgttattt aagcttgccc aaaaagaaga agagtcgaat 100
 10 gaactgtgtg cgcaggtaga agctttggag attatcgtca ctgcaatgct 150
 tcgcaatatg gcgcaaaatg accaacagcg gttgattgat caggtagagg 200
 gggcgctgta cgaggtaaag cccgatgcc a gattcctga cgacgatacg 250
 gagctgctgc gcgattacgt aaagaagtta ttgaagcatc ctcgtcagta 300
 aaaagttaat cttttcaaca gctgtcataa agttgtcacg gccgagactt 350
 15 atagtcgctt tgtttttatt ttttaatgta tttgtaacta gaattcgagc 400
 tcggtacccg gggatcctct cgaggttgag gtgattttat gaaaaagaat 450
 atcgcatctt ttcttgcac tatgttcgtt ttttctattg ctacaaacgc 500
 atacgctgat atccagatga cccagtcgcc gagctccctg tccgcctctg 550
 tgggcatag ggtcaccatc acctgcaggt caagtcaaag cttagtacat 600
 20 ggtatagggtg ctacgtattt aactgggtat caacgaaac caggaaaagc 650
 tccgaaacta ctgatttaca aagtatccaa tcgattctct ggagtcctt 700
 ctcgcttctc tggatccggt tctgggacgg atttcactct gaccatcagc 750
 agtctgcagc cagaagactt cgcaacttat tactgttcac agagtactca 800
 tgtcccgctc acgtttggac agggtaacaa ggtggagatc aaacgaactg 850
 25 tggctgcacc atctgtcttc atcttcccgc catctgatga gcagttgaaa 900
 tctggaactg cttctgttgt gtgcctgctg aataacttct atcccagaga 950
 ggccaaagta cagtgaagg tggataacgc cctccaatcg ggtaactccc 1000
 aggagagtgt cacagagcag gacagcaagg acagcaccta cagcctcagc 1050
 agcaccctga cgctgagcaa agcagactac gagaaacaca aagtctacgc 1100
 30 ctgcgaagtc acccatcagg gcctgagctc gcccgtcaca aagagcttca 1150
 acaggggaga gtgttaagct gatcctctac gccggacgca tcgtggccct 1200

agtacgcaac tagtcgtaaa aagggtatct agaggttgag gtgattttat 1250
 gaaaaagaat atcgcatctt ttcttgcatc tatgttcgtt ttttctattg 1300
 ctacaaacgc gtacgctgag gttcagctag tgcagtctgg cgggtggcctg 1350
 gtgcagccag ggggctcact ccgtttgtcc tgtgcagctt ctggctactc 1400
 5 cttctcgagt cactatatgc actgggtccg tcaggccccg ggtaagggcc 1450
 tggaatgggt tggatatatt gatccttcca atggtgaaac tacgtataat 1500
 caaaagtcca agggccgttt cactttatct cgcgacaact ccaaaaacac 1550
 agcatacctg cagatgaaca gcctgcgtgc tgaggacact gccgtctatt 1600
 actgtgcaag aggggattat cgctacaatg gtgactgggt cttcgacgtc 1650
 10 tggggccaag gaacctggt caccgtctcc tcggcctcca ccaagggccc 1700
 atcggctctc cccctggcac cctcctccaa gagcacctct gggggcacag 1750
 cggccctggg ctgcctggtc aaggactact tccccgaacc ggtgacgggtg 1800
 tcgtggaact caggcgccct gaccagcggc gtgcacacct tcccggctgt 1850
 cctacagtec tcaggactct actcctcag cagcgtgggt accgtgccct 1900
 15 ccagcagctt gggcacccag acctacatct gcaacgtgaa tcacaagccc 1950
 agcaacacca aggtcgacaa gaaagttgag cccaaatctt gtgacaaaac 2000
 tcacacatgc ccgccgtgcc cagcaccaga actgctgggc ggccgcatga 2050
 aacagctaga ggacaaggtc gaagagctac tctccaagaa ctaccaccta 2100
 gagaatgaag tggcaagact caaaaagctt gtcggggagc gctaagcatg 2150
 20 cgacggccct agagtccta acgtcgggt gccgccgggc gttttttatt 2200
 gttaactcat gtttgacagc ttatcatcga taagctttaa tgcggtagtt 2250
 tatcacagtt aaattgctaa cgcagtcagg caccgtgtat gaaatctaac 2300
 aatgcgtca tcgtcatcct cggcacgct accctggatg ctgtaggcat 2350
 aggcttggtt atgccggtac tgccgggcct cttgcgggat atcgtccatt 2400
 25 ccgacagcat cgccagtcac tatggcgtgc tgctagcgt atatgcgttg 2450
 atgcaatttc tatgcgcacc cgttctcgga gcactgtccg accgctttgg 2500
 ccgcgcacca gtctgtctg cttcgctact tggagccact atcgactacg 2550
 cgatcatggc gaccacacce gtctgtgga tcctctacgc cggacgcac 2600

gtggccggca tcaccggcgc cacaggtgcg gttgctggcg cctatatcgc 2650
 cgacatcacc gatggggaag atcgggctcg ccacttcggg ctcatgagcg 2700
 cttgtttcgg cgtgggtatg gtggcaggcc ccgtggccgg gggactgttg 2750
 ggcgccatct ccttgcacgc accattcctt ~~gcggggcg~~ ~~tgctcaacgg~~ 2800
 5 cctcaaccta ctactgggct gcttcctaata gcaggagtcg cataaggagg 2850
 agcgtcgtcc gatgcccttg agagccttca acccagtcag ctccttcggg 2900
 tgggcgcggg gcattgactat cgtcgccgca cttatgactg tcttctttat 2950
 catgcaactc gtaggacagg tgccggcagc gctctgggctc attttcggcg 3000
 aggaccgctt tcgctggagc gcgacgatga tcggcctgtc gcttgcggtta 3050
 10 ttcggaatct tgcacgccct cgctcaagcc ttcgtcactg gtcccgcac 3100
 caaacgtttc ggcgagaagc aggccattat cgccggcatg gcggccgacg 3150
 cgctgggcta cgtcttgctg gcgttcgca cgcgaggctg gatggccttc 3200
~~gccattatga~~ ~~ttcttctcgc~~ ~~ttccggcggc~~ ~~atcgggatgc~~ ~~ccgcgttgca~~ 3250
 ggccatgctg tccaggcagg tagatgacga ccatcaggga cagcttcaag 3300
 15 gatcgtcgc ggctcttacc agcctaactt cgatcactgg accgctgac 3350
 gtcacggcga tttatgccgc ctcggcgagc acatggaacg ~~ggttggcatg~~ 3400
 gattgtaggc gccgccctat accttgctc cctccccgcg ttgcgtcgcg 3450
 gtgcatggag ccgggccacc tcgacctgaa tggaagccgg cggcacctcg 3500
 ctaacggatt caccactcca agaattggag ccaatcaatt cttgcggaga 3550
 20 actgtgaatg cgcaaaccaa cccttggcag aacatatcca tcgcgtccgc 3600
 catctccagc agccgcacgc ggcgcactc gggcagcgtt gggtcctggc 3650
 cacgggtgcg catgatcgtg ctctgtcgt tgaggaccgc gctaggctgg 3700
 cgggggttgc ttactgggta gcagaatgaa tcaccgatac gcgagcgaac 3750
 gtgaagcgac tgctgctgca aaacgtctgc gacctgagca acaacatgaa 3800
 25 tggctcttcgg tttccgtgtt tcgtaaagtc tggaaacgcg gaagtcagcg 3850
 ccctgcacca ttatgttccg gatctgcac gcaggatgct gctggctacc 3900
 ctgtggaaca cctacatctg tattaacgaa gcgctggcat tgacctgag 3950
 tgatttttct ctggtcgcgc cgcattccata ccgccagttg tttacctca 4000

caacgttcca gtaaccgggc atgttcatca tcagtaaccc gtatcgtgag 4050
 catcctctct cgtttcatcg gtatcattac ccccatgaac agaaattccc 4100
 ccttacacgg aggcatacaag tgaccaaaaca ggaaaaaacc gcccttaaca 4150
 tggcccgctt tatcagaagc cagacattaa cgcttctgga gaaactcaac 4200
 5 gagctggacg cggatgaaca ggcagacatc tgtgaatcgc ttcacgacca 4250
 cgctgatgag ctttaccgca gctgcctcgc gcgtttcggg gatgacgggtg 4300
 aaaacctctg acacatgcag ctcccgagga cggtcacagc ttgtctgtaa 4350
 gcggatgccg ggagcagaca agcccgtcag ggcgcgtcag cgggtgttgg 4400
 cgggtgtcgg ggcgcagcca tgaccagtc acgtagcgat agcggagtgt 4450
 10 atactggctt aactatgcgg catcagagca gattgtactg agagtgcacc 4500
 atatgcgggtg tgaaataccg cacagatgcg taaggagaaa ataccgcac 4550
 aggcgctctt ccgcttctc gctcactgac tcgctgcgct cggtcgttcg 4600
 gctgcggcga gcggtatcag ctactcaaa ggcggtaata cggttatcca 4650
 cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa 4700
 15 aaggccagga accgtaaaaa ggcgcggttg ctggcgtttt tccataggct 4750
 ccgccccct gacgagcatc acaaaaatcg acgctcaagt cagaggtggc 4800
 gaaacccgac aggactataa agataccagg cgtttcccc tggaagctcc 4850
 ctgctgcgct ctctgttcc gaccctgccg cttaccggat acctgtccgc 4900
 ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt 4950
 20 atctcagttc ggtgtaggtc gttcgtcca agctgggctg tgtgcacgaa 5000
 cccccgttc agcccgaccg ctgcgcctta tccggttaact atcgtcttga 5050
 gtccaacccg gtaagacacg acttatcgcc actggcagca gccactggta 5100
 acaggattag cagagcgagg tatgtaggcg gtgctacaga gttcttgaag 5150
 tggtaggcta actacggcta cactagaagg acagtatttg gtatctgcgc 5200
 25 tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg 5250
 gcaaacaac caccgctggg agcggtggtt ttttgtttg caagcagcag 5300
 attacgcgca gaaaaaagg atctcaagaa gatccttga tcttttctac 5350
 ggggtctgac gctcagtga acgaaaactc acgttaaggg attttggta 5400

tgagattatc aaaaaggatc ttcacctaga tccttttaaa ttaaaaatga 5450
 agttttaaat caatctaaag tatatatgag taaacttggt ctgacagtta 5500
 ccaatgctta atcagtgagg cacctatctc agcgatctgt ctatttcgtt 5550
 catccatagt tgcctgactc cccgtcgtgt agataaetac gatacgggag 5600
 5 ggcttaccat ctggccccag tgctgcaatg ataccgag agccacgctc 5650
 accggctcca gatttatcag caataaacca gccagccgga agggccgagc 5700
 gcagaagtgg tctgcaact ttatccgctt ccatccagtc tattaattgt 5750
 tgccgggaag ctagagtaag tagttcgcca gttaatagtt tgcgcaacgt 5800
 tgttgccatt gctgcaggca tctggtgtc acgctcgtcg tttggtatgg 5850
 10 cttcattcag ctccggttcc caacgatcaa ggcgagttac atgatcccc 5900
 atgttgtgca aaaaagcggg tagctccttc ggtcctccga tctgtgtcag 5950
 aagtaagttg gccgcagtgt tatcactcat gggtatggca gcactgcata 6000
 attctcttac tgtcatgcca tccgtaagat gcttttctgt gactggtgag 6050
 tactcaacca agtcattctg agaatagtgt atgcggegac cgagttgctc 6100
 15 ttgcccggcg tcaacacggg ataataccgc gccacatagc agaactttaa 6150
 aagtgtcat cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc 6200
 ttaccgctgt tgagatccag ttcatgttaa cccactcgtg cacccaactg 6250
 atcttcagca tcttttactt tcaccagcgt ttctgggtga gcaaaaacag 6300
 gaaggcaaaa tgccgcaaaa aagggaataa ggcgacacg gaaatgttga 6350
 20 atactcatac tcttcctttt tcaatattat tgaagcattt atcaggggta 6400
 ttgtctcatg agcggataca tatttgaatg tatttagaaa aataaaciaa 6450
 taggggttcc gcgcacattt ccccgaaaag tgccacctga cgtctaagaa 6500
 accattatta tcatgacatt aacctataaa aataggcgta tcacgaggcc 6550
 ctttcgtctt caa 6563
 25 <210> 62
 <211> 242
 <212> PRT
 <213> Artificial Sequence
 <220>
 30 <221> Artificial Sequence
 <222> 1-242

<223> recombinant immunoglobulin

<400> 62

	Met	Lys	Lys	Asn	Ile	Ala	Phe	Leu	Leu	Ala	Ser	Met	Phe	Val	Phe	
	1				5					10					15	
5	Ser	Ile	Ala	Thr	Asn	Ala	Tyr	Ala	Asp	Ile	Gln	Met	Thr	Gln	Ser	
					20					25					30	
	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	
					35					40					45	
10	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Gly	Ile	Gly	Glu	Thr	Tyr	
					50					55					60	
	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	
					65					70					75	
	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	Ser	Arg	Phe	
					80					85					90	
15	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	
					95					100					105	
	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Ser	Gln	Ser	Thr	
					110					115					120	
20	His	Val	Pro	Leu	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	
					125					130					135	
	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	
					140					145					150	
	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	
					155					160					165	
25	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	
					170					175					180	
	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	
					185					190					195	
30	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	
					200					205					210	
	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	
					215					220					225	
	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	
					230					235					240	
35	Glu	Cys														
		242														

<210> 63

<211> 27

<212> DNA
 <213> Artificial Sequence

<220>
 <221> Artificial Sequence

5 <222> 1-27
 <223> recombinant immunoglobulin

<400> 63
 catggtatag gttaaactta tttacac 27

10 <210> 64
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <221> Artificial Sequence
 15 <222> 1-27
 <223> recombinant immunoglobulin

<220>
 <221> n
 <222> 13
 20 <223> a or g or c or t

<220>
 <221> n
 <222> 14
 <223> a or g or c or t

25 <400> 64
 catggtatag gtnnsactta tttacac 27

<210> 65
 <211> 780
 <212> DNA
 30 <213> Artificial Sequence

<220>
 <221> Artificial Sequence
 <222> 1-780
 <223> recombinant immunoglobulin

35 <400> 65
 atgaaaaaga atatcgcat tcttcttgca tctatgttcg ttttttctat 50
 tgctacaaac gcatacgtg atatccagat gacccagtcc ccgagctccc 100
 tgtccgctc tgtgggcgat agggtcacca tcacctgcag gtcaagtcaa 150
 agcttagtac atggtatagg tgagacgtat ttacactggg atcaacagaa 200
 40 accaggaaaa gctccgaaac tactgattta caaagtatcc aatcgattct 250
 ctggagtcctc ttctcgcttc tctggatccg gttctgggac ggatttcact 300

ctgaccatca gcagtctgca gccagaagac ttcgcaactt attactgttc 350
acagagtact catgtcccg ctcacgtttgg acagggtacc aagggtggaga 400
tcaaacgaac tgtggctgca ccactctgtct tcactctccc gccactctgat 450
gagcagttga aatctggaac tgcttctgtt gtgtgcctgc tgaataactt 500
5 ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 550
egggtaaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc 600
tacagcctca gcagcaccct gacgctgagc aaagcagact acgagaaaca 650
caaagtctac gcctgcgaag tcacccatca gggcctgagc tcgcccgta 700
caaagagctt caacagggga gagtgttaag ctgacccctt acgccggacg 750
10 catcgtggcc ctagtacgca actagtcgta 780

<210> 66
<211> 78
<212> DNA
<213> Artificial Sequence

15 <220>
<221> Artificial Sequence
<222> 1-78
<223> recombinant immunoglobulin

<400> 66

20 ctagtgcagt ctggcgggtg cctggtgcag ccagggggct cactcgttt 50
gtcctgtgca gcttctggct actccttc 78

<210> 67
<211> 82
<212> DNA

25 <213> Artificial Sequence

<220>
<221> Artificial Sequence
<222> 1-82
<223> recombinant immunoglobulin

30 <400> 67

tcgagaagga gtagccagaa gctgcacagg acaaacggag tgagccccct 50
ggctgcacca ggccaccgcc agactgcact ag 82

<210> 68
<211> 8120

35 <212> DNA
<213> Artificial Sequence

<220>

<221> Artificial Sequence
 <222> 1-8120
 <223> recombinant immunoglobulin

<400> 68

5 ~~ttc~~gagctcg cccgacattg attattgact agagtcgac gacagctgtg 50
 gaatgtgtgt cagttagggt gtggaaagtc cccagggtcc ccagcaggca 100
 gaagtatgca aagcatgcat ctcaattagt cagcaaccag gtgtggaaaag 150
 tccccaggct ccccgacagg cagaagtatg caaagcatgc atctcaatta 200
 gtcagcaacc atagtcccg ccctaactcc gcccatcccg ccctaactc 250
 10 cgcccagttc cgcccattct cgcgcccatg gctgactaat tttttttatt 300
 tatgcagagg ccgaggccgc ctcgccctct gagctattcc agaagtagtg 350
 aggaggcttt tttggaggcc taggcttttg caaaaagcta gcttatccgg 400
 ccgggaacgg tgcattggaa cgcggattcc ccgtgccaaag agtgacgtaa 450
 gtaccgccta tagagcgata agaggatttt atccccgctg ccatcatggt 500
 15 tgcaccattg aactgcatcg tcgccgtgtc ccaaaatacg gggattggca 550
 agaacggaga cctaccctgg cctccgctca ggaacgagtt caagtacttc 600
~~caaagaatga ccacaacctc~~ ttcagtgga ggtaaacaga atctgggtgat 650
 tatgggtagg aaaacctggt tctccattcc tgagaagaat cgacctttaa 700
 aggacagaat taatatagtt ctcatgtag aactcaaaga accaccacga 750
 20 ggagctcatt ttcttgccaa aagtttggat gatgccttaa gacttattga 800
 acaaccggaa ttggcaagta aagtagacat ggtttggata gtcggaggca 850
 gttctgttta ccaggaagcc atgaatcaac caggccacct tagactcttt 900
 gtgacaagga tcatgcagga atttgaaagt gacacgtttt tcccagaaat 950
 tgatttgggg aaatataaac ctctcccaga ataccaggc gtcctctctg 1000
 25 aggtccagga ggaaaaaggc atcaagtata agtttgaagt ctacgagaag 1050
 aaagactaac aggaagatgc tttcaagttc tctgctcccc tcctaaagct 1100
 atgcattttt ataagaccat gggacttttg ctggcttttag atccccctgg 1150
 cttcgttaga acgcagctac aattaatata taaccttatg tatcatacac 1200
 atacgattta ggtgacacta tagataacat ccactttgcc tttctctcca 1250
 30 caggtgtcca ctcccaggtc caactgcacc tcggttctat cgattgaatt 1300

ccaccatggg atggtcacgt atcacccttt ttctagtagc aactgcaact 1350
 ggagtacatt cagaagttca gctagtgcag tctggcggtg gcctgggtgca 1400
 gccagggggc tcaactcgtt tgcctgtgc agcttctggc tactccttct 1450
 cgagtcacta tatgcactgg gtccgtcagg ccccggttaa gggcctggaa 1500
 5 tgggttggat atattgatcc ttccaatggt gaaactacgt ataatacaaa 1550
 gttcaagggc cgtttcactt tatctcgca caactccaaa aacacagcat 1600
 acctgcagat gaacagcctg cgtgctgagg aactgccgt ctattactgt 1650
 gcaagagggg attatcgcta caatggtgac tggttcttcg acgtctgggg 1700
 tcaaggaacc ctggtcaccg tctcctcggc ctccaccaag ggcccatcgg 1750
 10 tcttccccct,ggcaccctcc tccaagagca cctctggggg cacagcggcc 1800
 ctgggctgcc tgggtcaagga ctacttcccc gaaccggtga cgggtgctgt 1850
 gaactcaggc gccctgacca gcggcgtgca caccttcccc gctgtcctac 1900
 agtcctcagg actctactcc ctacgagcg tggtgactgt gccctctagc 1950
 agcttgggca cccagaccta catctgcaac gtgaatcaca agccagcaa 2000
 15 caccaaggtg gacaagaaag ttgagcccaa atcttgtgac aaaactcaca 2050
 catgccacc gtgcccagca cctgaactcc tggggggacc gtcagtcttc 2100
 ctcttcccc caaaacccaa ggacaccctc atgatctccc ggaccctga 2150
 ggtcacatgc gtggtggtgg acgtgagcca cgaagaccct gaggtcaagt 2200
 tcaactggta cgtggacggc gtggaggtgc ataatgcca gacaaagccg 2250
 20 cgggaggagc agtacaacag cacgtaccgt gtggtcagcg tcctcaccgt 2300
 cctgcaccag gactggctga atggcaagga gtacaagtgc aaggtctcca 2350
 acaaagccct cccagcccc atcgagaaaa ccatctccaa agccaaagg 2400
 cagccccgag aaccacaggt gtacaccctg ccccatccc gggaagagat 2450
 gaccaagaac caggtcagcc tgacctgct ggtcaaaggc ttctatccca 2500
 25 gcgacatcg cgtggagtgg gagagcaatg ggcagccgga gaacaactac 2550
 aagaccagc ctcccggtgt ggactccgac ggctccttct tcctctacag 2600
 caagctcacc gtggacaaga gcaggtggca gcaggggaac gtcttctcat 2650
 gctccgtgat gcatgaggct ctgcacaacc actacacgca gaagagcctc 2700

tccctgtctc cgggtaaagt agtgcgacgg ccctagagtc gacctgcaga 2750
 agcttgcccg ccatggccca acttgtttat tgcagcttat aatggttaca 2800
 aataaagcaa tagcatcaca aatttcacaa ataaagcatt tttttcactg 2850
 cattctagtt gtggtttgc ~~caaactcacc~~ ~~aatgtatctt~~ ~~atcatgtctg~~ 2900
 5 gatcgatcgg gaattaattc ggcgacgac catggcctga aataacctct 2950
 gaaagaggaa cttgggttagg taccttctga ggcgaaaga accatctgtg 3000
 gaatgtgtgt cagttagggt gtggaaagtc cccaggtcc ccagcaggca 3050
 gaagtatgca aagcatgcat ctcaattagt cagcaaccag gtgtggaaag 3100
 tccccaggct cccagcagg cagaagtatg caaagcatgc atctcaatta 3150
 10 gtcagcaacc atagtccgc cctaactcc gccatcccg ccctaactc 3200
 cgcccagttc cgccattct cgcccatg gctgactaat ttttttatt 3250
 tatgcagagg ccgaggccgc ctggcctct gagctattcc agaagtagtg 3300
 aggaggcttt tttggaggcc taggcttttg caaaaagcta gcttatccgg 3350
 ccgggaacgg tgcattggaa cgcgattcc cctgccaag agtcaggtaa 3400
 15 gtaccgcta tagagtctat agggccacc ccttggttc gttagaacgc 3450
 ggctacaatt aatacataac cttttggatc ~~gacccactg~~ ~~acactgacat~~ 3500
 ccacttttcc tttttctcca cagggtgcca ctccaggtc caactgcacc 3550
 tcggttcgcy aagctagctt gggctgcac gattgaattc caccatggga 3600
 tggatcatga tcatcctttt tctagtagc actgcaactg gactacattc 3650
 20 agatatccag atgaccagtc ccccgagtc cctgtccgc tctgtgggcy 3700
 atagggtcac catcacctgc aggtcaagtc aaagcttagt acatgggtata 3750
 ggtgctacgt atttactg gtatcaacag aaaccaggaa aagctccgaa 3800
 actactgatt taaaagtat ccaatcgatt ctctggagtc ctttctcgct 3850
 tctctggatc cggttctggg acggatttca ctctgaccat cagcagctctg 3900
 25 cagccagaag acttcgcaac ttattactgt tcacagagta ctcatgtccc 3950
 gctcacgttt ggacagggtg ccaagggtga gatcaaaca actgtggctg 4000
 caccatctgt cttcatcttc ccgcatctg atgagcagtt gaaatctgga 4050
 actgcttctg ttgtgtgct gctgaataac ttctatccca gagaggccaa 4100

agtacagtgg aaggtggata acgccctcca atcgggtaac tcccaggaga 4150
 gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc 4200
 ctgacgctga gcaaagcaga ctacgagaaa cacaaagtct acgcctgcga 4250
 agtcacccat cagggcctga gctcgcccggt cacaaagagc ttcaacaggg 4300
 5 gagagtgtta agcttggccg ccatggccca acttgtttat tgcagcttat 4350
 aatggttaca aataaagcaa tagcatcaca aatttcacaa ataaagcatt 4400
 tttttcactg cattctagtt gtggtttgtc caaactcatc aatgtatctt 4450
 atcatgtctg gatcgatcgg gaattaattc ggcgagcac catggcctga 4500
 aataacctct gaaagaggaa cttggttagg taccttctga ggcggaaga 4550
 10 accagctgtg gaatgtgtgt cagttagggt gtggaaagtc cccaggctcc 4600
 ccagcaggca gaagtatgca aagcatgcat ctcaattagt cagcaaccag 4650
 gtgtggaaag tccccaggct cccagcagg cagaagtatg caaagcatgc 4700
 atctcaatta gtcagcaacc atagtccgc ccctaactcc gcccatcccg 4750
 ccctaactc cgccagttc cgccattct cgcgccatg gctgactaat 4800
 15 tttttttatt tatgcagagg ccgaggccgc ctggcctct gagctattcc 4850
 agaagtagtg aggaggcttt tttggaggcc taggcttttg caaaaagctg 4900
 ttacctcgag cgccgctta attaaggcgc gccatttaa tctgcaggt 4950
 aacagcttgg cactggccgt cgttttaca cgctgtgact gggaaaacc 5000
 tggcgttacc caacttaac gccttgtagc acatcccccc ttcgccagct 5050
 20 ggcgtaatag cgaagaggcc cgcaccgac gcccttccca acagttgcgt 5100
 agcctgaatg gcgaatggcg cctgatgcgg tattttctcc ttacgcatt 5150
 gtgcggtatt tcacaccgca tacgtcaaag caaccatagt acgcgcctg 5200
 tagcggcgca ttaagcgcg cggtgtggt ggttacgcgc agcgtgaccg 5250
 ctacacttgc cagcgcccta gcgccgctc ctttcgcttt cttcccttcc 5300
 25 tttctcgcca cgttcgcgg ctttccccgt caagctctaa atcgggggct 5350
 ccttttaggg ttccgattta gtgctttacg gcacctgac cccaaaaaac 5400
 ttgatttggg tgatggttca cgtagtgggc catcgccctg atagacggtt 5450
 tttcgccctt tgacgttga gtccacgttc ttaatagtg gactcttgtt 5500

ccaaactgga acaacactca accctatctc gggctattct tttgatttat 5550
aagggatttt gccgatttcg gcctattggt taaaaaatga gctgatttaa 5600
caaaaattta acgcgaattt taacaaaata ttaacgttta caattttatg 5650
gtgcactctc agtacaatct gctctgatgc cgcatagtta agccaactcc 5700
5 gctatcgcta cgtgactggg tcatggctgc gccccgacac ccgccaacac 5750
ccgctgacgc gccctgacgg gcttgtctgc tcccggcatc cgcttacaga 5800
caagctgtga ccgtctccgg gagctgcatg tgtcagaggt tttcaccgtc 5850
atcaccgaaa cgcgcgaggg agtattcttg aagacgaaag ggcctcgtga 5900
tacgcctatt tttatagggt aatgtcatga taataatggt ttcttagacg 5950
10 tcaggtggca cttttcgggg aaatgtgcgc ggaacccta tttgtttatt 6000
tttctaaata cattcaaata tgtatccgct catgagacaa taaccctgat 6050
aaatgcttca ataattattga aaaaggaaga gtatgagtat tcaacatttc 6100
cggtgtcgccc ttattccctt ttttgcgga ttttgccctc ctgtttttgc 6150
tcaccagaa acgctggtga aagtaaaaga tgctgaagat cagttgggtg 6200
15 cagagtggg ttacatcgaa ctggatctca acagcggtaa gatccttgag 6250
agttttcgcc ccgaagaacg ttttccaatg atgagcactt ttaaagttct 6300
gctatgtggc gcggtattat cccgtgatga cgccgggcaa gagcaactcg 6350
gtcgccgcat acactattct cagaatgact tggttgagta ctcaccagtc 6400
acagaaaagc atcttacgga tggcatgaca gtaagagaat tatgcagtgc 6450
20 tgccataacc atgagtgata aactgcggc caacttactt ctgacaacga 6500
tcggaggacc gaaggagcta accgcttttt tgcacaacat gggggatcat 6550
gtaactcgcc ttgatcggtg ggaaccggag ctgaatgaag ccataccaaa 6600
cgacgagcgt gacaccacga tgccagcagc aatggcaaca acgttgcgca 6650
aactattaac tggcgaacta cttactctag cttcccggca acaattaata 6700
25 gactggatgg aggcggataa agttgcagga ccacttctgc gctcggccct 6750
tccggtggc tggttttattg ctgataaatc tggagccggt gagegtgggt 6800
ctcgcggtat cattgcagca ctggggccag atggtaagcc ctcccgtatc 6850
gtagttatct acacgacggg gagtcaggca actatggatg aacgaaatag 6900

acagatcgct gagatagggtg cctcactgat taagcattgg taactgtcag 6950
accaagttta ctcatatata ctttagattg atttaaaact tcatttttaa 7000
tttaaaagga tctagggtgaa gatccttttt gataatctca tgacccaaat 7050
cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga 7100
5 tcaaaggatc ttcttgagat cctttttttc tgcgcgtaat ctgctgcttg 7150
caaacaaaaa aaccaccgct accagcgggtg gtttgtttgc cggatcaaga 7200
gctaccaact ctttttccga aggtaactgg cttcagcaga gcgcagatac 7250
caaatactgt ccttctagtg tagccgtagt taggccacca cttcaagaac 7300
tctgtagcac cgctacata cctcgtcttg ctaatcctgt taccagtggc 7350
10 tgctgccagt ggcgataagt cgtgtcttac cgggttggac tcaagacgat 7400
agttaccgga taaggcgag cggtcgggct gaacgggggg ttcgtgcaca 7450
cagcccagct tggagcgaac gacctacacc gaactgagat acctacagcg 7500
tgagcattga gaaagcgcca cgcttcccga agggagaaaag gcggacaggt 7550
atccggtaag cggcagggtc ggaacaggag agcgcacgag ggagcttcca 7600
15 gggggaaacg cctggtatct ttatagtcct gtcgggttcc gccacctctg 7650
acttgagcgt cgatttttgt gatgctcgtc agggggggcg agcctatgga 7700
aaaacgccag caacgcggcc tttttacggt tcttggcctt ttgctggcct 7750
tttgtcaca tgttctttcc tgcgttatcc cctgattctg tggataaccg 7800
tattaccgcc tttgagttag ctgataccgc tcgccgcagc cgaacgaccg 7850
20 agcgcacgca gtcagtgagc gaggaagcgg aagagcgccc aatacgcaaa 7900
ccgcctctcc ccgcgcgttg gccgattcat taatccagct ggcacgacag 7950
gtttcccgac tggaaagcgg gcagtgagcg caacgcaatt aatgtgagtt 8000
acctcactca ttaggcaccc caggctttac actttatgct tccggctcgt 8050
atgttgtgtg gaattgtgag cggataacaa tttcacacag gaaacagcta 8100
25 tgaccatgat tacgaattaa 8120

<210> 69

<211> 800

<212> DNA

<213> Artificial Sequence

30 <220>

<221> Artificial Sequence
 <222> 1-800
 <223> recombinant immunoglobulin

<400> 69

```

5   aaaagggtat ctagagggtg aggtgatttt atgaaaaaga atatcgcat 50

    tcttcttgca tctatgttcg ttttttctat tgctacaaac gcgtacgctg 100
    aggttcagct agtgagctct ggcggtggcc tggcgcagcc agggggctca 150
    ctccgtttgt cctgtgcagc ttctggctac tccttctcga gtcactatat 200
    gcactgggtc cgtcaggccc cgggtaaggg cctggaatgg gttggatata 250
10  ttgatccttc caatggtgaa actacgtata atcaaaagtt caagggccgt 300
    ttcactttat ctgcgcacaa ctccaaaaac acagcatacc tgcagatgaa 350
    cagcctgcgt gctgaggaca ctgccgtcta ttactgtgca agaggggatt 400
    atcgctacaa tggtgactgg ttcttcgacg tctgggggtca aggaacctg 450
    gtcaccgtct cctcggcctc caccaagggc ccatcggtct tccccctggc 500
15  accctcctcc aagagcacct ctggggggcac agcgggcctg ggctgcctgg 550
    tcaaggacta ctccccgaa ccggtgacgg tgcgtggaa ctcaggcgcc 600
    ctgaccagcg gcgtgcacac ctcccggt gtccctacagt cctcaggact 650
    ctactcctc agcagcgtgg tgaccgtgcc ctccagcagc ttgggcaccc 700
    agacctacat ctgcaacgtg aatcacaagc ccagcaacac caaggtcgac 750
20  aagaaagttg agcccaaatc ttgtgacaaa actcacacat gcccgctga 800
  
```

<210> 70
 <211> 256
 <212> PRT
 <213> Artificial Sequence

25 <220>
 <221> Artificial Sequence
 <222> 1-256
 <223> recombinant immunoglobulin

<400> 70

```

30  Met Lys Lys Asn Ile Ala Phe Leu Leu Ala Ser Met Phe Val Phe
     1           5           10           15

    Ser Ile Ala Thr Asn Ala Tyr Ala Glu Val Gln Leu Val Gln Ser
           20           25           30

    Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys
35           35           40           45
  
```

	Ala Ala Ser Gly Tyr Ser Phe Ser Ser His Tyr Met His Trp Val	
	50	60
	Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Tyr Ile Asp	
	65	75
5	Pro Ser Asn Gly Glu Thr Thr Tyr Asn Gln Lys Phe Lys Gly Arg	
	80	90
	Phe Thr Leu Ser Arg Asp Asn Ser Lys Asn Thr Ala Tyr Leu Gln	
	95	105
10	Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala	
	110	120
	Arg Gly Asp Tyr Arg Tyr Asn Gly Asp Trp Phe Phe Asp Val Trp	
	125	135
	Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly	
	140	150
15	Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly	
	155	165
	Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu	
	170	180
20	Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val	
	185	195
	His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu	
	200	210
	Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr	
	215	225
25	Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp	
	230	240
	Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro	
	245	255
30	Pro	
	256	
	<210> 71	
	<211> 452	
	<212> PRT	
	<213> Artificial Sequence	
35	<220>	
	<221> Artificial Sequence	
	<222> 1-452	
	<223> recombinant immunoglobulin	

	<400>	71	Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln Pro Gly	1 5 10 15
			Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Ser Phe Ser	20 25 30
			Ser His Tyr Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu	35 40 45
			Glu Trp Val Gly Tyr Ile Asp Pro Ser Asn Gly Glu Thr Thr Tyr	50 55 60
10			Asn Gln Lys Phe Lys Gly Arg Phe Thr Leu Ser Arg Asp Asn Ser	65 70 75
			Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp	80 85 90
15			Thr Ala Val Tyr Tyr Cys Ala Arg Gly Asp Tyr Arg Tyr Asn Gly	95 100 105
			Asp Trp Phe Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val	110 115 120
			Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro	125 130 135
20			Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu	140 145 150
			Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser	155 160 165
25			Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln	170 175 180
			Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser	185 190 195
			Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys	200 205 210
30			Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys	215 220 225
			Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu	230 235 240
35			Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr	245 250 255
			Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp	260 265 270
			Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp	

		275		280		285
	Gly Val Glu Val His Asn Ala Lys Thr	Lys Pro Arg Glu Glu Gln				
	290	295			300	
5	Tyr Asn Ser Thr Tyr Arg Val Val Ser	Val Leu Thr Val Leu His				
	305	310			315	
	Gln Asp Trp Leu Asn Gly Lys Glu Tyr	Lys Cys Lys Val Ser Asn				
	320	325			330	
	Lys Ala Leu Pro Ala Pro Ile Glu Lys	Thr Ile Ser Lys Ala Lys				
	335	340			345	
10	Gly Gln Pro Arg Glu Pro Gln Val Tyr	Thr Leu Pro Pro Ser Arg				
	350	355			360	
	Glu Glu Met Thr Lys Asn Gln Val Ser	Leu Thr Cys Leu Val Lys				
	365	370			375	
15	Gly Phe Tyr Pro Ser Asp Ile Ala Val	Glu Trp Glu Ser Asn Gly				
	380	385			390	
	Gln Pro Glu Asn Asn Tyr Lys Thr Thr	Pro Pro Val Leu Asp Ser				
	395	400			405	
	Asp Gly Ser Phe Phe Leu Tyr Ser Lys	Leu Thr Val Asp Lys Ser				
	410	415			420	
20	Arg Trp Gln Gln Gly Asn Val Phe Ser	Cys Ser Val Met His Glu				
	425	430			435	
	Ala Leu His Asn His Tyr Thr Gln Lys	Ser Leu Ser Leu Ser Pro				
	440	445			450	
25	Gly Lys					
	452					
	<210> 72					
	<211> 219					
	<212> PRT					
	<213> Artificial Sequence					
30	<220>					
	<221> Artificial Sequence					
	<222> 1-219					
	<223> recombinant imunoglobulin					
	<400> 72					
35	Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val					
	1	5	10	15		
	Gly Asp Arg Val Thr Ile Thr Cys Arg Ser Ser Gln Ser Leu Val					
	20	25	30			
	His Gly Ile Gly Ala Thr Tyr Leu His Trp Tyr Gln Gln Lys Pro					

		35		40		45
		Gly Lys Ala Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe				
		50		55		60
		Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp				
5		65		70		75
		Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr				
		80		85		90
		Tyr Tyr Cys Ser Gln Ser Thr His Val Pro Leu Thr Phe Gly Gln				
		95		100		105
10		Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val				
		110		115		120
		Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala				
		125		130		135
15		Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys				
		140		145		150
		Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln				
		155		160		165
		Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu				
		170		175		180
20		Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys				
		185		190		195
		Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val				
		200		205		210
25		Thr Lys Ser Phe Asn Arg Gly Glu Cys				
		215		219		

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/01081

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 36-64 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/01081

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9523865	A	08-09-1995	CA 2181787 A	08-09-1995
			EP 0749488 A	27-12-1996
			JP 9509837 T	07-10-1997
			US 5707622 A	13-01-1998
			US 5702946 A	30-12-1997
			US 5686070 A	11-11-1997
			US 5677426 A	14-10-1997
EP 770628	A	02-05-1997	US 5874080 A	23-02-1999
			AU 701342 B	28-01-1999
			AU 2936395 A	16-02-1996
			CA 2194907 A	01-02-1996
			HU 77420 A	28-04-1998
			WO 9602576 A	01-02-1996
			JP 8217799 A	27-08-1996
			ZA 9505832 A	10-04-1996

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/US 99/01081

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/13 C07K19/00 A61K47/48 C07K16/24 C12N15/85 C12N5/10		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 23865 A (GENENTECH, INC. & INDIANA UNIVERSITY FOUNDATION) 8 September 1995 see examples see claims	1,26-29, 36-46, 61-64
A	N. KATRE: "The conjugation of proteins with polyethylene glycol and other polymers. Altering properties of proteins to enhance their therapeutic potential." ADVANCED DRUG DELIVERY REVIEWS, vol. 10, no. 1, 1993, pages 91-114, XP002084717 Amsterdam, The Netherlands see figure 3	1-25, 30-35, 47-60
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search	Date of mailing of the international search report	
3 June 1999	16/06/1999	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040. Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Nooij, F	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/01081

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	E. ENOAMOOQUAYE ET AL.: "Altered biodistribution of an antibody-enzyme conjugate modified with polyethylene-glycol." BRITISH JOURNAL OF CANCER, vol. 73, no. 11, June 1996, pages 1323-1327, XP002084718 London, GB see page 1324, left-hand column, line 33 - line 54	1-25, 30-35, 47-60
A	E. MAINOLFI ET AL. 'REDUCTION OF IMMUNOGENICITY OF A MURINE ANTI-ICAM-1 ANTIBODY THROUGH PEGYLATION CHEMISTRY.' "In: THE 9TH INTERNATIONAL CONGRESS OF IMMUNOLOGY (abstract book)," July 1995, SAN FRANCISCO, CA, USA XP002104756 see page 885, abstract 5247	1-25, 30-35, 47-60
A	C. CUNNINGHAM-RUNDLES ET AL.: "Biological activities of polyethylene-glycol immunoglobulin conjugates." JOURNAL OF IMMUNOLOGICAL METHODS, vol. 152, no. 2, 10 August 1992, pages 177-190, XP000471626 Amsterdam, The Netherlands see 'Material and Methods'	1-25, 30-35, 47-60
A	C. DELGADO ET AL.: "Enhanced tumour specificity of an anti-carcinoembryonic antigen Fab' fragment by poly(ethylene glycol) (PEG) modification." BRITISH JOURNAL OF CANCER, vol. 73, no. 2, January 1996, pages 175-182, XP002084719 London, GB see the whole document	1-25, 30-35, 47-60
A	EP 0 770 628 A (CHUGAI SEIYAKU KK ET AL.) 2 May 1997 see page 2, line 15 - page 3, line 25 see examples see claims	1, 26-29, 36-46, 61-64